Cite this: *Chem. Commun.*, 2012, **48**, 4821–4823

www.rsc.org/chemcomm

COMMUNICATION

Trifluoromethyl acting as stopper in [2]rotaxane†

Suvankar Dasgupta,^a Kuo-Wei Huang^b and Jishan Wu^{*a}

Received 11th February 2012, Accepted 23rd March 2012

DOI: 10.1039/c2cc31009k

A modified dumbbell obtained by replacing one of the phenyl groups of the dibenzylammonium with a strong electron-withdrawing trifluoromethyl group templated the synthesis of the smallest [2]rotaxane reported so far. The trifluoromethyl group not only enhances the templating effect of the dumbbell but also acts as the stopper to prevent dethreading of a [20]crown ether macrocycle.

Mechanically interlocked molecules (MIM)¹ in addition to being aesthetically appealing, also find potential applications in molecular nanotechnology. For example, rotaxane² based molecular switches³ have been visioned as potential candidates for molecular scale electronics,⁴ molecular actuators,⁵ molecular elevators,⁶ smart surfaces⁷ and controlled drug release.⁸ For constructing rotaxanes, the template-directed *clipping*⁹ methodology involves self-assembly of acyclic precursors around the recognition site of the dumbbell followed by reversible condensation^{9a,b,d,f,h} or metathesis^{9c,e,g,i} to generate the macrocycle encompassing the recognition site. The ring closing metathesis (RCM) of an acyclic diolefin polyether, in the presence of the “dumbbell” (ArCH₂)₂NH₂⁺ (Ar = aryl), is known to produce interlocked structures containing [24]crown ether.^{9b–d}

Very recently, using RCM we obtained a series of [2]rotaxanes on dibenzylammonium dumbbell (**1**, Fig. 1) with smaller macrocycles such as [20], [21] and [22]crown ethers, by systematically varying the alkyl chain length of the acyclic diolefin polyether.¹⁰ Inspired by the success with dibenzylammonium dumbbell and knowing that [21]crown ether was the smallest crown ether to be threaded by dialkylammonium ion (**2**, Fig. 1),¹¹ one can contemplate that clipping a [20]crown ether onto a dialkylammonium ion will give a [2]rotaxane with alkyl groups acting as stoppers! Therefore, salt **2**·PF₆¹² and acyclic diolefin polyether **5** were prepared.¹⁰ A mixture of two equivalents of **5** with one equivalent of **2**·PF₆ was stirred in CHCl₃–CH₃CN (3 : 1) for 24 h, followed by removing the solvent under reduced pressure

Fig. 1 Molecular structures of the dumbbells **1–4**, acyclic diolefin polyether **5**, and [20]crown ether **6**.

without heating. The residue, expected to be the complex **7**·PF₆ (soluble in CH₂Cl₂), was subjected to RCM with Grubbs' 2nd generation catalyst under reflux conditions in dry DCM for 60 h (Scheme 1).

Although the peak corresponding to the desired compound **10**·PF₆ was detected in nominal ESI-MS (Fig. S1, ESI[†]), purification by column chromatography failed to isolate **10**·PF₆. Dethreading during purification may explain this finding if **10**·PF₆ is a pseudorotaxane. However, no threading

Scheme 1 RCM of dumbbells **2**·PF₆, **3**·BAR₄ and **4**·BAR₄.

^a Department of Chemistry, National University of Singapore, 3 Science Drive 3, Singapore 117543. E-mail: chmwuj@nus.edu.sg; Fax: (+65) 6779 1691; Tel: (+65) 6516 2677

^b King Abdullah University of Science and Technology (KAUST), Division of Chemical and Life Sciences and Engineering and KAUST Catalysis Center, Thuwal 23955-6900, Saudi Arabia

† Electronic supplementary information (ESI) available: Synthetic procedures, nominal ESI-MS and HR ESI-MS spectra, 2D NOESY spectrum, stacked ¹H NMR spectra, decay curve, ¹H, ¹³C and ¹⁹F NMR spectra. See DOI: 10.1039/c2cc31009k

Fig. 2 Partial ^1H NMR spectra (500 MHz, CD_3CN), displaying the increase in acidity of methylene protons $\text{g} \rightarrow \text{e} \rightarrow \text{c}$ in dumbbells $2 \rightarrow 3 \rightarrow 4$. Benzylic protons a in **1** is for reference.

could be detected by ^1H NMR on refluxing a mixture of **2**• PF_6 with three equivalents¹³ of [20]crown ether **6** (for synthesis, see ESI†) for 7 days in CH_2Cl_2 followed by another 7 days in CHCl_3 . **10**• PF_6 was expected to be formed in extremely low yield (if at all), which would make it difficult to be isolated. This may be ascribed to the weaker templating effect of the dumbbell **2** than **1** due to the lower acidity of $-\text{CH}_2\text{NH}_2^+\text{CH}_2-$ protons (responsible for templating effect) in **2** (protons $\text{g} \sim 2.9$ ppm, Fig. 2(iv)) compared to **1** (protons $\text{a} \sim 4.2$ ppm, Fig. 2(i)). To increase the templating effect of the dumbbell, *N*-benzylalkylammonium dumbbells **3** and **4** having electron-withdrawing fluoromethyl ($-\text{CH}_2\text{F}$) and trifluoromethyl ($-\text{CF}_3$) stopper groups were prepared. However, the dumbbell with difluoromethyl ($-\text{CHF}_2$) stopper group could not be synthesized since the precursors $\text{F}_2\text{CHCH}_2\text{NH}_2$ or F_2CHCHO was not easily available.

The electronic impact of $-\text{CH}_2\text{F}$ and $-\text{CF}_3$ stopper groups can be observed from the downfield chemical shifts of the protons e (~ 3.4 ppm, Fig. 2(iii)) and c (~ 3.9 ppm, Fig. 2(ii)) in **3** and **4**, respectively. The dumbbells **3** and **4** were prepared with BAR_4 counter ion because their chloride precursors (**15**• Cl and **16**• Cl , see ESI†) could not be dissolved even in hot water and thus could not undergo a clean counter ion exchange reaction by mixing with aqueous NH_4PF_6 . Moreover, the BAR_4 counter ion is a weakly associating ion and extremely lipophilic, making **3** and **4** easily dissolve in DCM.¹⁴ The salts **3**• BAR_4 and **4**• BAR_4 were then mixed with two equivalents of **5** to afford **8**• BAR_4 and **9**• BAR_4 , respectively, which as for **7**• PF_6 above, were subjected to RCM with the Grubbs' 2nd generation catalyst under refluxing condition in dry DCM for 60 h.

The [2]rotaxane **11**• BAR_4 , corresponding to the salt **3**• BAR_4 , could only be detected in nominal ESI-MS (Fig. S2, ESI†) while [2]rotaxane **12**• BAR_4 obtained from salt **4**• BAR_4 was isolated as a white solid in reasonably good yield (54%, Scheme 1), reflecting the strong templating effect of **4**• BAR_4 .

Fig. 3 ^1H NMR spectra (500 MHz, CD_3CN) of (i) **4**• BAR_4 , and (ii) **12**• BAR_4 . Dotted lines indicate the downfield shift observed ($\text{b} \rightarrow \text{d}$) due to the effect of H-bonding interaction.

The ^1H NMR spectrum of [2]rotaxane **12**• BAR_4 (Fig. 3(ii)), displaying an $-\text{NH}_2^+$ hump (~ 8.6 ppm), olefinic proton peak (protons $\text{e} \sim 6.1$ ppm), and splitting of benzylic protons (protons $\text{c} \sim 4.3$ ppm), clearly indicated a [2]rotaxane structure encircled with a [20]crown ether.¹⁰ The large downfield shift of the methylene protons b in **4**• BAR_4 to d in **12**• BAR_4 can be explained by the participation of protons d in hydrogen bonding with the oxygen atoms of the [20]crown ether, substantiating an interlocked geometry for [2]rotaxane **12**• BAR_4 . The observed HR ESI-MS result for **12**• BAR_4 showed a peak at 480.2573 (Fig. S3, ESI†) with -1.2 ppm deviation from the calculated HRMS (480.2567) for $[\text{M} - \text{BAR}_4]^+$, showing that pure **12**• BAR_4 can be obtained by column chromatography. Repeated attempts to grow suitable single crystals failed, consequently 2D NOESY NMR (Fig. S4, ESI†) was performed in CDCl_3 to reveal the interlocked structure.

To ascertain the kinetic stability of **12**• BAR_4 , its ^1H NMR spectrum was recorded in $\text{DMSO}-d_6$, and a tiny peak (~ 5.75 ppm, Fig. S5(ii), ESI†) corresponding to the free crown ether was observed. We became suspicious of dethreading, however a threading experiment performed with the salt **4**• BAR_4 and three equivalents of [20]crown ether **6** failed despite refluxing the mixture in CH_2Cl_2 for 7 days followed by another 7 days in CHCl_3 . Moreover, in the ^1H NMR spectrum of **12**• BAR_4 recorded in $\text{DMSO}-d_6$ (Fig. S5, ESI†), the absence of peak corresponding to the free dumbbell suggested decay of [2]rotaxane **12**• BAR_4 was not by dethreading. Some electronic effects, operative in the component $-\text{NH}_2^+\text{CH}_2\text{CF}_3$ of **12**• BAR_4 , may be responsible for the slow decomposition in polar solvent (DMSO). A time-based concentration study of **12**• BAR_4 in $\text{DMSO}-d_6$ over a week suggested a first-order decay with half-life of 26.1 h (Fig. S6, ESI†). However, the compound **12**• BAR_4 did not show any decomposition after standing in CDCl_3 over two weeks and no dethreading was observed on heating up to 373 K in $\text{C}_2\text{D}_2\text{Cl}_4$ (Fig. S7, ESI†).

Fig. 4 Capped stick (a) and space-filling (b) representation for the calculated structure of [2]rotaxane **12-Bar₄**, showing short contact distances (in Å). (a) In capped stick structures, [C–H···O] and [N–H···O] distances are shown. Hydrogen atoms not involved in short contact distances are omitted for clarity. (b) In space-filling representation, oxygen atoms of crown ether and fluorine atoms of the dumbbell are represented in red and yellow, respectively. All the hydrogen atoms are omitted for clarity.

Theoretical calculations at the B3LYP/6-31+G(d,p) level of theory were conducted to understand the geometry of the [2]rotaxane **12** cation.¹⁵ This method was validated by comparing the optimized structure of an analogous [2]rotaxane with dibenzylammonium dumbbell interlocked with a [20]crown ether¹⁰ with its crystallographic structure, and it was confirmed that the computed geometry was in good agreement with the experimental observation. Similar calculations on cation **12** predict an interlocked [2]rotaxane structure involving multiple [C–H···O] and [N–H···O] interactions (Fig. 4a). The space-filling representation (Fig. 4b) gives an estimate of the cavity size of the [20]crown ether being smaller than the volume of the CF₃ group, which is consistent with the experimental results confirming the role of the CF₃ moiety as a stopper for the [20]crown ether.

In conclusion, the potential of alkyl or fluorinated alkyl groups acting as stoppers has been studied in detail, with the [20]crown ether as the encircling macrocycle. RCM has been utilized to clip the macrocycle onto the dumbbell. Indeed, we have obtained a very strongly interlocked [2]rotaxane on the PhCH₂NH₂⁺CH₂CF₃ dumbbell encircled with a [20]crown ether, highlighting the role of trifluoromethyl group in enhancing the template effect (needed for clipping) along with acting as a stopper. The [2]rotaxane **12-Bar₄**, though not stable in polar solvents, remains extremely stable in chlorinated solvents even at 373 K, underlying the steric resistance offered by the trifluoromethyl group is sufficient for acting as a stopper, which is also supported by theoretical calculations. The [2]rotaxane **12-Bar₄** represents the smallest [2]rotaxane reported so far, in terms of the molar mass and the total number of the atoms for the cationic part.¹⁶

This work was financially supported by the NUS AcRF Tier 1 grant (R-143-000-467-112) and KAUST. We thank Han Yanhui and Ong Wei Qiang for 2D NMR analysis.

Notes and references

- (a) D. B. Amabilino and J. F. Stoddart, *Chem. Rev.*, 1995, **95**, 2725–2828; (b) J. F. Stoddart, *Chem. Soc. Rev.*, 2009, **38**, 1802–1820; (c) J. E. Beves, B. A. Blight, C. J. Campbell, D. A. Leigh and R. T. McBurney, *Angew. Chem., Int. Ed.*, 2011, **50**, 9260–9327.
- (a) I. T. Harrison and S. Harrison, *J. Am. Chem. Soc.*, 1967, **89**, 5723–5724; (b) G. Schill, W. Beckmann and W. Vetter, *Angew. Chem., Int. Ed. Engl.*, 1973, **12**, 665–666; (c) C. A. Schalley, K. Beizai and F. Vögtle, *Acc. Chem. Res.*, 2001, **34**, 465–476; (d) A. Harada, A. Hashidzume, H. Yamaguchi and Y. Takashima, *Chem. Rev.*, 2009, **109**, 5974–6023; (e) H. Lahlali, K. Jobe, M. Watkinson and S. M. Goldup, *Angew. Chem., Int. Ed.*, 2011, **50**, 4151–4155.
- (a) B. L. Feringa, *Acc. Chem. Res.*, 2001, **34**, 504–513; (b) R. Klajn, J. F. Stoddart and B. A. Grzybowski, *Chem. Soc. Rev.*, 2010, **39**, 2203–2237; (c) H. Li, A. C. Fahrenbach, A. Coskun, Z. Zhu, G. Barin, Y. L. Zhao, Y. Y. Botros, J. P. Sauvage and J. F. Stoddart, *Angew. Chem., Int. Ed.*, 2011, **50**, 6782–6788.
- (a) C. P. Collier, G. Mattersteig, E. W. Wong, Y. Luo, K. Beverly, J. Sampaio, F. M. Raymo, J. F. Stoddart and J. R. Heath, *Science*, 2000, **289**, 1172–1175; (b) J. E. Green, J. W. Choi, A. Boukai, Y. Bunimovich, E. Johnston-Halperin, E. Delonno, Y. Luo, B. A. Sheriff, K. Xu, Y. S. Shin, H. R. Tseng, J. F. Stoddart and J. R. Heath, *Nature*, 2007, **445**, 414–417.
- (a) M. C. Jimenez, C. Dietrich-Buchecker and J. P. Sauvage, *Angew. Chem., Int. Ed.*, 2000, **39**, 3284–3287; (b) Y. Liu, A. H. Flood, P. A. Bonvallet, S. A. Vignon, B. H. Northrop, H. R. Tseng, J. O. Jeppesen, T. J. Huang, B. Brough, M. Baller, S. Magonov, S. D. Solares, W. A. Goddard, C. M. Ho and J. F. Stoddart, *J. Am. Chem. Soc.*, 2005, **127**, 9745–9759; (c) J. Wu, K. C. Leung, D. Benitez, J. Y. Han, S. J. Cantrill, L. Fang and J. F. Stoddart, *Angew. Chem., Int. Ed.*, 2008, **47**, 7470–7474.
- (a) J. D. Badjic, V. Balzani, A. Credi, S. Silvi and J. F. Stoddart, *Science*, 2004, **303**, 1845–1849; (b) J. D. Badjic, C. M. Ronconi, J. F. Stoddart, V. Balzani, S. Silvi and A. Credi, *J. Am. Chem. Soc.*, 2006, **128**, 1489–1499.
- J. B. Berna, D. A. Leigh, M. Lubomska, S. M. Mendoza, E. M. Perez, P. Rudolf, G. Teobaldi and F. Zerbetto, *Nat. Mater.*, 2005, **4**, 704–710.
- (a) T. D. Nguyen, H. R. Tseng, P. C. Celestre, A. H. Flood, Y. Liu, J. F. Stoddart and J. I. Zink, *Proc. Natl. Acad. Sci. U. S. A.*, 2005, **102**, 10029–10034; (b) S. Saha, K. C. F. Leung, T. D. Nguyen, J. F. Stoddart and J. I. Zink, *Adv. Funct. Mater.*, 2007, **17**, 685–693.
- (a) W. Clegg, C. Gimenez-Saiz, D. A. Leigh, A. Murphy, A. M. Z. Slawin and S. J. Teat, *J. Am. Chem. Soc.*, 1999, **121**, 4124–4129; (b) P. T. Glink, A. I. Oliva, J. F. Stoddart, A. J. P. White and D. J. Williams, *Angew. Chem., Int. Ed.*, 2001, **40**, 1870–1875; (c) A. F. Kilbinger, S. J. Cantrill, A. W. Waltman, M. W. Day and R. H. Grubbs, *Angew. Chem., Int. Ed.*, 2003, **42**, 3281–3285; (d) J. Wu, K. C. Leung and J. F. Stoddart, *Proc. Natl. Acad. Sci. U. S. A.*, 2007, **104**, 17266–17271; (e) P. G. Clark, E. N. Guidry, W. Y. Chan, W. E. Steinmetz and R. H. Grubbs, *J. Am. Chem. Soc.*, 2010, **132**, 3405–3412; (f) M. E. Belowich, C. Valente and J. F. Stoddart, *Angew. Chem., Int. Ed.*, 2010, **49**, 7208–7212; (g) A. M. Fuller, D. A. Leigh and P. J. Lusby, *J. Am. Chem. Soc.*, 2010, **132**, 4954–4959; (h) D. M. D'Souza, D. A. Leigh, L. Mottier, K. M. Mullen, F. Paolucci, S. J. Teat and S. Zhang, *J. Am. Chem. Soc.*, 2010, **132**, 9465–9470; (i) N. Momcilovic, P. G. Clark, A. J. Boydston and R. H. Grubbs, *J. Am. Chem. Soc.*, 2011, **133**, 19087–19089.
- S. Dasgupta and J. Wu, *Chem. Sci.*, 2012, **3**, 425–432.
- C. Zhang, S. Li, J. Zhang, K. Zhu, N. Li and F. Huang, *Org. Lett.*, 2007, **9**, 5553–5556.
- P. R. Ashton, P. J. Campbell, E. J. T. Chrystal, P. T. Glink, S. Menzer, D. Philp, N. Spencer, J. F. Stoddart, P. A. Tasker and D. J. Williams, *Angew. Chem., Int. Ed. Engl.*, 1995, **34**, 1865–1869.
- P. R. Ashton, I. Baxter, M. C. T. Fyfe, F. M. Raymo, N. Spencer, J. F. Stoddart, A. J. P. White and D. J. Williams, *J. Am. Chem. Soc.*, 1998, **120**, 2297–2307.
- B. A. Blight, A. Camara-Campos, S. Djurdjevic, M. Kaller, D. A. Leigh, F. M. McMillan, H. McNab and A. M. Slawin, *J. Am. Chem. Soc.*, 2009, **131**, 14116–14122.
- See ESI† for computational details.
- (a) C. C. Hsu, N. C. Chen, C. C. Lai, Y. H. Liu, S. M. Peng and S. H. Chiu, *Angew. Chem., Int. Ed.*, 2008, **47**, 7475–7478; (b) H. Lahlali, K. Jobe, M. Watkinson and S. M. Goldup, *Angew. Chem., Int. Ed.*, 2011, **50**, 4151–4155.