Accepted Manuscript

Arabidopsis Yak1 protein (AtYak1) is a dual specificity protein kinase

Dongjin Kim, Valentine Otang Ntui, Nianshu Zhang, Liming Xiong

PII: S0014-5793(15)00874-1

DOI: <http://dx.doi.org/10.1016/j.febslet.2015.09.025>

Reference: FEBS 37383

To appear in: *FEBS Letters*

Received Date: 11 July 2015

Revised Date: 8 September 2015

Accepted Date: 24 September 2015

Please cite this article as: Kim, D., Ntui, V.O., Zhang, N., Xiong, L., Arabidopsis Yak1 protein (AtYak1) is a dual specificity protein kinase, *FEBS Letters* (2015), doi: <http://dx.doi.org/10.1016/j.febslet.2015.09.025>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Arabidopsis Yak1 protein (AtYak1) is a dual specificity protein kinase

Dongjin Kim^{a,b,*}, Valentine Otang Ntui^{a,b}, Nianshu Zhang^c, Liming Xiong^{a,*}

^a Biological and Environmental Sciences and Engineering Division, King Abdullah University of Science and Technology (KAUST), 23955-6900, Thuwal, Kingdom of Saudi Arabia

^b Center for Desert Agriculture, Biological and Environmental Sciences and Engineering Division, King Abdullah University of Science and Technology (KAUST), 23955-6900, Thuwal, Kingdom of Saudi Arabia

^c Cambridge Systems Biology Centre, Department of Biochemistry, The Sanger Building, Tennis Court Road, Cambridge CB2 1GA, UK

* Corresponding authors at: Biological and Environmental Sciences and Engineering Division, King Abdullah University of Science and Technology (KAUST), 23955-6900, Thuwal, Kingdom of Saudi Arabia. Tel: +966-12-808-2668

E-mail addresses: liming.xiong@kaust.edu.sa (L. Xiong);
dongjin.kim@kaust.edu.sa, djkim73@gmail.com (D. Kim)

ABSTRACT

Yak1 is a member of dual-specificity Tyr phosphorylation-regulated kinases (DYRKs) that are evolutionarily conserved. The downstream targets of Yak1 and their functions are largely unknown. Here, a homologous protein AtYAK1 was identified in *Arabidopsis thaliana* and the phosphoprotein profiles of the wild type and an *atyak1* mutant were compared on two-dimensional gel following Pro-Q Diamond phosphoprotein gel staining. Annexin1, Annexin2 and RBD were phosphorylated at serine/ threonine residues by the AtYak1 kinase. Annexin1, Annexin2 and Annexin4 were also phosphorylated at tyrosine residues. Our study demonstrated that AtYak1 is a dual specificity protein kinase in Arabidopsis that may regulate the phosphorylation status of the annexin family proteins.

Keywords

Serine / Threonine protein kinase
Tyrosine protein kinase
Dual specificity protein kinase
Phosphorylation
Phosphopeptide

Abbreviations

PK, protein kinase; RBD, RNA binding family protein; DYRKs, dual specificity of Tyr phosphorylation regulated kinases; 2-D PAGE, two dimensional polyacrylamide gel electrophoresis; GST, glutathione S-transferase

1. Introduction

Phosphorylation is the most frequent type of protein post-translational modification, which is often associated with the molecular activation/de-activation of signal-transduction pathways [1]. In higher eukaryotes, an estimated one third of all proteins are regulated by phosphorylation by protein kinases (PKs). The genomes of plant species encode more than 1000 PKs [2]. However, the substrates of only a small fraction of these kinases are known [3].

Protein kinases are generally classified into serine (Ser)/threonine (Thr) kinases and tyrosine (Tyr) kinases, based on their substrate specificity. Several studies have shown that plants have dual protein kinases that are classified into a serine/threonine kinase family but catalyze phosphorylation on both serine/threonine and tyrosine residues [4-5], indicating that these plant kinases exhibit dual specificity. Several plants kinases have also been shown to autophosphorylate on Tyr, Ser and Thr *in vitro* [6]. In *Arabidopsis thaliana*, a few protein Tyr-specific phosphatases and 22 dual-specific phosphatases have been identified [1].

Yak1 is a member of Ser/Thr protein kinases family known as dual specificity of Tyr phosphorylation regulated kinases (DYRKs) in *Saccharomyces cerevisiae* [7]. In *S. cerevisiae*, Yak1 was originally identified as a negative regulator downstream of Ras/PKA signaling pathway. Yak1 is phosphorylated by PKA *in vitro* at several serine and threonine residues and regulated glucose starvation *in vivo*. However, the regulatory mechanisms of Yak1 activity and functions are largely unknown.

The present study was undertaken to identify proteins that are phosphorylated by Yak1 in *Arabidopsis thaliana*. Of the seven candidate proteins selected by 2-DE analysis, Annexin1, Annexin2, Annexin4 and RNA binding family protein (RBD) were confirmed with phosphorylation assay and MS analysis. Annexin1, Annexin2 and RBD were phosphorylated at serine/ threonine residues by AtYak1 kinase. Annexin1, Annexin2 and Annexin4 were additionally phosphorylated at tyrosine residues. Our data suggest that AtYak1 is dual specificity protein kinase in *Arabidopsis*.

2. Materials and methods

2.1. Plant materials and growth conditions

Arabidopsis (*Arabidopsis thaliana*) wild-type plants, T-DNA insertion mutant plants used in this work were of the Col-0 ecotype. The *atyak1-1* (SAIL_31_C10) and *atyak1-2* (SALK_0746) T-DNA insertion mutant alleles were obtained from the Arabidopsis Biological Resource Center (<http://www.arabidopsis.org>). Seeds were sterilized and germinated as described elsewhere [8]. Seven-day-old seedlings were harvested for total protein extraction.

2.2. Total protein extraction

Proteins were extracted from *atyak1-1* T-DNA mutant and wild-type plants by trichloroacetic acid (TCA)/acetone precipitation as described [9], with slight modifications. Briefly, 500 mg of tissue were ground to a fine powder in liquid nitrogen, suspended in 10% (w/v) TCA/acetone containing 0.06% (w/v) sodium sulfite and incubated overnight at -20 °C. The extract was centrifuged at 12,000 g for 20 min at 4 °C, washed three times in 80% (v/v) ice-cold acetone and air-dried for 10 min to remove residual acetone. Pellet was resuspended in rehydration buffer consisting of 7 M urea, 2 M thiourea, 4% (w/v) CHAPS, 40 mM Tris-HCl (pH 8.0), 0.001% (v/v) tributylphosphine (TBP), vortexed overnight and centrifuged at 12,000 g for 20 min. Protein was quantified by Bradford assay.

2.3. 2-D PAGE and image analysis

For 2-D PAGE (two dimensional polyacrylamide gel electrophoresis) 100 µg of protein was adjusted to 125 µl with rehydration buffer supplemented with 1% (v/v) ASB-14, 0.2% (w/v) DTT and 0.5% (v/v) ampholyte. Proteins were focused using the IPGphor II (GE Healthcare, Bio-Sciences AB, Uppsala, Sweden) at 20 °C, 10,000 V. Strips were equilibrated in 6 M urea, 50 mM Tris-HCl (pH 8.8), 2% (w/v) SDS, 30% (v/v) glycerol containing 1% (w/v) DTT for 20 min followed by incubation in the same buffer containing 2.5% (w/v) iodoacetamide for another 20 min and then run in 12% SDS-PAGE. Phosphoproteins were stained using Pro-Q diamond phosphoprotein gel staining kit (Invitrogen Cat. No# MPP33000) following the manufacturer's instructions. Proteins were visualized using excitation at 532 nm with 560 nm longpass emission filter by Typhon 9410 scanner (GE Healthcare). Image analysis was performed with Delta2D Software Version 4.3 (DECODON,

www.decodon.com). Only proteins with significant and reproducible changes were considered to be differentially expressed.

2.4. *In-gel trypsin digestion and MALDI-TOF analysis*

Enzymatic digestion was performed as previously described [10]. Briefly, digestion was performed with 10 ng/mL of trypsin and 100 mM ammonium bicarbonate and incubated overnight at 37 °C. Following enzymatic digestion, the resultant peptides were extracted with 20 mM ammonium bicarbonate and twice with 5 % formic acid in 50 % acetonitrile and dried in SpeedVac. The peptides for LC-MS were suspended in 5 % acetonitrile and 0.1 % formic acid, and those for MALDI-TOF were suspended in 5 % acetonitrile and 0.1 % trifluoroacetic acid. Peptide masses were measured on a MALDI-TOF (maXis UHR-TOF, Bruker) [11]. Peptide mass fingerprint data were matched to the TAIR10 Arabidopsis protein sequence database by using MASCOT and scaffold software.

2.5. *Expression and purification of recombinant protein*

GST fusion protein, pGEX6p-AtYak1, pGEX6p-AnnAt1, pGEX6p-AnnAt2, pGEX6p-AnnAt4, pGEX6p-CAB3, pGEX6p-RCA, and pGEX6p-RBD were generated using primers sets listed in Table 1. Protein expression and purification was performed as described [12] with modifications. Briefly, 50 ml of bacterium cell culture were harvested by centrifugation, washed once with 20 ml of ice-cold STE [10 mM Tris, pH 8.0, 150 mM NaCl, 1 mM EDTA] and resuspended by repeated pipetting in 5 ml of STE containing 100 µg/ml of lysozyme. After incubation on ice for 20 min, dithiothreitol (DTT) was added to a final concentration of 5 mM. Protease inhibitors were also added. Bacteria were lysed by adding 1.5% N-laurylsarcosine (Sarkosyl). The cells were then vortexed for 5 s and sonicated on ice for 1 min. Thereafter, the lysate was centrifuged for 10 min at full speed at 4 °C. The supernatant was collected and mixed with 2% Triton X-100. Then, 200 µl of washed glutathione agarose bead suspension (50% v/v in PBS) was added and the lysate was incubated at 4 °C for 15 min. The beads were washed 6 times with ice-cold PBS [8.4 mM Na₂HPO₄, 1.9 mM NaH₂PO₄, pH 7.4, 150 mM NaCl] by repeated centrifugation and were resuspended in elution buffer [75 mM Hepes, pH 7.4, 150 mM NaCl, 10 mM

Reduced Glutathione, 5 mM DTT]. For storage, glycerol was added to a final concentration of 10% (v/v) and the protein solution was kept at -80 °C.

2.6. *In Vitro* Phosphorylation assays

AtYak1 kinase activity assays were performed with 20 ng of GST-AtYak1 purified protein and 1 µg of target protein in a 50 µl reaction volume. Assay buffer contained 50 mM HEPES, pH 7.5, 10 mM MgCl₂, 2 mM DTT, 2 mM EGTA, and 2.5 mM CaCl₂ with 60 mM ATP as a phosphate donor. Kinase assay components were assembled in each tube on ice. After being transferred to 30 °C, ATP was added and the mixture was incubated for 30 min. Reactions were stopped by adding 10 µl of SDS sample loading buffer. Aliquots of 30 µl of each reaction mixture were resolved in 12% SDS-PAGE gel and stained with Coomassie Blue.

2.7. *Mass spectrometry*

The peptides were fractionated by C₁₈ RP chromatography and analyzed online by mass spectrometry (LC-MS/MS). Mass spectrometry analysis was carried out using LTQ Orbitrap Velos (Thermo-Scientific) mass spectrometer connected to a nano-LC system (Proxeon). Peptides (5 µl injection) were loaded onto a C₁₈ trap column (Michrom), washed, followed by elution at 400 µl/min flow rate via on line analytical column (Michrom 3 µM, 200 Å, Magic C18AQ, 0.1 x 150 mm) connected to a nano-spray source (Proxeon). The peptide elution from C₁₈ column was achieved by a 25 min linear gradient of 5–35% (v/v) acetonitrile in 0.1% (v/v) formic acid, and a 5 min linear gradient of 35–80% (v/v) acetonitrile in 0.1% (v/v) formic acid, followed by 10 min isocratic flow of 80% (v/v) acetonitrile in 0.1% (v/v) formic acid. The eluted peptides were analyzed online by electrospray ionization tandem mass spectrometry, Xcalibur (v2.1) software (Thermo-Scientific) was used to collect mass spectrometry data, and mass range for the MS survey scans was set at m/z 350–1600. The m/z value of ions during first full scan was determined at high accuracy using Orbitrap mass analyzer. Each MS scan was followed by 10 MS/MS scans, and the data was collected using LTQ-linear ion trap. MS3 scan was selectively triggered upon neutral loss ion (32.7, 49, 80, and 98) detection. We used a dynamic exclusion method where a specific ion was sequenced twice at maximum and excluded from the list for 45 s.

The RAW mass spectrometry data files were converted to mgf format using Proteome Discoverer 1.2 (Thermo-Scientific), and the MS/MS data sets were compared with the TAIR10 Arabidopsis protein sequence database using Mascot search engine (v2.3, Matrix Science).

3. Results

3.1. *AtYak1* T-DNA insertion line genotyping

We obtained two alleles of *atyak1* T-DNA insertion mutants from the Arabidopsis Biological Resource Center. The T-DNA was inserted in the first exon in the *atyak1-1*(SAIL_31_C10) mutant and in the ninth intron in the *atyak1-2* (SALK_074646) mutant (Fig. 1A). RT-PCR screening of these homozygous lines revealed complete knockout of *AtYak1* in the *atyak1-1* mutant yet the *AtYAK1* transcripts were still observed in *atyak1-2* (Fig. 1B). We checked the phenotype of *atyak1-1*, *atyak1-2* and wild-type plants under normal growth conditions. *atyak1-1* plants had smaller and fewer leaves and displayed reduced biomass than the wild-type plants, whereas there was no difference in growth and biomass between *atyak1-2* plants and wild-type plants (unpublished data). The results show that *atyak1-1* was a null allele and thus was used in the current study.

3.2. Classification of identified phosphorylated proteins

Total proteins isolated from wild-type and *atyak1-1* mutant seedlings were resolved by 2D gel electrophoresis and phosphoproteins were identified by staining using Pro-Q diamond phosphoprotein gel staining kit. In either the mutant or wild-type samples, about 150 protein spots were found to distribute evenly between pH 4 and 7 and the molecular masses of these proteins range from 10 to 80 kD. To classify these phosphorylated proteins, we used Classification Superviewer. Proteins were assigned to 13 classes based on their domain composition or functional description in the literature (Fig. 2). Phosphorylated proteins involved in metabolic and cellular processes were the most abundant, whereas proteins involved in developmental processes, transport and other signaling processes were the least abundant. Between the *atyak1-1* mutant and the wild-type plants, three categories of proteins were identified: Proteins that show the same phosphopeptide abundance in both the wild-type and the *atyak1-1* mutant, proteins that show decreased phosphopeptide

abundance in the *atyak1-1* mutant plants compared with the wild-type, and proteins that show increased phosphopeptide abundance in the *atyak1-1* mutant compared with the wild-type. For example, phosphopeptide proteins involved in metabolic and cellular processes showed the same abundance in both the *atyak1-1* mutant and the wild-type plants. Phosphorylated proteins involved in electron transport or energy pathways showed decreased abundance in the *atyak1-1* mutant compared with wild-type, whereas those involved in biotic and abiotic stress response showed increased abundance in the *atyak1-1* mutant compared to wild-type plants (Fig. 2). Full lists of the identified proteins are provided in Table S2.

3.3. Identification of phosphorylated proteins

Among the phosphoproteins separated on the 2-D gel, we selected 50 spots based on differences in their intensity between the wild-type and the mutant and revealed the identities of these proteins with MALDI-TOF MS (Fig. 3A, Table 1). To analyze the changes in spot intensity between the wild-type and *atyak1-1* mutant, samples were quantified (section 2.3). Six spots (numbers 13, 14, 16, 20, 21, 22 and 26) representing protein p13, p14, p16, p20, p21 p22 and p26, respectively, were found to be differentially expressed or phosphorylated between the wild-type and the mutant (Fig. 3B). As compared to the wild-type, the level of expression or phosphorylation of p13 increased, whereas that of p14, p16, p20, p21 p22 and p26 seemed to decrease or not detected in the *atyak1-1* mutant.

Among these proteins, p26 (spot number 26) was identified as Annexin1 (AnnAt1), which has been implicated in responses to heat, cold, salt, osmotic stress and abscisic acid (ABA) in plants [12]. The intensity of the p26 spot was higher in the wild-type than in the *atyak1-1* mutant (Fig. 3A). Protein p26 migrated with a molecular mass of 42.0 kD (Table 1), which is larger than the standard molecular size of AnnAt1 (36.2 kD). This difference may be because of the different developmental stages or growth conditions of the plants under investigation [13]. The apparent pI value of p26 is 5.2, similar to the standard value of 5.0. Phosphoproteins p21 and p22 (spot numbers 21 and 22) were identified as chlorophyll A/B binding protein 3 (CAB3, AB180) that functions in photosynthesis. p21 and p22 spots were only observed in the wild-type but not in the *atyak1-1* mutant (Table 1), suggesting that their expression and/or phosphorylation is dependent on AtYak1. These proteins and their annotations are listed in Table 1.

3.4. *In vitro* phosphorylation assay

Among the phosphorylated proteins identified, we selected AnnAt1, AnnAt2, AnnAt4, CAB3, RBD and RCA as target proteins for further characterization. We purified six GST fused target proteins from the *Escherichia coli* BL21. For *in vitro* phosphorylation, target and substrate protein combination and enzyme reactions are shown in Supplementary Figure 1. After enzyme reaction, each protein band was isolated for phosphoprotein detection by LC-MS/MS (LTQ Orbitrap). Our *in vitro* phosphorylation assay analysis shows that all 3 members of annexin family, AnnAt1, AnnAt2 and AnnAt4, were phosphorylated by AtYak1 kinase. And AtYak1 kinase was phosphorylated by itself. On the other hand, CAB3 and RCA were not phosphorylated by AtYak1 kinase (Supplementary Figure 1).

3.5. MS identification of phosphorylation sites

To identify the sites where the AtYak1 kinase and target proteins, annexins and RBD, were phosphorylated, we conducted LC-MS/MS analysis. Annotated LC-MS/MS spectra showing peptide fragmentation and predicted phosphorylation sites are presented in Figures 4 and 5, and in Table 2. Fragmentation patterns of the doubly charged ions 448.76 m/z from the AtYak1 and 805.86 m/z from the RBD peptide digests (Fig. 4 A and B) resulted in the match of the sequence of peptide fragment 895.51 Da of AtYak1 and 1609.72 Da of RBD. The phosphorylation site of AtYak1 was deduced to be located at the Ser-202 by autophosphorylation (Fig. 4A, Table 2). Interestingly, this autophosphorylation site differs from yeast where autophosphorylation was observed at tyrosine residues in the activation loop among the DYRKs [14]. The phosphorylation sites of RBD were located at Ser-6 and Ser-12 (Fig. 4B, Table 3). The identification of the phosphorylation sites in the AnnAt1 and AnnAt2 was performed similarly. Fragmentation pattern of the doubly charged ions 484.24 m/z from AnnAt1 and 382.15 m/z from AnnAt2 matched the sequence of peptide fragment 1448.77 Da of AnnAt1 (Fig. 4C) and 1146.40 Da of AnnAt2 (Fig. 4D). The amino acid sequences of the peptide fragment 1448.77 Da showed that the phosphorylation site was located at Ser-289 (Fig. 4C, Table 2). The phosphorylation sites of AnnAt2 were identified at Ser- 95 and Thr-100 (Fig 4D, Table 2).

The Annexin family members (AnnAt1, AnnAt2 and AnnAt4) also phosphorylated at tyrosine by AtYak1 kinase. Fragmentation patterns of the doubly

charged ions 537.25 m/z from the AnnAt1, 554.27 m/z from the AnnAt2 and 673.85 m/z from AnnAt4 matched the sequence of the peptide fragments 1072.45 Da of AnnAt1 (Fig. 5A, Table 2), 1106.53 of AnnAt2 (Fig. 5B) and 1345.69 of AnnAt4 (Fig. 5C). The phosphorylation site of AnnAt1, AnnAt2 and AnnAt4 was located at Tyr-284, Tyr-126 and Tyr-211, respectively (Fig. 5A-C, Table 2).

4. Discussion

The aim of this study was to identify components of the AtYAK1 signaling pathway in Arabidopsis by examining the phosphorylation status of the proteomes of wild-type and the *atyak1-1* mutant plants. In this study, we used two-dimensional gel electrophoresis and LC-MS/MS to identify a number of candidates that showed altered phosphoprotein/peptide abundance in the *atyak1-1* mutant. Several protein spots were identified that showed altered phosphopeptide abundance in the *atyak1-1* mutant. Through proteomic analysis, we identified and selected Annexins for further characterization. Annexins are a family of Ca^{2+} -dependent membrane binding proteins that exist in almost all species from fungi to humans [15]. These proteins have been shown to play important roles in various cellular processes such as membrane trafficking and organization, regulation of ion channel activity, inflammatory response, mitotic signaling, and phospholipid metabolism [16]. Our results demonstrate that the abundance or phosphorylation of Annexin family proteins was lower in the *atyak1-1* mutant (Fig. 3B). *In vitro* experiments suggest that AtYak1 may directly phosphorylate annexin proteins in Arabidopsis.

Although in most plant species no more than two annexin genes have been reported, in Arabidopsis, seven distinct Annexin genes have been identified, which are AnnAt1-AnnAt7 [17], with AnnAt4 being the most divergent. AnnAt1 and AnnAt4 have been suggested to play important roles in osmotic and ABA signaling [13]. Here, we identified three Annexins, AnnAt1, AnnAt2 and AnnAt4, which were phosphorylated at serine / threonine and tyrosine residues by AtYak1 kinase.

Yak1 belongs to a dual-specificity tyrosine-regulated protein kinase (DYRK) family that autophosphorylate an essential tyrosine in their activation loop, but phosphorylate their substrate at several serine and threonine residues [14] and regulate glucose starvation *in vivo*. Studies have shown that Yak1 acts downstream of the PKA pathway [18] and that it controls transcriptional factors involved in stress response

[19] and ribosomal biogenesis [18]. In previous studies, Yak1 was shown to phosphorylate casein [20] and autophosphorylate at tyrosine residues when expressed in yeast [14]. In this study, we showed that AtYak1 protein kinase phosphorylates its substrates on Ser/Thr residues and undergoes Ser-202 autophosphorylation, indicating that Yak1 is a dual specificity protein kinase. In yeast, autophosphorylation was observed at tyrosine residues in the activation loop among the DYRKs [14]. In our experiment, autophosphorylation was detected at the Ser-202 residue. This difference may be due to the different studied organisms or incomplete peptide sequencing of AtYak1.

In conclusion, this study demonstrates that AtYak1 is a dual specificity protein kinase in Arabidopsis, which phosphorylates substrate proteins on Ser/Thr and Tyr residues. We are currently undertaking studies to find out the role of AtYak1 in stress response in Arabidopsis.

Acknowledgements

This work was partly supported by the AEA program between KAUST and the University of Cambridge. Special thanks are due to Dr. Ludivine Thomas in Biosciences core laboratory of KAUST for advice on the proteomic work.

References

- [1] Wan, L., Ross, A.R.S., Yang, J., Hegedus D.D. and Kermode A.R. (2007) Phosphorylation of the 12 S globulin cruciferin in wild-type and *abil-1* mutant *Arabidopsis thaliana* (thale cress) seeds. *Biochem. J.* 404, 247-256.
- [2] The Arabidopsis Genome Initiative (2000) Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature* 408, 796-815.
- [3] van Bentem, S.F., Anrather, D., Dohnal, I., Roitinger E., Csaszar, E., Joore, J., Buijnink, J., Carreri, A., Forzani, C., Lorkovi, Z.J., Barta, A., Lecourieux, D., Verhounig, A., Jonak, C. and Hirt H. (2008) Site-specific phosphorylation profiling of Arabidopsis proteins by mass spectrometry and peptide chip analysis. *J. Proteome Res.* 7, 2458-2470.

- [4] Douville E., Duncan, P., Abraham, N. and Bell, J. (1994) Dual specificity kinases- a new family of signal transducers. *Cancer Metastasis Rev.* 13, 1-7.
- [5] Lindberg R.A., Quin, A.M. and Hunter T. (1992) Dual-specificity protein kinases: will any hydroxyl do? *Trends Biochem. Sci.* 17, 114-119.
- [6] Rudrabhatla P., Reddy M.M. and Rajasekharan R. (2006) Genome-wide analysis and experimentation of plant serine/threonine/tyrosine-specific protein kinases. *Plant Mol. Biol.* 60, 293-319.
- [7] Yoshida K. (2008) Role for DYRK family kinases on regulation of apoptosis. *Biochem. Pharmacol.* 76, 1389-1394.
- [8] Joana S., Sara F., Rui M. T., Teresa L. and Herlander A. (2014) Phenotype analysis of the Arabidopsis heat stress response during germination and early seedling development. *Plant Methods*, 10, 7-18.
- [9] Wang, W., Vignani, R., Scali, M. and Cresti, M. (2006) A universal and rapid protocol for protein extraction from recalcitrant plant tissues for proteomic analysis. *Electrophoresis* 27, 2782-2786.
- [10] Shevchenko, A., Wilm, M., Vorm, O. and Mann, M. (1996) Mass spectrometric sequencing of proteins silver-stained polyacrylamide gels. *Anal. Chem.* 1, 850-858.
- [11] Landry, F., Lombardo C.R. and Smith, J.W. (2000) A method for application of samples to matrix-assisted laser detection. *Anal. Biochem.* 279, 1-8.
- [12] Frangioni J.V. and Neel B.G. (1993) Solubilization and purification of enzymatically active glutathione S-transferase (pGEX) fusion protein. *Anal. Biochem.* 210, 179-187.
- [13] Lee, S., Lee E.J., Yang, E.J., Lee, J.E, Park, A.R., Song, W.H. and Park, O.K. (2004) Proteomic identification of Annexins, calcium-dependent membrane binding proteins that mediate osmotic stress and abscisic acid signal transduction in Arabidopsis. *Plant Cell* 16, 1378-1391.
- [14] Becker, W. and Joost H.G. (1999) Structural and functional characteristics of DYRK, a novel subfamily of protein kinases with dual specificity. *Prog. Nucleic Acid Res. Mol. Biol.* 62, 1-17.
- [15] Gerke, V. and Moses, S.E. (2002) Annexins: From structure to function. *Physiol. Rev.* 82, 331-371.
- [16] Knight, H. and Knight, M.R. (2001) Abiotic stress signaling pathways: Specificity and cross-talk. *Trends Plant Sci.* 6, 262-278.

- [17] Clark G. B. and Roux S. J. (1995) Annexins of plant cells. *Plant Physiol.* 109, 1133-1139.
- [18] Martin, D.E., Soulard, A. and Hall, M.N. (2004) TOR regulates ribosomal protein gene expression via PKA and the Forkhead transcriptional factor FHL1. *Cell* 119, 969-979.
- [19] Lee, P., Cho, B., Joo, H. and Hahn J. (2008) Yeast Yak1 kinase, a bridge between PKA and stress-responsive transcriptional factors, Hsf1 and Msn2/Msn4. *Mol. Microbiol.* 70, 882-895.
- [20] Garrett, S., Menold, M.M. and Broach, J.R. (1991) The *Saccharomyces cerevisiae* YAK1 gene encodes a protein kinase that is induced by arrest early in the cell cycle. *Mol. Cell Biol.* 11, 4045-4052.

Figure legends

Figure 1. Genotyping and gene expression analyses of the AtYak1 T-DNA insertion mutants. (A) Scheme of *AtYak1* gene. The triangle indicates the position of the T-DNA insertion. Genomic AtYak1 DNA sequence is represented by exons (solid black bars), introns (thin black lines) and UTRs (white bars). Homozygous lines were genotyped with LP, RP and BP primers. (B) RT-PCR analysis of *AtYak1* expression in wild-type and homozygous *atyak1-1* and *atyak1-2* mutant plants. Red arrows (in Fig. 1A) indicate positions of primers used in RT-PCR. *ACT2* was used as a control.

Figure 2. Functional classification of the identified phosphorylated proteins. (A) The pie chart shows the identified phosphoprotein of wild-type Col-0. One hundred and fifteen peptides were identified. (B) Identified phosphoprotein of *atyak1-1* mutant. One hundred and fifty peptides were identified. Identified proteins were grouped according to functional classification in Classification SuperViewer (http://bar.utoronto.ca/ntools/cgi-bin/ntools_classification_superviewer.cgi).

Figure 3. Comparison of phosphorylated proteins between wild-type and *atyak1-1* mutant plants. The gels were stained with Pro-Q Diamond phosphorylation stain kit. (A) Total phosphorylated proteins in the range of pH 4 to 7. Total phosphoproteins in the wild-type (left) and *atyak1-1* mutant (right). Protein spots identified by MALDI-

TOF MS are numbered and listed in Table 1. (B) Selected phosphoproteins with changed expression levels. Circles indicate absence of phosphoprotein in the *atyak1-1* mutant.

Figure 4. Serine / threonine phosphorylation site identification of AtYak1, RBD, AnnAt1 and AnnAt2 by LC-MS/MS. Fragmentation patterns of the doubly charged ions 448.76 m/z from the AtYak1. The amino acid sequence of the peptide fragments 895.51 Da of AtYak1 is shown with the phosphorylation site of Ser-202 by autophosphorylation (A) and 805.86 m/z from the RBD peptide digests. The amino acid sequence of the peptide fragments 1609.72 Da of RBD is shown with the phosphorylation sites of Ser-6 and Ser-12 by AtYak1 kinase (B). Fragmentation pattern of the doubly charged ions 484.24 m/z from AnnAt1 (C). Fragmentation pattern of the doubly charged ions 383.15 m/z from AnnAt2 (D). The amino acid sequences of the peptide fragments 1448.77 Da and 1146.40 Da are shown with the phosphorylation site of Ser-289 from AnnAt1 and Ser- 95 and Thr-100 from AnnAt2 by AtYak1 kinase.

Figure 5. Tyrosine Phosphorylation site identification of AnnAt1 and AnnAt2 by LC-MS/MS. Fragmentation patterns of the doubly charged ions 537.25 m/z from the AnnAt1. The amino acid sequence of the peptide fragments 1072.45 Da of AnnAt1 is shown with the phosphorylation site of Tyr-284 (A) and 554.27 m/z from the AnnAt2 peptide digests. The amino acid sequence of the peptide fragments 1106.53 Da of AnnAt2 is shown with the phosphorylation site of Tyr-126 (B). Fragmentation pattern of the doubly charged ions 673.85 m/z from AnnAt4. The amino acid sequences of the peptide fragments 1345.69 Da are shown with the phosphorylation site of Tyr-211 from AnnAt4. Annexin family(AnnAt1, AnnAt2 and AnnAt4) shown tyrosine phosphorylation by AtYak1 kinase.

Supplementary Figure 1. Phosphorylation assay of candidate substrate proteins by AtYak1 kinase *in vitro*. Coomassie blue stained SDS-PAGE gel indicates AtYak1 kinase and substrate loading. See Material and Methods for details.

B

Table 1. Functional annotation of classified phosphoproteins

Spot No. ^a	Protein name ^b	Accession No.	Match MW(kD)/pI ^c	Apparent MW(kD)/pI ^d	Identification probability ^e	Percent Covered ^f	Fold change	
5	GTP binding Elongation factor Tu family protein	AT1G07920.1	49.5 / 9.6	41.5 / 4.3	100.00%	23.50%	-1.51	
6	Tubulin beta-7 chain (TUB7)	AT2G29550.1	50.7 / 4.5	32.0 / 4.3	100.00%	18.90%	-0.84	Salt stress response
8	Glyceraldehyde-3-phosphate dehydrogenase of plastid 2 (GAPCP-2)	AT1G16300.1	44.8 / 8.9	25.0 / 4.5	100.00%	20.50%	-0.38	Root development
13	Chloroplast RNA-binding protein 29 (CP29)	AT3G53460.1	38.4 / 5.2	30.0 / 4.8	100.00%	19.90%	3.58	Cold stress response
14	Pyrophosphorylase 6 (AtPPa6)	AT5G09650.1	33.8 / 5.8	30.0 / 4.9	100.00%	17.30%	-5.42	Cold stress, Salt stress response
15	RNA-binding (RRM/RBD/RNP motifs) family protein	AT2G37220.1	30.7 / 4.7	24.5 / 4.6	99.80%	11.40%	-0.50	Light reaction, ABA & cold response
16	RNA-binding (RRM/RBD/RNP motifs) family protein	AT2G37220.1	30.7 / 4.7	24.5 / 4.6	100.00%	17.60%	-2.16	Light reaction, ABA & cold response
20	Ribulose-bisphosphate carboxylases (RBCL)	ATCG00490.1	52.9 / 6.2	60.0 / 5.0	100.00%	34.40%	-9.15	ABA response
20	Trigger factor type chaperone family protein	AT5G55220.1	61.7 / 5.0	61.7 / 5.0	100.00%	25.60%	9.15	
21	Chlorophyll A/B binding protein 3 (CAB3, AB180)	AT1G29910.1	28.2 / 5.0	40.0 / 4.8	99.80%	3.75%	-7.99	Photosynthesis
22	Chlorophyll A/B binding protein 3 (CAB3, AB180)	AT1G29910.1	28.2 / 5.0	40.0 / 4.8	100.00%	8.99%	-55.80	
23	FtsH extracellular protease family (VAR1, FTSH5)	AT5G42270.1	75.2 / 5.1	72.0 / 5.2	100.00%	11.80%	-2.01	
25	Phosphoribulokinase (PRK)	AT1G32060.1	44.4 / 5.9	44.0 / 5.2	100.00%	10.10%	-3.09	
26	Annexin 1 (ANNAT1)	AT1G35720.1	36.2 / 5.0	42.0 / 5.2	100.00%	16.10%	-1.32	Cold, Heat, osmotic, salt and ABA response
27	Ribosomal protein L10 family protein	AT3G11250.1	34.4 / 4.7	40.0 / 5.1	100.00%	8.36%	-2.78	Salt stress response
28	Thioredoxin superfamily protein	AT3G52960.1	24.7 / 9.5	15.0 / 5.0	100.00%	13.20%	2.39	
30	Ureidoglycolate amidohydrolase (ATAAH-2, UAH)	AT5G43600.1	51.5 / 5.3	60.0 / 5.5	99.80%	3.99%	2.31	
31	Insulinase (Peptidase family M16) protein	AT1G51980.1	54.4 / 6.2	55.0 / 5.5	100.00%	14.50%	-0.96	Salt stress response
34	Rubisco activase (RCA)	AT2G39730.1	51.9 / 5.9	54.0 / 5.1	100.00%	7.17%	-0.52	Cold, light & JA response

^a Number of spot^b Identified protein name^c Predicted molecular weight (MW) and pI of match peptide sequence^d Observed molecular weight (MW) and pI of spot from the gel^e Percentage of protein identification probability^f Percentage of predicted protein sequence covered by matched peptides

Table 2. Phosphorylation sites identified within AtYak1 and target protein/*In Vitro* phosphorylation assay by LC-MS/MS

Protein	Peptide and phosphorylation site(s)	Locus	Residues	Describe*
AtYak1	GL p SLSIVK	At5g35980.1	220-227	Protein kinase, Autophosphorylation
AnnAt1	YIGEE p YQR	At1g35720.1	279-286	Annexin gene family, Tyr
AnnAt1	RN p SIPLEKAITK	At1g35720.1	287-298	Annexin gene family, Ser/Thr
AnnAt1*	RWTS p SNQVLMEVAC p TRTSTQLLHAR	At1g35720.1	98-122	Annexin gene family, Ser/Thr
AnnAt1*	LLVSLVT pSp YRYEGDEVNMTLAK	At1g35720.1	148-169	Annexin gene family, Ser/Thr, Tyr
AnnAt2	QAYQAR p YK	At5g65020.1	123-130	Annexin gene family, Tyr
AnnAt2	Ep ST KMF p TK	At5g65020.1	94-101	Annexin gene family, Ser/Thr
AnnAt4	SKLHLQHL p YK	At2g38750.1	203-212	Annexin gene family, Tyr
AnnAt4*	ADH p SDEMNEIK	At2g38750.1	274-284	Annexin gene family, Ser/Thr
RBD	AAS p SSLALS p SFNPCK	At2g37220.1	2 – 17	Chloroplast RNA binding protein, Ser/Thr

pS, phosphoserine; pT, phosphothreonine; pY, phosphotyrosine

* Ions Score below 10

Highlights

- One hundred fifty phosphoprotein spots identified by 2-D PAGE
- Four proteins confirmed to be phosphorylated by AtYak1 kinase
- AnnAt1, AnnAt2, AnnAt4 and RBD phosphorylated Ser/Thr residues by AtYak1 kinase
- AnnAt1, AnnAt2 and AnnAt4 phosphorylated Tyr residues by AtYak1 kinase