ORGANIC CHEMISTRY

FRONTIERS

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: S. Nolan, L. Cavallo, A. Poater, S. Vummaleti and G. Talarico, *Org. Chem. Front.*, 2015, DOI: 10.1039/C5QO00281H.

This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Journal Name

ARTICLE

How easy is CO₂ fixation by M-C bond containing complexes (M = Cu, Ni, Co, Rh, Ir)?

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Sai V. C. Vummaleti,^a Giovanni Talarico,^b Steven P. Nolan^{*,c} Luigi Cavallo^{*,a} and A. Poater^{*,d}

A comparison between different M-C bonds (M = Cu(I), Ni(II), Co(I), Rh(I) and Ir(I)) has been reported by using density functional theory (DFT) calculations to explore the role of the metal in the fixation or incorporation of CO₂ into such complexes. The systems investigated are various metal based congeners of the Ir-complex **8** [(cod)(iPr)Ir-CCPh], with a ligand scaffold based on cod and iPr ligands (cod=1,5-cyclooctadiene; iPr = 1,3-bis(isopropyl)imidazol-2-ylidene). Results of this study show that the calculated CO₂ insertion barriers follow the trend: Cu(I) (20.8 kcal/mol) < Rh(I) (30.0 kcal/mol) < Co(I) (31.3 kcal/mol) < Ir(I) (37.5 kcal/mol) < Ni(II) (45.4 kcal/mol), indicating that the Cu(I) based analogue is the best CO₂ fixer, while Ni(II) is the worst in the studied series.

Development of catalysts capable of utilising CO₂ as feedstock,¹⁻⁴ possibly in large volume transformations,⁵⁻⁷ is considered to be one of the key challenges in modern chemistry. However, the wide spread use of CO₂ in chemical transformations is limited due to its chemical inertness.⁸ In recent past, despite some attempts with metal-free processes,^{9,10} even organocatalyzed trapping of CO₂ through C-C and C-O bond formation,¹¹ progress has been made in exploring the CO₂ reactivity with a large number of its reactions being insertion into transition metal-X bond (where X = C, O, N) supported with appropriate ligands.¹²⁻¹⁷ These reactions are believed to be one of the most challenging organometallic reactions as CO₂ is thermodynamically stable and kinetically inert.¹⁸⁻²⁴ To this end, a novel organometallic complex as well as chemical mechanism able to perform or explain the capture of CO₂ would be extremely interesting and valuable in directing catalyst design efforts.²⁵⁻⁴⁶ Therefore, a thorough understanding of the mechanism of CO₂ insertion into the M-X bonds is of fundamental importance and could guide advances in the development of this important reaction.

Despite the very recent large number of experimental⁴⁷⁻⁵² and theoretical⁵³⁻⁵⁹ studies focusing on CO₂ insertion, reports on direct

comparison between theoretical and experimental results remain very limited.^{53,60,61} To partially fill this gap, we have recently reported density functional theory based investigations of CO₂ fixation mechanism as proposed by Nolan *et al* for a new Ir(I) complex,⁶¹⁻⁶³ [(Ir(cod)iPr(OH))] (**1**) (cod=1,5-cyclooctadiene; iPr = 1,3-bis(isopropyl)imidazol-2-ylidene) and its derivatives containing Ir-O (complexes **3** and **4**), Ir-N (complex **5**), and Ir-C bonds (complexes **6**, **7**, **8** and **9**; see Scheme 1).⁶⁴ According to the experiments, complexes containing Ir-heteroatom bonds (**1**, **3-5**) display high reactivity towards CO₂, while complexes with Ir-C bonds (**6-9**) are completely unreactive towards the green house gas.

Scheme 1.

^a KAUST Catalysis Center, Physical Sciences and Engineering Division, King Abdullah University of Science and Technology, Thuwal 23955-6900, Saudi Arabia. E-mail: luigi.cavallo@kaust.edu.sa

^b Dipartimento di Scienze Chimiche, Università di Napoli Federico II, Via Cintia, 80126 Napoli, Italy

^c Chemistry Department, College of Science, King Saud University, Riyadh 11451, Saudi Arabia. Email: stevenpnolan@gmail.com

^d Institut de Química Computacional i Catàlisi and Departament de Química, Universitat de Girona, Campus Montilivi, 17071 Girona, Catalonia, Spain. E-mail: albert.poater@udg.edu

Electronic Supplementary Information (ESI) available: Supporting information for this article is given via a link at the end of the document, providing energies, cartesian coordinates, and 3D view for all DFT optimized species discussed in this work. See DOI: 10.1039/x0xx00000x

ARTICLE

Journal Name

Based on the experimental findings, a plausible mechanism for CO₂ insertion into the M-X bonds (X = O or N) proposed by Nolan *et al* is outlined in Scheme 2. Considering complex **A**, the nucleophilic attack of the heteroatom X onto CO₂ does not proceed via the oxygen atoms but through the carbon atom, involving a four-membered cyclic center, hypothetically structure **B**, i.e. most probably **B** is the transition state itself. The particular structural nature of **1** allows for a second molecule of **1** to react yielding a dimeric structural product **2**; this occurs once a water molecule has been released from intermediate **C**.

Scheme 2.

Our detailed investigation of the proposed mechanism in Scheme 2 for complex **1** revealed that the CO₂ insertion step is the rate determining step rather than the subsequent dimerization step, **C**→**2**, (15.8 kcal/mol vs 9.4 kcal/mol, respectively).⁶² This finding contradicts the recent study by Truscott *et al.*⁶¹ as they argued that dimerization is probably the rate-limiting step for complex **1**. In fact, we pointed out that the facile dimerization step is due to the cooperative nature of the iridium centers. Additionally, in agreement with experiments, the calculated low CO₂ insertion barriers for complex **1** and its derivatives **3-5** (**3** (11.8 kcal/mol) < **4** (22.5 kcal/mol) < **5** (17.7 kcal/mol)) indicated that the CO₂ insertion into Ir-O and Ir-N bonds is kinetically facile. On the other hand, significantly higher barrier values were estimated for complexes **6-9** (ranges from 35.0-50.0 kcal/mol), suggesting that CO₂ insertion into Ir-C bonds is kinetically more challenging and supports the experimental observation that the Ir-C bond are unreactive towards CO₂.

Having rationalized the behaviour of the systems proposed in the literature (**1**, **3-9**), in the present study we turn our attention to the effect of changing the metal (but keeping the same ligand scaffold; cod and iPr ligands) with the aim of finding one metal that can induce CO₂ insertion into M-C bonds. It is well documented in the literature that CO₂ can be inserted into the M-C bonds of Rh and other transition metals such as Cu and Ni.⁶⁵⁻⁶⁷ To this end, we studied the CO₂ insertion step for different metal (Cu(I), Rh(I) and Ni(II)) based congeners of **8** [(cod)(iPr)Rh-CCPh],⁶⁸ which presented the lowest CO₂ insertion barrier (37.5 kcal/mol) among the investigated Ir-C bond containing complex in the series **6-9**.

To clarify the role of the metal in the CO₂ insertion DFT calculations were carried out. All DFT static calculations have been performed at the GGA level with the Gaussian09 set of programs,⁶⁹ using the BP86 functional of Becke and Perdew.^{70,71} The electronic configuration of the molecular systems was described with the standard split-valence basis set with a polarization function of Ahlrichs and co-workers for H, C, N, and O (TZVP keyword in

Gaussian).⁷² For Cu, Ni, Rh and Ir, the quasi-relativistic Stuttgart/Dresden effective core potential was used, with the associated valence basis set (standard SDD keywords in Gaussian09).⁷³⁻⁷⁵ The geometry optimizations were carried out without symmetry constraints, and the characterization of the stationary points was performed by analytical frequency calculations. Single point calculations on the BP86 optimized geometries were performed using the M06 functional⁷⁶ with the triple- ζ basis set of Ahlrichs for main-group atoms (TZVP keyword in Gaussian),⁷⁷ while for Ir the SDD basis set has been employed. These single point calculations also used the polarizable continuous model PCM to address solvent effects, using benzene as a solvent.^{78,79} The reported free energies in this work include energies obtained at the M06/TZVP level corrected with zero-point energies, thermal corrections and entropy effects evaluated at 298 K with the BP86/SVP method in the gas phase.^{80,81}

Figure 1 presents the computed energy profile for the CO₂ insertion into M-C bond of different metal analogues of **8**. The reference point is the energy of complex **8** analogues plus free CO₂. For the selected metals, the calculated CO₂ insertion barriers (for simplicity named TS₈, in analogy with our Ir(I)-complexes of Scheme 1) follows the trend: Cu(I) (20.8 kcal/mol) < Rh(I) (30.0 kcal/mol) < Co(I) (31.3 kcal/mol) < Ir(I) (37.5 kcal/mol) < Ni(II) (45.4 kcal/mol), indicating that the Cu(I) based analogue is the best CO₂ fixer, while Ni(II) is the worst of the studied metal series. However, from a thermodynamic stand point, the overall CO₂ insertion reaction is endergonic in nature for all examined metals (see the energy values for **II**₈ in Figure 1), being less endergonic in the case of Cu(I) (5.0 kcal/mol) than Ni(II) (22.3 kcal/mol). It is worth mentioning here that we calculated the CO₂ insertion barrier for high spin Co(II) complex and is predicted to be 32.9 kcal/mol, which is slightly higher in energy (by 1.6 kcal/mol) than the calculated barrier for low spin Co complex (31.3 kcal/mol).

Figure 1. Computed free energy profile for the CO₂ insertion into M-C bond of different metal analogues of Ir(I)-complex **8** (where R¹ = CCPh). Free energies are given in kcal/mol. Ni(II) is a cationic complex (overall charge +1).

Figure 2. Calculated transition states of CO₂ fixation mechanism promoted by different metals analogues of Ir(I)-complex **8**: (a) Cu(I); (b) Rh(I); (c) Co(I); (d) Ir(I); and (e) Ni(II). The most notable distances are given in Å.

A plausible explanation for the remarkable metal effect on the calculated CO₂ insertion barriers can be drawn from the geometries of the TS structures (see Figure 2) and by considering the Hammond postulate. In detail, the calculated high energy TS for Ni(II) can be considered as a textbook example of a “late TS” closer to product ($r(\text{Ni-O}) = 2.269$ Å; see Figure 2e). By changing the metal from Ni(II) to Cu(I), the

transition state moves towards the reactants as the reaction becomes less endothermic. Thus, the calculated low energy TS for Cu(I) can be considered as an “early TS” closer to the reactants ($r(\text{Cu-O}) = 3.053$ Å; see Figure 2a). Intermediate situations are reported for Rh and Ir ($r(\text{Rh-O}) = 2.891$ Å and $r(\text{Ir-O}) = 2.716$ Å; Figure 2b and c, respectively).

Having these results at hand, we then calculated the CO₂ insertion barrier for the experimentally known Cu(I) complex, [Cu(IPr)(CCPh)],⁸² and further, to be compared with other group 11 metals Ag(I) and Au(I). The calculated CO₂ insertion barrier follows the trend: Ag(I) (30.9 kcal/mol) < Cu(I) (34.6 kcal/mol) < Au(I) (37.0 kcal/mol), indicating that the Ag(I) based analogue is the best CO₂ fixer, followed by Cu(I) and Au(I) based analogues. However, the calculated CO₂ insertion barriers are still above 30.0 kcal/mol, suggesting that the CO₂ fixation is kinetically challenging in the considered group 11 metal series. Calculations attempting to screen a longer series of ligands to characterize in detail how the CO₂ insertion barrier is affected by the nature of the ligand are ongoing.^{82,83}

Last but not least, even though the energy data include the correction for dispersion implicitly, the geometry optimization is not included, as previously reported,^{62,79,80} however new tests performed on the transition states for Ni and Cu displayed in Figure 2 reveal an increase in the energy barrier of +0.4 kcal/mol for Ni and a decrease of 0.8 kcal/mol for Cu, which suggest that in absolute value the difference is small to negligible and that there is no obvious trend concerning the effect of dispersion on the CO₂ insertion barriers.

Conclusions

We have explored the role of metal in the CO₂ insertion into M-C bond by considering various metal (M = Cu(I), Co(I), Rh(I) and Ni(II)) analogues of Ir(I)-complex **8**. Results show that among the considered metals, Cu(I) was the best CO₂ fixer with an estimated barrier of 20.8 kcal/mol, while Ni(II) was found to be the least reactive with a barrier of 45.4 kcal/mol. As a final note, the large spread in the activation energies estimated by changing the metal but with the same ligand scaffold (cod and I¹Pr ligands) strongly suggests that there is a large space to explore for a suitable combination of ligand and metal to add important components to improve/affect the thermodynamics of this fascinating reaction.

Acknowledgements

A.P. thanks the Spanish MINECO for a project CTQ2014-59832-JIN, and European Commission for a Career Integration Grant (CIG09-GA-2011-293900). S.V.C.V, S.P.N. and L.C. thank King Abdullah University of Science and Technology (CCF-1974-3 project) for support. G.T. thanks the KAUST for the visiting contract KAUST-2015-C0546.

Notes and references

- 1 M. Aresta, *Carbon Dioxide as Chemical Feedstock*, Wiley-VCH, Weinheim, 2010.

- 1 B. Dutcher, M. Fan and A. G. Russell, *ACS Appl. Mater. Inter.*, 2015, **7**, 2137-2148.
- 2 C. C. Wang, Y. Q. Zhang, J. Li and P. Wang, *J. Mol. Struct.*, 2015, **1083**, 127-136.
- 3 N. von der Assen, P. Voll, M. Peters and A. Bardow, *Chem Soc. Rev.*, 2014, **43**, 7982-7994.
- 4 V. D'Elia, A. A. Ghani, A. Monassier, J. Sofack-Kreutzer, J. D. A. Pelletier, M. Drees, S. V. C. Vummaleti, A. Poater, L. Cavallo, M. Cokoja, J. M. Basset and F. E. Kuhn, *Chem.-Eur. J.*, 2014, **20**, 11870-11882.
- 5 V. D'Elia, H. Dong, A. J. Rossini, C. M. Widdifield, S. V. C. Vummaleti, Y. Minenkov, A. Poater, E. Abou-Hamad, J. D. A. Pelletier, L. Cavallo, L. Emsley and J. M. Basset, *J. Am. Chem. Soc.*, 2015, **137**, 7728-7739.
- 6 M. Cokoja, C. Bruckmeier, B. Rieger, W. A. Herrmann and F. E. Kuhn, *Angew. Chem., Int. Ed.*, 2011, **50**, 8510-8537.
- 7 C. Song, *Catal. Today*, 2006, **115**, 2-32.
- 8 J. Y. Hu, J. Ma, Q. G. Zhu, Z. F. Zhang, C. Y. Wu and B. X. Han, *Angew. Chem., Int. Ed.*, 2015, **54**, 5399-5403.
- 9 E. Theuergarten, T. Bannenberg, M. D. Walter, D. Holschumacher, M. Freytag, C. G. Daniliuc, P. G. Jones and M. Tamm, *Dalton Trans.*, 2015, **43**, 1651-1662.
- 10 Z. Xin, C. Lescot, S. D. Friis, K. Daasbjerg and T. Skrydstrup, *Angew. Chem., Int. Ed.*, 2015, **54**, 6862-6866.
- 11 W. Leitner, *Coord. Chem. Rev.*, 1996, **153**, 257-284.
- 12 D. H. Gibson, *Coord. Chem. Rev.*, 1999, **185-186**, 335-355.
- 13 F. J. Fernández-Alvarez, M. Iglesias, L. A. Oro and V. Polo, *ChemCatChem*, 2013, **5**, 3481-3494.
- 14 G. E. Dobereiner, J. Wu, M. G. Manas, N. D. Schley, M. K. Takase, R. H. Crabtree, N. Hazari, F. Maseras and A. Nova, *Inorg. Chem.*, 2012, **51**, 9683-9693.
- 15 H.-W. Suh, T. J. Schmeier, N. Hazari, R. A. Kemp and M. K. Takase, *Organometallics*, 2012, **31**, 8225-8236.
- 16 T. J. Schmeier, A. Nova, N. Hazari and F. Maseras, *Chem.-Eur. J.*, 2012, **18**, 6915-6927.
- 17 A. Behr, *Angew. Chem., Int. Ed.*, 1988, **27**, 661-678.
- 18 T. Herskovitz and L. J. Guggenberger, *J. Am. Chem. Soc.*, 1976, **98**, 1615-1616.
- 19 D. W. Lee, C. M. Jensen and D. Morales-Morales, *Organometallics*, 2003, **22**, 4744-4749.
- 20 G. Parkin, *Chem. Rev.*, 2004, **104**, 699-768.
- 21 X. Yin and J. R. Moss, *Coord. Chem. Rev.*, 1999, **181**, 27-59.
- 22 J. Langer, W. Imhof, M. J. Fabra, P. García-Orduña, H. Görls, F. J. Lahoz, L. A. Oro and M. Westerhausen, *Organometallics*, 2010, **29**, 1642-1651.
- 23 M. Aresta and A. Dibenedetto, *Dalton Trans.*, 2007, 2975-2992.
- 24 R. A. Allred, L. H. McAlexander, A. M. Arif and L. M. Berreau, *Inorg. Chem.*, 2002, **41**, 6790-6801.
- 25 N. Kitajima, S. Hikichi, M. Tanaka and Y. Morooka, *J. Am. Chem. Soc.*, 1993, **115**, 5496-5508.
- 26 R. Johansson and O. F. Wendt, *Organometallics*, 2007, **26**, 2426-2430.
- 27 Z.-W. Mao, F. W. Heinemann, G. Liehr and R. v. Eldik, *Dalton Trans.*, 2001, 3652-3662.
- 28 U. Singh, P. Babbar, P. Tyagi and T. Weyhermüller, *Transition Met. Chem.*, 2008, **33**, 931-940.
- 29 R. A. Jewsbury, *Inorg. Chim. Acta*, 1981, **49**, 141-143.
- 30 V. D. Bianco, S. Doronzo and N. Gallo, *Inorg. Nucl. Chem. Lett.*, 1979, **15**, 187-189.
- 31 R. S. Haszeldine, *Science*, 2009, **325**, 1647-1652.
- 32 A. Collado, A. Gómez-Suárez, P. B. Webb, H. Kruger, M. Buhl, D. B. Cordes, A. M. Z. Slawin and S. P. Nolan, *Chem. Commun.*, 2014, **50**, 11321-11324.
- 33 B. P. Buffin, A. M. Arif and T. G. Richmond, *Chem. Commun.*, 1993, 1432-1434.
- 34 D. S. Glueck, L. J. N Winslow and R. G. Bergman, *Organometallics*, 1991, **10**, 1462-1479.
- 35 L. J. Newman and R. G. Bergman, *J. Am. Chem. Soc.*, 1985, **107**, 5314-5315. DOI: 10.1039/C5QO00281H
- 36 S. Krogsrud, S. Komiya, T. Ito, J. A. Ibers and A. Yamamoto, *Inorg. Chem.*, 1976, **15**, 2798-2805.
- 37 M. H. Chisholm and M. Extine, *Chem. Commun.*, 1975, 438-439.
- 38 G. Gattow and W. Behrendt, *Angew. Chem., Int. Ed.*, 1972, **11**, 534-535.
- 39 M. Rahim, C. White, A. L. Rheingold and K. J. Ahmed, *Organometallics*, 1993, **12**, 2401-2403.
- 40 J. M. Seul and S. Park, *Bull. Korean Chem. Soc.*, 2010, **31**, 3745-3748.
- 41 S. Park, *Bull. Korean Chem. Soc.*, 2001, **22**, 1410-1412.
- 42 D. J. Darensbourg, G. Groetsch, P. Wiegrefe and A. L. Rheingold, *Inorg. Chem.*, 1987, **26**, 3827-3830.
- 43 R. Johansson, M. Jarenmark and O. F. Wendt, *Organometallics*, 2005, **24**, 4500-4502.
- 44 T. J. Schmeier, N. Hazari, C. D. Incarvito and J. A. Raskatov, *Chem. Commun.*, 2011, **47**, 1824-1826.
- 45 M. T. Johnson, M. Janse Van Rensburg, J.; M. Axelsson, M. S. G. Ahlquist and O. F. Wendt, *Chem. Sci.*, 2011, **2**, 2373-2377.
- 46 F.-Y. Yi, H.-L. Jiang and X.-M. Sun, *Chem. Commun.*, 2015, **51**, 8446-8449.
- 47 G. G. Zhang, Z.-A. Lan and X. C. Wang, *ChemCatChem*, 2015, **7**, 1422-1423.
- 48 T. Kurahashi and S. Matsubara, *Acc. Chem. Res.*, 2015, **48**, 1703-1716.
- 49 I. P. Koronaki, L. Prentza and V. Papaefthimiou, *Renew. Sust. Energ. Rev.*, 2015, **50**, 547-566.
- 50 D. Yu, S. P. Teong and Y. Zhang, *Coord. Chem. Rev.*, 2015, **293**, 279-291.
- 51 J. Jeromenok, W. Böhlmann, C. Jäger and J. Weber, *ChemistryOpen*, 2013, **2**, 17-20.
- 52 A. Paparo, J. S. Silvia, C. E. Kefalidis, T. P. Spaniol, L. Maron, J. Okuda and C. C. Cummins, *Angew. Chem., Int. Ed.*, 2015, **54**, 9115-9119.
- 53 S. Marmitt and P. F. B. Goncalves, *J. Comput. Chem.*, 2015, **36**, 1322-1333.
- 54 S. Gennen, M. Alves, R. Mereau, T. Tassaing, B. Gilbert, C. Detrembleur, C. Jerome and B. Grignard, *ChemSusChem*, 2015, **8**, 1845-1849.
- 55 K. Sokolowski, W. Bury, A. Tulewicz, A. M. Cieslak, I. Justyniak, D. Kubicki, E. Krajewska, A. Milet, R. Moszynski and J. Lewinski, *Chem.-Eur. J.*, 2015, **21**, 5496-5503.
- 56 S. Rasul, D. H. Anjum, A. Jedidi, Y. Minenkov, L. Cavallo and K. Takanabe, *Angew. Chem., Int. Ed.*, 2015, **54**, 2146-2150.
- 57 T. Ema, Y. Miyazaki, J. Shimonishi, C. Maeda and J.-Y. Hasegawa, *J. Am. Chem. Soc.*, 2014, **136**, 15270-15279.
- 58 W. K. Offermans, C. Bizzarri, W. Leitner and T. E. Müller, *Beilstein J. Org. Chem.*, 2015, **11**, 1340-1351.
- 59 T. Fan, X. Chen and Z. Lin, *Chem. Commun.*, 2012, **48**, 10808-10828.
- 60 B. J. Truscott, H. Kruger, P. B. Webb, M. Bühl and S. P. Nolan, *Chem.-Eur. J.*, 2015, **21**, 6930-6935.
- 61 S. V. C. Vummaleti, S. P. Nolan, L. Cavallo, G. Talarico and A. Poater, *Eur. J. Inorg. Chem.*, 2015, 4653-4657.
- 62 B. J. Truscott, D. J. Nelson, C. Lujan, A. M. Z. Slawin and S. P. Nolan, *Chem.-Eur. J.*, 2013, **19**, 7904-7916.
- 63 B. J. Truscott, D. J. Nelson, A. M. Z. Slawin and S. P. Nolan, *Chem. Commun.*, 2014, **50**, 286-288.
- 64 T. Suga, H. Mizuno, J. Takaya and N. Iwasawa, *Chem. Commun.*, 2014, **50**, 14360-14363.
- 65 I. S. Kolomnikov, A. O. Gusev, T. S. Belopotapova, M. Kh. Grigoryan, T. V. Lysyak, Y. T. Struchkov and M. E. Vol'pin, *J. Organomet. Chem.*, 1974, **69**, C10-C12.
- 66 K. Ukai, M. Aoki, J. Takaya and N. Iwasawa, *J. Am. Chem. Soc.*, 2006, **128**, 8706-8707.
- 67 B. J. Truscott, G. C. Fortman, A. M. Z. Slawin and S. P. Nolan, *Org. Biomol. Chem.*, 2011, **9**, 7038-7041.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Journal Name

ARTICLE

69 Gaussian 09, Revision C.01, M. J. Frisch et al, Gaussian, Inc., Wallingford CT, 2009.

70 A. Becke, *Phys. Rev. A*, 1988, **38**, 3098-3100.

71 J. P. Perdew, *Phys. Rev. B*, 1986, **33**, 8822-8824.

72 A. Schaefer, H. Horn and R. Ahlrichs, *J. Chem. Phys.*, 1992, **97**, 2571-2577.

73 U. Haeusermann, M. Dolg, H. Stoll and H. Preuss, *Mol. Phys.*, 1993, **78**, 1211-1224.

74 W. Kuechle, M. Dolg, H. Stoll and H. Preuss, *J. Chem. Phys.*, 1994, **100**, 7535-7542.

75 T. Leininger, A. Nicklass, H. Stoll, M. Dolg and P. Schwerdtfeger, *J. Chem. Phys.*, 1996, **105**, 1052-1059.

76 Y. Zhao and D. G. Truhlar, *Theor. Chem. Acc.*, 2008, **120**, 215-241.

77 F. Weigend and R. Ahlrichs, *Phys. Chem. Chem. Phys.*, 2005, **7**, 3297-3305.

78 V. Barone and M. Cossi, *J. Phys. Chem. A*, 1998, **102**, 1995-2001.

79 J. Tomasi and M. Persico, *Chem. Rev.*, 1994, **94**, 2027-2094.

80 A. Poater, E. Pump, S. V. C. Vummaleti and L. Cavallo, *J. Chem. Theory Comput.*, 2014, **10**, 4442-4448.

81 J. P. Martínez, M. Solà and A. Poater, *ChemistryOpen*, 2015, DOI: 10.1002/open.201500093.

82 G. C. Fortman, A. M. Z. Slawin and S. P. Nolan, *Organometallics*, 2010, **29**, 3966-3972.

83 A. H. Christian, P. Müller and S. Monfette, *Organometallics*, 2014, **33**, 2134-2137.

View Article Online
DOI: 10.1039/C5QO00281H

Organic Chemistry Frontiers Accepted Manuscript