DOI: 10.1016/j.marpolbul.2015.11.045

REFERENCE

Robitzch V.S.N., Lozano-Cortés D, Kandler NM, Salas E, Berumen ML
(2015) Productivity and sea surface temperature are correlated with the pelagic larval
duration of damselfishes in the Red Sea. Marine Pollution Bulletin.
doi:10.1016/j.marpolbul.2015.11.045

ARTICLE TYPE

Research Report

TITLE

Productivity and sea surface temperature are correlated with the pelagic larval duration of
damselfishes in the Red Sea

CORRESPONDING AUTHOR

Vanessa S. N. Robitzch

Affiliation: Red Sea Research Center, Division of Biological and Environmental Science
and Engineering, King Abdullah University of Science and Technology, Thuwal, 23955-
6900, Saudi Arabia; Phone: +966 54 0497300;

Email: Vanessa.RobitzchSierra@kaust.edu.sa

AUTHORS

Vanessa S. N. Robitzch¹

Diego Lozano-Cortés^{1,2}

Nora M. Kandler¹

Eva Salas^{3,4}

Michael L. Berumen¹

¹ Red Sea Research Center, Division of Biological and Environmental Science and
Engineering, King Abdullah University of Science and Technology, Thuwal, 23955-
6900, Saudi Arabia

² Coral Reef Ecology Research Group, Department of Biology, Universidad del Valle,
Apartado Aéreo 25360, Cali, Colombia

³ Department of Ecology and Evolutionary Biology, University of California, 100
Shaffer road, Santa Cruz, California, 95060, USA

⁴ Section of Ichthyology, California Academy of Sciences, 55 Music Concourse Drive,
San Francisco, California, 94118, USA

ABSTRACT

We examined the variation of pelagic larval durations (PLDs) among three damselfishes, *Dascyllus aruanus*, *D. marginatus*, and *D. trimaculatus*, which live under the influence of an environmental gradient in the Red Sea. PLDs were significantly correlated with latitude, sea surface temperature (SST), and primary production (CHLA; chlorophyll a concentrations). We find a consistent decrease in PLDs with increasing SST and primary production (CHLA) towards the southern Red Sea among all species. This trend is likely related to higher food availability and increased metabolic rates in that region. We suggest that food availability is a potentially stronger driver of variation in PLD than temperature, especially in highly oligotrophic regions. Additionally, variations in PLDs were particularly high among specimens of *D. marginatus*, suggesting a stronger response to local environmental differences for endemic species. We also report the first average PLD for this species over a broad geographic range (19.82 ± 2.92 days).

KEYWORDS

Biogeography; Endemic; Environment; Food Availability; PLD; Reef Fish

1. INTRODUCTION

The time a fish larva spends in the plankton before recruiting onto a reef, known as its pelagic larval duration (PLD), is a valuable source of information on how ecological and oceanographic processes affect a fish's early life. Such information can help to better understand organisms with complex life cycles, such as many coral reef inhabitants; subsequently, this fundamental life history information can be used to improve management of marine ecosystems and their natural resources. PLDs of coral reef fishes

are known to exhibit variation driven by a wide range of intrinsic and extrinsic factors (McCormick and Molony, 1995; Searcy and Sponaugle, 2000; Sponaugle, 2010; Wellington and Victor, 1992; Wilson and Meekan, 2002). Two of the most relevant extrinsic factors that cause intraspecific variations in larval growth and development are temperature and food availability (Heath, 1992; Houde and Zastrow, 1993; Mcleod et al., 2011; Takahashi and Watanabe, 2005), which can be estimated from remote sensing satellite data on sea surface temperature (SST) and chlorophyll a (CHLA) concentrations respectively. Warmer temperatures increase metabolic rate and growth rates (Green and Fisher, 2004; Meekan et al., 2003; Sponaugle et al., 2006), which can lead to shorter PLDs (Bergenius et al., 2005; Green and Fisher, 2004; Grorud-Colvert and Sponaugle, 2011; McCormick and Molony, 1995; Sponaugle et al., 2006). Similarly, higher food availability can increase larval growth (Bikhtashev et al., 2003; Fontes et al., 2010; Houde and Hoyt, 1987; Houde, 1989; Landaeta and Castro, 2006; Sponaugle, 2010, 2009) and potentially decrease the time a fish larva spends in the pelagic before recruiting onto a coral reef (Sponaugle and Grorud-Colvert, 2006). Variations in PLDs can be indicative of differences in dispersal potential, growth rates, and survival of fish larvae. Furthermore, the conditions a fish larva experiences during its PLD seem to even influence success, fitness, and growth in later life stages as juvenile and adult fish (Cushing and Horwood, 1994; Houde and Hoyt, 1987; Houde, 1989; Rankin and Sponaugle, 2011; Sponaugle and Grorud-Colvert, 2006; Sponaugle et al., 2011). However, most studies that have assessed the influence of temperature and food availability on the larval stage of fishes have focused on larval growth rate (Mcleod et al., 2015), size at settlement (McCormick and Molony, 1995), or recruitment cohort size (Lo-Yat et al., 2011) rather than on the length of PLDs in the standing population, and still fewer of these studies target coral reef fishes. In our study we focus specifically on PLD variations within the standing populations (i.e., post-settlement and adult fishes, as opposed to recruitment cohorts) of three congeneric coral reef damselfishes present in the Red Sea.

The Red Sea is an ideal location to study the effect of a variety of environmental parameters on coral reefs (Berumen et al., 2013). The Red Sea harbors thriving and continuous coral reefs along both sides of its narrow basin. Due to its shape, location, and only a single shallow and narrow connection to the Indian Ocean, it displays a unique

environmental gradient of temperature, salinity, productivity, turbidity, and essentially reef-scape (Racault et al., 2015; Raitos et al., 2013). These physical and environmental characteristics have made the Red Sea a natural location to assess the environmental impact on several coral reef organisms (e.g., Froukh and Kochzius, 2007; Giles et al., 2015; Lozano-Cortés and Berumen, 2015; Nanninga et al., 2014; Ngugi et al., 2012; Roberts et al., 1992; Robitzch et al., 2015; Sawall et al., 2014a, 2014b).

Here, we focus on the two seemingly major parameters that may drive PLD variations in the environmental gradient of the Red Sea: temperature and food availability. For this purpose we use the three *Dascyllus* species that live in the Red Sea (*D. aruanus*, *D. marginatus*, and *D. trimaculatus*). These species serve as a good model group to study PLD variations in coral reef fishes for several reasons. They are biologically similar, small in size, and commonly present along the entire environmental gradient of the Red Sea at overlapping depth ranges. They are zooplanktivorous, demersal spawners, and have relatively similar recruitment periods and larval development times (PLD). However, they have subtly different ecological preferences, which make them an interesting group to study the effect of the environment on early life history traits in the context of ecological specialization. We consider *D. trimaculatus* as the most ecologically versatile species within this group. It can be found on exposed reef walls, sheltered backside-reefs, and lagoons; it is not dependent on live coral; and it only uses reef structure for shelter, recruitment, and as spawning habitat. While *D. trimaculatus* mostly recruits onto anemones in the Red Sea, it can also recruit into branching corals, sea urchins, and other microhabitats. Once they reach juvenile stages, *D. trimaculatus* leave their settlement substrate, range throughout the entire reef structure, and spend most of their lifetime foraging in the water column. Intermediate on the ecological specialization ranking is *D. aruanus*. This species shows stronger habitat preferences, only recruiting to and living its entire life in branching live corals that are located in sheltered sandy reefs and lagoons. *Dascyllus aruanus* and our third species, *D. marginatus*, both show similar habitat preferences and occasionally even co-inhabit the same coral colony. However, the degree of ecological specialization is stronger in *D. marginatus*, which cannot be found on very shallow reef-tops and lagoons, nor can it be

found on offshore reefs in most of the Red Sea. Therefore, we refer to *D. marginatus* as the most ecologically specialized species in our group.

The *Dascyllus* model also seems suitable to study variation in PLDs in the context of biogeography of coral reef associated fishes. Of the three species that occur in the Red Sea, *D. marginatus* is an Arabian endemic while *D. trimaculatus* and *D. aruanus* are both widespread species, common in most of the Indo-Pacific. In general, pomacentrid species with restricted distributions have shorter PLDs than wide-ranging species (Cowen and Sponaugle, 1997; Wellington and Victor, 1989), although the opposite is observed for labrids (Cowen and Sponaugle, 1997; Victor, 1986). The expected trend is that endemic or geographically-constrained species have shorter PLDs potentially restricting dispersal (Lester and Ruttenberg, 2005; Macpherson and Raventos, 2006; Mora et al., 2003). However, in endemic pomacentrid species the PLDs seem to have a rather non-linear relationship with range size (Thresher et al., 1989). Similarly, most studies assessing correlations between PLDs and biogeographic ranges over a number of species find no link between the two (Macpherson and Raventos, 2006; Victor and Wellington, 2000; Zapata and Herrón, 2002). The exact nature of PLD vs. biogeography remains an open question.

Thus, the objectives of our study are to use the model species-group of *D. marginatus*, *D. aruanus*, and *D. trimaculatus* to 1) assess the influence of environmental parameters such as SST and CHLA on the PLDs of coral reef associated damselfishes with different degrees of ecological specialization in a natural environment inside the Red Sea; 2) investigate variations in PLDs related to differences in biogeographic ranges among closely related species; 3) provide the first PLD-measurements for *D. trimaculatus* and *D. aruanus* from the Red Sea and additional measurements from two sites outside the Red Sea as well as 4) provide the first PLD-measurements the first PLDs for *D. marginatus* from a wide geographic range.

2. MATERIALS AND METHODS

2.1 Fish sampling:

Clove oil, tweezers, hand nets, and spears were used to collect juvenile and adult specimen from the three *Dascyllus* species (*D. aruanus*, *D. marginatus*, and *D.*

1 *trimaculatus*) present in the Red Sea (RS) (Fig. 1). Fishes of different sizes (Fig. 2)
2 and/or from different sampling years (Tab. 1) were collected for PLD assessment to cover
3 potential temporal variations as much as possible. These were caught at 18 reef sites off
4 the Saudi Arabian coastline and at two sites outside the RS (Indian Ocean “IO” and Indo-
5 Australian-Archipelago “IA”; Tab. 1 and Fig. 1). Samples were immediately preserved in
6 90% ethanol. The sampling range (>1600 km of coastline) covers most of the latitudinal
7 span of the RS.

8

9 **Fig. 1: Sampling sites of *Dascyllus marginatus*, *D. aruanus*, and *D. trimaculatus*.** A
10 total of 20 sites are grouped into four regions within the Red Sea (RS) and two outside
11 the Red Sea (IA: Indo-Australian Archipelago; and IO: Indian Ocean). Those inside the
12 Red Sea are represented by white circles (GAQ: Gulf of Aqaba), squares (NCRS: north-
13 central RS), diamonds (SCRS: south-central RS), or sails (SRS: southern RS) according
14 to the respective region. The regions were assigned in consensus with those in Raitso et
15 al. (2013). The two locations outside the Red Sea are represented by a white nabla and a
16 white triangle (IO: Indian Ocean and IA: Indo-Australian-Archipelago, respectively).
17 Map axes indicate latitude (°N) and longitude (°E). Additional information on exact
18 number of samples and coordinates for each site can be found in Tab. 1. The chlorophyll
19 *a* (CHLA, in mg m⁻³) concentrations of the Indo-Pacific and Red Sea are displayed from
20 the NASA Giovanni website (<http://oceancolor.gsfc.nasa.gov>) to visualize approximate
21 differences between locations.

Fig. 2: Distribution of fish sizes (total length, in mm) of *Dascyllus marginatus* (left panel) and *D. aruanus* (right panel) used for PLD measurements from each site. Sites are given with three letter abbreviations on the x-axis (reef names can be found in Tab. 1). The sites are in latitudinal order from north to south from left to right, respectively along the axis. Black circles represent mean sizes and the whiskers display the maximum and minimum sizes of the fishes.

Tab. 1: Sampling sites, including number of samples, reef names and coordinates.

Location	Region	Site	N (Dt)	N (Dm)	N (Da)	Latitude	Longitude	Reef Name
RS	GAQ	HAQ	2	7	7	29°15'11.45"N	34°56'20.11"E	Haql
	NCRS	YRM	1	-	-	23°46'21.12"N	38°16'34.02"E	Yanbu-Ras Majiz
		RMA	-	9	10	23°18'36.12"N	38°26'12.48"E	Ras Masturah
		AKA	-	3	11	22°56'15.90"N	38°45'56.58"E	Shib Al Karrah
		SNA	2	-	12	22°20'51.96"N	38°51'09.42"E	Shib Nazar
		FSA	-	8	-	22°17'46.80"N	39°04'26.64"E	Inner Fsar
		AFA	4	-	-	22°17'49.20"N	38°57'32.40"E	Al Fahal
		SAK	-	-	7	21°40'16.53"N	38°50'36.60"E	Shib Al Kabir
		AMA	-	-	10	22°04'25.75"N	38°46'40.26"E	Abu Madafi
	Total		7	20	50			
	NSRS	MAN	4	7	8	20°08'05.10"N	40°06'04.39"E	Manila Bay
		SAU	2	-	7	19°53'15.43"N	40°09'23.94"E	Saut Reef
		LCG	-	2	-	20°09'44.52"N	40°13'36.72"E	Coast Guard Reef
		Total	6	9	15			
SRS		GHU	-	-	1	17°06'37.20"N	42°04'03.10"E	Ghurab
		DAH	2	-	-	16°52'22.60"N	41°26'24.50"E	Dhi Dahaya
		ZDU	1	7	-	16°50'03.30"N	42°18'38.20"E	Zahrat Durakah
		BAG	1	-	1	16°58'44.00"N	41°23'05.60"E	Al Baglah
		DUR	-	13	-	16°51'36.20"N	42°19'18.00"E	Durakah
		DUM	4	-	11	16°33'50.80"N	42°03'30.60"E	Dumsuq
		Total	8	20	13			
IA	OUT	IND	-	-	7	02°11'11.04"N	118°32'55.33"E	Manado Nth Sulawesi
IO	OUT	COC	5	-	-	12°7'42.24"S	96°55'0.42"E	Cocos Keeling
Total			28	56	92			

The three main geographic locations are the Red Sea (RS), the Indo-Australian-Archipelago (IA), and the Indian Ocean (IO). The RS location is subdivided in four major

regions (GAQ: Gulf of Aqaba, NCRS: north-central RS, SCRS: south-central RS, and SRS: south RS). The sites represent the reefs sampled for each region (three letter codes based on the reef names). The IA and IO locations are represented by one sampling site each: Mandando in North Sulawesi, Indonesia (IND) and the island of Cocos Keeling, Australia (COC) respectively. N represents the number of individuals for which pelagic larval durations (PLDs) were assessed for statistical analyses of each studied species: *D. trimaculatus* (Dt), *D. marginatus* (Dm), and *D. aruanus* (Da). Samples from the RS were collected in 2013, 2014, and 2015; from the IO in 2010 and 2014; and from the IA in 2003.

Whenever possible, specimens from all three species were collected from the same reef site for coherent interspecific comparisons. In case of absence of any of the species, specimens from the nearest possible site were sampled. The sampling sites were divided into four major Red Sea (RS) regions: “GAQ” (Gulf of Aqaba), “NCRS” (north-central RS), “SCRS” (south-central RS); “SRS” (southern RS) (in consensus with Raitso et al., 2013; Fig. 1 and Tab. 1).

2.2 Otolith preparation and PLD assessment:

PLDs for all three species were estimated from daily increments up to the settlement marks of sagittal otoliths. Sagittae were extracted and mounted onto a microscopy glass slide with thermoplastic resin, then ground and polished (following Wilson and McCormick, 1997) using aluminum oxide lapping films (South Bay Technology, Inc.) of three different thicknesses (30, 12, and 5 micrometers [μm]). Three to ten pictures were taken of each polished otolith with the program AxioVision Rel. (V. 4.8.2.0; copyright 2006-2010 Carl Zeiss Micro Imaging GmbH) utilizing a Zeiss AXIO Scope A1 microscope (200x magnification) and a Zeiss AxioCam ICc1. From the pictures, three independent PLD readings were made and if the counts deviated by less than 10% the mean count for each individual was calculated and included in the analyses. The same reader calculated PLDs for all species and lacked any information or metadata about the origin of the sample. A total of 28 *D. trimaculatus*, 56 *D. marginatus*, and 92 *D. aruanus* otolith readings were used for the statistical analyses (Tab. 1).

2.3 Chlorophyll a (CHLA) and sea surface temperature (SST) correlations:

Two environmental parameters of the Red Sea were tested for correlations to PLD data: Chlorophyll *a* (CHLA) and sea surface temperature (SST). CHLA content is a good proxy for phytoplankton biomass (Håkanson et al., 2007) and is herein used as a proxy of food availability for the pelagic larva of the three studied species. Validated regional averages of CHLA concentrations (mg m^{-3}) and SST ($^{\circ}\text{C}$) were provided by D. Raitzos, based on Raitzos et al. (2013) for the regions NRS, NCRS, SCRS, and SRS; and by D. Dreano for the Gulf of Aqaba (GAQ). These estimates were produced from a 10-year high-resolution data set of satellite remote-sensing CHLA estimates (see Raitzos et al., (2013) for a detailed description of data acquisition). These estimates were subsequently used for Pearson correlations with PLD for each species.

2.4 Statistical analyses:

All data were tested for normality (Kolmogorov–Smirnov test) and homoscedasticity (Levene’s test) prior to analyses. The differences in PLD among the three species in the four Red Sea regions (GAQ, NCRS, SCRS, and SRS) were explored with a factorial ANOVA. The spatial variation of the PLD within each species was tested using an independent one-way ANOVA for each of the three *Dascyllus* species. A post-hoc test applying the unequal N Tukey’s HSD means comparisons was used to detect any significant differences within the two aforementioned analyses. Subsequently, the PLD data was correlated (Pearson correlation) with latitude, SST, and CHLA (i.e., food availability) measurements to assess the impact of the latitudinal environmental gradient of the Red Sea. Finally, the PLDs from the Red Sea sites were compared against those from locations outside the Red Sea for the two widespread species (*D. trimaculatus* and *D. aruanus*). For this analysis, data from the four Red Sea regions for each of the two species was merged and compared independently with the IO (*D. trimaculatus*) and the IA (*D. aruanus*) using a t-student test. All analyses were conducted with STATISTICA 8.0 (StatSoft, Inc. 2007).

3. RESULTS

3.1 PLD measurements:

3.1.1 Within the Red Sea:

The total mean PLDs of all three species differed significantly from each other ($F_{(2,144)}=50.453$, $p<0.001$, Fig. 3). Overall, *D. trimaculatus* showed the highest mean PLD (24.33 ± 2.02), followed by *D. aruanus* (23.09 ± 1.92), while *D. marginatus* exhibited the lowest (19.82 ± 2.92) (Tab. 2). Significant interspecific differences in PLDs were found among regions ($F_{(3,144)}=31.723$, $p<0.001$, Fig. 5) and intraspecific differences among all three species: between sampling regions ($F_{(3,144)}=31.723$, $p<0.001$, Fig. 5) and sites (only assessed for *D. marginatus* ($F_{(7,44)}=10.099$, $p<0.001$) and *D. aruanus* ($F_{(11,76)}=7.6954$, $p<0.001$ due to low number of *D. trimaculatus* specimen per site; see Fig. 5). The interaction between species and regions was not significant. In addition, a consistent latitudinal decrease in PLD was found to be present from north to south among all of the species between the four regions (see Fig. 4). PLD ranges were the largest for the endemic species *D. marginatus* (15-28 d; 13 d difference) and almost double of the ranges found in *D. aruanus* (19-27 d; 8 d difference) and *D. trimaculatus* (20-27 d; 7 d difference). Within sampling sites, individuals of the endemic species also showed the highest PLD range (19-28 d, 9 d difference in HAQ) compared to those of the two widespread species (*D. aruanus*: 6 d difference in MAN; *D. trimaculatus*: 4 d difference in AFA).

Fig. 3: Mean pelagic larval duration (PLD, in days) for *Dascyllus marginatus*, *D. aruanus*, and *D. trimaculatus* averaged among all study sites in the Red Sea. Whiskers represent 95% confidence intervals around the mean (black circle).

Tab. 2: Average pelagic larval durations (PLD in days; mean $\pm$ standard deviation) of *Dascyllus marginatus*, *Dascyllus aruanus*, and *Dascyllus trimaculatus* based on replicate counts of daily growth increments from otoliths.

Locat.	Region	Site	PLD Dm	PLD Da	PLD Dt	Lat.	mean SST [° C]	mean CHLA [mg m ⁻³]
RS	GAQ	HAQ	23.43 $\pm$ 2.89	25.10 $\pm$ 0.94	26.50 $\pm$ 1.17	29.253	24.2	0.20
	(NRS)						25.8	0.15
	NCRS	YRM	-	-	27.00	23.773		
		RMA	19.56 $\pm$ 1.97	23.77 $\pm$ 1.32	-	23.310		
		AKA	24.55 $\pm$ 2.04	24.61 $\pm$ 1.89	-	22.938		
		SNA	-	23.45 $\pm$ 1.46	26.67 $\pm$ 0.94	22.348		
		FSA	21.67 $\pm$ 1.00	-	-	22.296		
		AFA	-	-	24.92 $\pm$ 2.06	22.297		
		SAK	-	23.24 $\pm$ 1.33	-	21.671		
		AMA	-	23.73 $\pm$ 1.37	-	22.074		
		Total	20.65 $\pm$ 2.33	23.79 $\pm$ 1.52	25.72 $\pm$ 1.81		27.2	0.14
	SCRS	MAN	19.29 $\pm$ 2.69	21.95 $\pm$ 1.77	24.58 $\pm$ 1.20	20.135		
		SAU	-	22.52 $\pm$ 1.14	24.84 $\pm$ 0.23	19.888		
		LCG	20.67 $\pm$ 0.94	-	-	20.162		
		Total	19.59 $\pm$ 2.43	22.22 $\pm$ 1.48	24.67 $\pm$ 0.94		28.8	0.27
	SRS	GHU	-	21.00	-	17.110		
		DHI	-	-	22.00 $\pm$ 0.95	16.873		
		ZDU	20.09 $\pm$ 0.71	-	20.33	16.834		
		BAG	-	24.00	25.33	16.979		
		DUR	17.83 $\pm$ 1.63	-	-	16.860		
		DUM	-	20.09 $\pm$ 0.71	22.17 $\pm$ 1.57	16.564		
		Total	17.34 $\pm$ 1.43	20.63 $\pm$ 1.44	22.29 $\pm$ 1.76		28.9	1.64
	Total		19.82 $\pm$ 2.92	23.22 $\pm$ 1.96	24.32 $\pm$ 2.17			
IO			-	-	24.40 $\pm$ 1.25	12.128		
IA			-	21.05 $\pm$ 1.10	-	2.186		
All			19.82 $\pm$ 2.92	23.09 $\pm$ 1.92	24.33 $\pm$ 2.02			
R ² [%]	Dm					43.6	39.4	33.0
	Da					34.8	30.6	27.6
	Dt					46.2	30.1	50.1

Average PLDs for each species (*D. marginatus* (Dm), *D. aruanus* (Da), and *D. trimaculatus* (Dt)) are given among all (All) and within each geographic location: of the Indian Ocean (IO), the Indo-Australian-Archipelago (IA), and the Red Sea (RS); of each region within the RS (GAQ: Gulf of Aqaba, NCRS: north-central RS, SCRS: south-central RS, and SRS: south RS); and of each site within the RS (three-letter coded reef name). The latitude (Lat., in decimal degrees) is given for each site as used for correlation analysis. Estimates of mean surface sea temperature (mean SST, in °C) and chlorophyll *a* content (CHLA, in mg m⁻³) are provided by D. Raitos and D. Dreano from 10-years satellite remote sensing high-resolution data (in consensus with Raitos et al., 2013).

Fig. 4: Mean pelagic larval durations (PLD, in days) of three *Dascyllus* species in the four regions of the Red Sea (RS) (GAQ: Gulf of Aqaba, NCRS: north-central RS, SCRS: south-central RS, and SRS: south RS). Average PLDs are given by blue triangles for *D. marginatus*, red squares for *D. aruanus*, and green diamonds for *D. trimaculatus*. Whiskers represent 95% confidence intervals around the mean.

3.1.2 Inside vs. outside the Red Sea:

We report the PLD measurements for *D. aruanus* and *D. trimaculatus* from sites outside the Red Sea (RS) (*D. aruanus*: 21.05 ± 1.10 in the IA; *D. trimaculatus*: 24.40 ± 1.25 in the IO) and compare these to our measurements from inside the Red Sea. The mean PLD of both widespread species is longer outside the Red Sea than within the Red Sea but only significantly longer for *D. aruanus* (*D. aruanus*: RS vs. IA: t-value=2.88, df=86, p=0.04; *D. trimaculatus*: RS vs. IO: t-value=0.07, df=26, p=0.93; see Fig. 5).

Fig. 5: Mean pelagic larval duration (PLD, in days) of *Dascyllus aruanus* and *D. trimaculatus* from locations inside the Red Sea (RS) and outside the Red Sea (Indo-Australian Archipelago (IA) and Indian Ocean (IO)). Whiskers represent 95% confidence intervals and boxes represent one standard error around the mean. Mean PLDs are given by black squares for *D. aruanus* and diamonds for *D. trimaculatus*.

3.2 Environmental correlations with PLD in the Red Sea:

Regional means of SST and CHLA concentrations were used as environmental data for Pearson correlations (Tab. 2). Their respective mean annual profiles are visualized in Fig. 6. The profiles of each region display a similar trend-curve except for SRS and GAQ. The SRS has overall much higher CHLA concentrations and its average annual CHLA profile has an additional peak in the summer. Different from all other regional CHLA profiles, the GAQ displays a sharp peak later in the spring and has both the highest concentrations of CHLA over the entire Red Sea during the spring and the lowest during the summer (Fig. 6). These unique seasonal changes of CHLA in GAQ result in one of the largest regional ranges ($>0.6 \text{ mg m}^{-3}$) in CHLA concentrations in the Red Sea (besides the southernmost SRS region; Tab. 2 and Fig. 6, left panel). In contrast, the SST profiles for all regions have similar ranges of about 6°C (Fig. 6, right panel).

Fig. 6: Validated monthly regional averages of chlorophyll *a* concentrations (CHLA, in mg m^{-3} , left panel) and sea surface temperature (SST, in $^\circ\text{C}$, right panel). Each panel shows monthly mean averages for an average calendar year (three-letter coded months, x-axis) for each region (GAQ: Gulf of Aqaba, NCRS: north-central RS, SCRS: south-central RS, and SRS: south RS). Averages were produced from 10-year high-resolution data of satellite remote-sensing CHLA and SST estimates provided by D. Raitsos (for the regions NRS, NCRS, SCRS and SRS; sensu Raitsos et al. (2013)) and D. Dreano for the Gulf of Aqaba (GAQ). The trend lines are color-coded for each region.

3.2.1 Correlations with Latitude:

Pearson correlation tests show a positive highly significant ($p < 0.001$) correlation between latitude and PLD among all species (Fig. 7). Latitude is also significantly negatively correlated with both CHLA and SST, indicating the latitudinal character of the Red Sea environmental gradient ($p < 0.001$). Even though latitude shows the highest correlations with PLDs (except among *D. trimaculatus*, $R^2 = 46.2\%$, Tab. 2), latitude itself is most likely not the driving factor but the correlation is rather a result of the combined effects of the environmental parameter at each site (e.g. SST and CHLA).

Fig. 7: Plots of Pearson correlations of pelagic larval durations (PLD, in days, y-axis) of three *Dascyllus* species (*Dascyllus marginatus*, *D. aruanus*, *D. trimaculatus*, from left to right) to: **A-C)** regional means of sea surface temperature (SST in °C, first row), **D-F)** regional means of chlorophyll *a* concentrations (CHLA, in mg m⁻³, middle row), and **G-I)** latitude (in degrees, last row; values in Tab. 2). The plots are arranged in columns according to species (i.e., *D. marginatus* shown in panels A, D, G; *D. aruanus* in panels B, E, H; and *D. trimaculatus* in panels C, F, I) and regional means of: SST, CHLA, and latitude. The blue circles represent the measured PLD values of each species (y-axis, in

1 days). The regression line is in red and the dotted curves represent the respective 95%
2 confidence intervals of the correlation tests.

3.2.2 Correlations with SST:

5 Temperature (SST) is significantly negatively correlated with PLDs of all three species
6 (longer PLDs at lower SST) and shows the second highest R^2 values (except among *D.*
7 *trimaculatus*, $R^2 = 30.1\%$, Tab. 2). This correlation is negative and significant for all
8 species (Fig. 7).

3.2.3 Correlations with CHLA:

11 CHLA concentration is significantly ($p < 0.001$) negatively correlated to the PLDs of all
12 three *Dascyllus* species. The correlation is the strongest for *D. trimaculatus* ($R^2 = 50.1\%$;
13 Tab. 2, Fig. 7). However, the sample size for this species is low and limits the overall
14 interpretation of this result. Moreover, the R^2 values of SST and CHLA do not differ much
15 and make it difficult to interpret which of the two parameters has the stronger influence
16 on PLDs.

4. DISCUSSION AND CONCLUSIONS

19 Our study shows that there is a significant link between the pelagic larval duration (PLD)
20 of *Dascyllus* species and environmental parameters such as mean sea surface temperature
21 (SST) and chlorophyll *a* (CHLA) content in the Red Sea. In all three study species, we
22 found a gradual and consistent decrease in mean PLD with decreasing latitude along the
23 environmental gradient. Moreover, intra-specific differences in PLD between sampling
24 regions along this gradient were higher in *D. marginatus* than the other two widespread
25 species (*D. aruanus* and *D. trimaculatus*), suggesting that the endemic species might be
26 more responsive to environmental differences within its habitat potentially related to a
27 higher degree of specialization. We provide the first PLD estimate for a broad geographic
28 range of the endemic *D. marginatus* (19.8 ± 2.9 d), which displays a consistently lower
29 PLD at all sites in comparison to PLDs of *D. aruanus* and *D. trimaculatus*. Our results
30 suggest that environmental factors can act as drivers of biogeography (as well as
31 endemism) and dispersal among coral reef fishes. We also find the PLDs within species

1 to be overall spatially and potentially also temporally highly plastic, which should be
2 considered when evaluating correlations of PLDs and other parameters.

3 4 **4.1 Pelagic larval durations (PLDs) in the chlorophyll a (CHLA) and sea surface** 5 **temperature (SST) gradient of the Red Sea**

6
7 The Red Sea is a good living laboratory for examining the effect of strong environmental
8 gradients on various aspects of community ecology (Roberts et al., 1992; Ngugi et al.,
9 2012; Sawall et al., 2014a,b) and genetic connectivity (Froukh and Kochzius, 2007;
10 Nanninga et al., 2014; Giles et al., 2015; Reimer et al. in rev.; Robitzsch et al., 2015).
11 However, very few studies have examined variation in life history traits or demographics
12 (e.g., growth rates or PLD). In our study, we specifically look at the influence of SST and
13 CHLA levels on early life stages of three *Dascyllus* species and find both to be
14 consistently negatively correlated to PLDs of the studied fishes. A reduction in PLDs can
15 be the result of an increase in the metabolic rate of larvae in warmer waters (Atkinson et
16 al., 1996; Björnsson et al., 2001; Green and Fisher, 2004; Hunt et al., 1996) or of higher
17 availability of food (Sponaugle and Grorud-Colvert, 2006) or of a combination of the
18 two. In this context, CHLA explained up to 50% of the variation in PLD data (e.g., of *D.*
19 *trimaculatus*), suggesting it is a strong environmental driver of PLDs in *Dascyllus*.
20 Nevertheless, the coefficients of correlation (R^2 values) for both SST and CHLA are quite
21 similar, which makes it difficult to infer which parameter has a stronger effect on PLDs,
22 and a higher metabolic rate in warmer waters also demands more energy and higher food
23 consumption (Houde and Zastrow, 1993; Houde, 1989). Since both SST and CHLA co-
24 vary with latitude in the Red Sea, it is not surprising that the strongest significant
25 correlation was a between PLD and latitude. However, latitude itself is unlikely to be
26 influencing the PLD of coral reef fishes (Booth and Parkinson, 2011). We interpret our
27 findings as the synergic effect of greater food availability (higher CHLA concentrations)
28 and higher metabolism (higher SST) on larval growth rates, subsequently reducing PLDs
29 for all species in the warmer and more productive southern regions unique to the Red
30 Sea. Furthermore, the decrease of PLDs along the SST and CHLA gradient seems to be
31 stronger and more significant among individuals of the endemic and ecologically more
32 specialized *D. marginatus* in comparison to *D. aruanus* and *D. trimaculatus*. We thus

1 propose that the influence of small-scale regional changes in the environment might be
2 stronger and positively correlated with the degree of ecological specialization and habitat
3 range of damselfishes.

4 This is best illustrated in our study by the observation of large differences in PLDs
5 among the endemic species, *D. marginatus* at one specific site in the Gulf of Aqaba
6 (GAQ) (a 9 d range of difference in HAQ). These differences in PLDs were the highest
7 among all *Dascyllus* in the Red Sea. If we then focus on the regional seasonal changes of
8 the two environmental parameters SST and CHLA, the range of seasonal changes of CHLA
9 concentrations for the GAQ region is distinctly higher compared to adjacent regions to
10 the south, further suggesting that the GAQ is a variable environment itself. In contrast,
11 the seasonal changes in SST in all other regions are rather homogeneous. In other words,
12 larvae that hatch during the spring/early summer in GAQ have up to six times more food
13 available compared to cohorts hatching during other seasons. The extreme variability in
14 the food available to these cohorts could be responsible for the large range of PLDs
15 within the GAQ region. We therefore hypothesize that if the range of PLD differences
16 within sites is a result of seasonal changes in the pelagic environment, seasonal changes
17 in CHLA (food availability) rather than SST could more likely be the driver of
18 differences in PLDs of *Dascyllus* in the Red Sea. However, the proposed hypothesis
19 warrants further investigation in view of the correlation between temperature and food
20 availability (CHLA) (McCormick, 1994; Lo-Yat, et al. 2011); and the implementation of
21 statistical models to discretely analyze the influence of these two environmental
22 parameters on coral reef fishes. In terms of biological data, recruit collections could be
23 targeted at different times of the year to assess whether longer PLDs correlate with
24 seasons of lower productivity. For the more “homogeneously” productive regions of the
25 Red Sea one could assess if the differences in PLDs are also more homogeneously
26 distributed throughout the year. Unfortunately, we were unable to reliably back-calculate
27 birthdates for our individual samples to determine specific intra-annual patterns.
28 Alternate hypotheses that could explain the large within-region differences in PLDs
29 include seasonal changes in survival strategies and/or competition during the larval stage.
30 For example, there may be a benefit in early settlement if there is seasonal variability in
31 settlement habitat availability and early settlement increases the opportunity to secure

space on the reef. Alternatively, delayed recruitment may confer a benefit due to larger size-at-settlement if post-settlement competition varies seasonally. During seasons that may have less settlement competition, delaying settlement may confer advantages in the form of decreased post-settlement mortality (bigger-is-best hypothesis, Anderson, 1988; Miller et al., 1988).

4.2 Pelagic larval durations (PLDs) of *Dascyllus* and the biogeography of coral reef fishes

It is thought that spatial-temporal variation in the environment can select for or against dispersal (Duputié and Massol, 2013), ultimately shaping specialization, evolutionary processes, and the biogeography of species (Berdahl et al., 2015; Heinz et al., 2009). In theory, endemic species should have reduced dispersal potentially linked to a shorter PLD than widespread species, and biogeographic provinces with steep environmental gradients may thus have higher rates of endemism than more homogeneous environments.

The Red Sea has long been recognized as a hotspot of endemism (Roberts, et al., 1992; DiBattista et al., 2015a,b) and in its sharp environmental gradient (Raitos et al., 2013) the endemic damselfish species *D. marginatus* consistently exhibits shorter PLDs than the two widespread species. Of the two more ecologically versatile species, *D. trimaculatus* shows the longest PLD and the global average for this species is even longer (Luiz et al., 2013, Dataset S1). From a biogeographical point of view, this might indicate a successive increase of average PLD with spatial distribution among the three *Dascyllus* species in our study.

The relationship between dispersal potential and biogeographic distribution of marine species has long been an open discussion and many studies have approached this question using PLD data. However, most of these studies have found either weak, non-linear, or no correlation between distribution ranges and PLDs at different geographic scales and for a wide number of species. There can be several reasons for conflicting results among studies on the relationship between dispersal potential and biogeography of marine species. From our data we find it important to point out the extreme plasticity observed in PLDs of *Dascyllus* in relation to the environment and how such variations can radically

change the outcome of the study. For instance, in our study, *D. marginatus* consistently has a shorter PLD than *D. aruanus* at the same geographic site (i.e., spatially coherent data) and we thus find a positive correlation between distribution range and dispersal potential. However, if our reference PLDs for the two species within the Red Sea were not spatially coherent but we instead use a reference PLD for *D. marginatus* from the northern Red Sea (GAQ; the only measurement available in previous literature) and a reference PLD for *D. aruanus* from the southern Red Sea (SRS), we would conclude the opposite, that is, we would find that the endemic species has a longer PLD than the widespread species. Broad comparisons of PLDs to range sizes may therefore be complicated by large intraspecific variations in PLD among sampling locations (see also Victor, 1991).

We suggest that appropriately designed studies may still have the potential to provide insight into the unresolved relationship of PLD and biogeography. The cautious collection of adequate temporal and spatial data is crucial. A study focused on long-term/evolutionary processes like biogeography in relation to contemporary demographic data like PLDs, would ideally meet the following criteria: A) address closely related species to eliminate variation related to phylogenetic effects, B) sample from standing (adult) populations to capture as much temporal variation as possible, also to ensure that the data represent the ‘effective’ population’s PLD (from adults from different recruitment cohorts), and C) sample with spatial coherency in order to isolate extrinsic environmental variables from intrinsic biological differences. The genus *Dascyllus* may offer an opportunity to address such questions. The genus has an endemic representative almost in every biogeographic province in the Indo-Pacific, which will allow future comparative studies using this genus to further assess the relationship between PLD and biogeography over a wide geographic range. Other genera (e.g., *Chaetodon*, *Thalassoma*, *Eviota*, etc.) may provide similar opportunities for congeneric comparison of endemic vs. widespread species. We further advocate the importance of analyzing PLD data accounting for sources of temporal and spatial variation at small scales.

Much work remains to be done to resolve the many potential drivers of plasticity in the PLDs of larval reef fishes at both ecological and evolutionary timescales. The role of various environmental factors and the inclusion of genetic, biological, and ecological data

are a prime target for future studies, particularly in connection with understanding how ecological specialization of a species interacts with these factors.

ACKNOWLEDGMENTS

We would like to thank fieldwork and logistic support provided by KAUST Coastal and Marine Resources Core Lab. For assistance acquiring samples, we thank A. Gusti, M. Roberts, R. Gatins, A. Kattan, G. Bernardi, M. Erdmann, J.-P. Hobbs. For support with environmental data, we thank D. Raitos and D. Dreano. We also thank two anonymous referees and S. Agrawal for comments and feedback, as well as J. S. Berdahl for formatting the artwork. Financial support came from KAUST baseline research funds and an award from the KAUST Office of Competitive Research Funds (award no. URF/1/1389-01-01) to M.LB).

REFERENCES

- Anderson, J.T., 1988. A review of size dependent survival during pre-recruit stages of fishes in relation to recruitment. *J. Northwest Atl. Fish. Sci.* 8, 55–66.
doi:10.1.1.467.4638
- Atkinson, D., Johnston, I.A., Bennett, A.F., 1996. Ectotherm life-history responses to developmental temperature. *Anim. Temp. Phenotypic Evol. Adapt* (ed. by I.A. Johnston & A.F. Bennett). 183–204. New York: Cambridge University Press.
doi:10.1017/CBO9780511721854.009
- Berdahl, A., Torney, C.J., Schertzer, E., Levin, S. a, 2015. On the evolutionary interplay between dispersal and local adaptation in heterogeneous environments. *Evolution*. 69(6), 1390-1405. doi:10.1111/evo.12664
- Bergenius, M.A.J., McCormick, M.I., Meekan, M.G., Robertson, D.R., 2005. Environmental influences on larval duration, growth and magnitude of settlement of a coral reef fish. *Mar. Biol.* 147(2), 291–300. doi:10.1007/s00227-005-1575-z
- Berumen, M.L., Hoey, A.S., Bass, W.H., Bouwmeester, J., Catania, D., Cochran, J.E.M., Khalil, M.T., Miyake, S., Mughal, M.R., Spaet, J.L.Y., Saenz-Agudelo, P., 2013. The status of coral reef ecology research in the Red Sea. *Coral Reefs* 32(3), 737–748.
doi:10.1007/s00338-013-1055-8
- Bikhtashev, V.N., Brindley, J., Horwood, J.W., 2003. Phytoplankton blooms and fish

1 recruitment rate: Effects of spatial distribution. J. Plankton Res. 25(1), 21–33.
2 doi:10.1016/j.bulm.2003.08.008

3 Björnsson, B., Steinarsson, a, Oddgeirsson, M., 2001. Optimal temperature for
4 growth and feed conversion of immature cod (*Gadus morhua* L.). ICES J. Mar.
5 Sci. 58(1), 29–38. doi:10.1006/jmsc.2000.0986

6 Booth, D.J., Parkinson, K., 2011. Pelagic larval duration is similar across 23° of
7 latitude for two species of butterflyfish (Chaetodontidae) in eastern Australia.
8 Coral Reefs 30, 1071–1075. doi:10.1007/s00338-011-0815-6

9 Cowen, R.K., Sponaugle, S., 1997. Relationships between early life history traits and
10 recruitment among coral reef fishes, in: Chambers, C.R., Trippel, E.A. (Eds.),
11 Early Life History and Recruitment in Fish Populations. Chapman and Hall, pp.
12 423–450. doi:10.1007/978-94-009-1439-1_15

13 Cushing, D.H., Horwood, J.W., 1994. The growth and death of fish larvae. J. Plankton
14 Res. 16(3), 291–300. doi:10.1093/plankt/16.3.291

15 DiBattista J.D., Roberts M., Bouwmeester J., Bowen B.W., Coker D., Lozano-Cortés
16 D.F., Choat J.H., Gaither M.R., Hobbs J.P., Khalil M.T., Kochzius M., Myers R.F.,
17 Paulay G., Robitzch V., Rocha L.A., Saenz-Agudelo P., Salas E., Sinclair-Taylor
18 T.H., Toonen R.J., Westneat M., Williams S.T., Berumen M.L., 2015a. A review of
19 contemporary patterns of endemism for shallow water reef fauna in the Red Sea. J.
20 Biogeogr. doi:10.1111/jbi.12649

21 DiBattista J.D., Howard Choat, J., Gaither, M. R., Hobbs, J. P. A., Lozano-Cortés, D. F.,
22 Myers, R. F., Paulay G., Rocha L.A., Toonen R.J., Westneat M., Berumen, M. L.,
23 2015b. On the origin of endemic species in the Red Sea. J. Biogeogr.
24 doi:10.1111/jbi.12631

25 Duputié, A., Massol, F., 2013. An empiricist's guide to theoretical predictions on the
26 evolution of dispersal. Interface Focus 3, 20130028.
27 doi:10.1098/rsfs.2013.0028

28 Fontes, J., Afonso, P., Santos, R.S., Caselle, J.E., 2010. Temporal variability of larval
29 growth, size, stage duration and recruitment of a wrasse, *Coris julis* (Pisces:
30 Labridae), from the Azores. Sci. Mar. 74(4), 721–729.
31 doi:10.3989/scimar.2010.74n4721

32 Froukh, T.J., Kochzius, M., 2007. Genetic population structure of the endemic
33 fourline wrasse (*Larabicus quadrilineatus*) suggests limited larval dispersal
34 distances in the Red Sea. Mol. Ecol. 16(7), 1359–1367. doi:10.1111/j.1365-
35 294X.2007.03236.x

- 1 Giles, E.C., Saenz-Agudelo, P., Hussey, N.E., Ravasi, T., Berumen, M.L., 2015. Exploring
2 seascape genetics and kinship in the reef sponge *Stylissa carteri* in the Red Sea.
3 Ecol. Evol. 5(13), 2487–2502. doi:10.1002/ece3.1511
- 4 Green, B.S., Fisher, R., 2004. Temperature influences swimming speed, growth and
5 larval duration in coral reef fish larvae. J. Exp. Mar. Bio. Ecol. 299(1), 115–132.
6 doi:10.1016/j.jembe.2003.09.001
- 7 Grorud-Colvert, K., Sponaugle, S., 2011. Variability in water temperature affects
8 trait-mediated survival of a newly settled coral reef fish. Oecologia 165(3),
9 675–686. doi:10.1007/s00442-010-1748-4
- 10 Håkanson, L., Bryhn, A.C., Hytteborn, J.K., 2007. On the issue of limiting nutrient and
11 predictions of cyanobacteria in aquatic systems. Sci. Total Environ. 379(1), 89–
12 108. doi:10.1016/j.scitotenv.2007.03.009
- 13 Heath, M.R., 1992. Field investigations of the early life stages of marine fish. Adv.
14 Mar. Biol. 28, 1–174. doi:10.1016/S0065-2881(08)60039-5
- 15 Heinz, S.K., Mazzucco, R., Dieckmann, U., 2009. Speciation and the evolution of
16 dispersal along environmental gradients. Evol. Ecol. 23(1), 53–70.
17 doi:10.1007/s10682-008-9251-7
- 18 Houde, E.D., 1989. Comparative growth, mortality, and energetics of marine fish
19 larvae: temperature and implied latitudinal effects. Fish. Bull. 87(3), 471–495.
20 doi:10.1.1.475.1985
- 21 Houde, E.D., Hoyt, R.D., 1987. Fish early life dynamics and recruitment variability, in:
22 Am. Fish. Soc. Symp. 2, 17-29.
- 23 Houde, E.D., Zastrow, C.E., 1993. Ecosystem- and taxon-specific dynamic and
24 energetics properties of larval fish assemblages. Bull. Mar. Sci. 53(2), 290–335.
- 25 Hunt, I., Boutilier, R.G., Miyake, T., Hall, B.K., 1996. Effects of temperature on
26 morphological landmarks critical to growth and survival in larval Atlantic cod
27 (*Gadus morhua*). Mar. Biol. 124(4), 593–606. doi:10.1007/BF00351041
- 28 Landaeta, M.F., Castro, L.R., 2006. Larval distribution and growth of the rockfish,
29 *Sebastes capensis* (Sebastidae, Pisces), in the fjords of southern Chile. ICES J.
30 Mar. Sci. 63(4), 714–724. doi:10.1016/j.icesjms.2006.01.002
- 31 Lester, S.E., Ruttenberg, B.I., 2005. The relationship between pelagic larval duration
32 and range size in tropical reef fishes: a synthetic analysis. Proc. Biol. Sci. 272,
33 585–591. doi:10.1098/rspb.2004.2985

- 1 Lo-Yat, A., Simpson, S.D., Meekan, M., Lecchini, D., Martinez, E., Galzin, R., 2011.
2 Extreme climatic events reduce ocean productivity and larval supply in a
3 tropical reef ecosystem. *Glob. Chang. Biol.* 17, 1695–1702. doi:10.1111/j.1365-
4 2486.2010.02355.x
- 5 Lozano-Cortés, D. F., Berumen, M. L., 2015. Colony size-frequency distribution of
6 pocilloporid juvenile corals along a natural environmental gradient in the Red
7 Sea. *Mar. Poll. Bull.* doi:10.1016/j.marpolbul.2015.10.051
- 8 Luiz, O.J., Allen, A.P., Robertson, D.R., Floeter, S.R., Kulbicki, M., Vigliola, L., Becheler,
9 R., Madin, J.S., 2013. Adult and larval traits as determinants of geographic range
10 size among tropical reef fishes. *PNAS* 110(41), 16498–16502.
11 doi:10.1073/pnas.1304074110
- 12 Macpherson, E., Raventos, N., 2006. Relationship between pelagic larval duration
13 and geographic distribution of Mediterranean littoral fishes. *Mar. Ecol. Prog.*
14 *Ser.* 327, 257–265. doi:10.3354/meps327257
- 15 McCormick, M., 1994. Variability in age and size at settlement of the tropical goatfish
16 *Upeneus tragula* (Mullidae) in the northern Great Barrier Reef lagoon. *Mar. Ecol.*
17 *Prog. Ser.* 103, 1–15. doi:10.3354/meps103001
- 18 McCormick, M., Molony, B., 1995. Influence of water temperature during the larval
19 stage on size, age and body condition of a tropical reef fish at settlement. *Mar.*
20 *Ecol. Prog. Ser.* 118, 59–68. doi:10.3354/meps118059
- 21 Mcleod, I.M., McCormick, M.I., Brooker, R.M., 2015. Latitudinal variation in larval
22 development of coral reef fishes: implications of a warming ocean. *Mar. Ecol.*
23 *Prog. Ser.* 521, 129–141. doi:10.3354/meps11136
- 24 Mcleod, I.M., Rummer, J.L., Clark, T.D., Jones, G.P., McCormick, M.I., Wenger, A.S.,
25 Munday, P.L., 2011. Climate change and the performance of larval coral reef
26 fishes: the interaction between temperature and food availability *Conserv.*
27 *Physiol.* 1, 1–12. doi:10.1093/conphys/cot024
- 28 Meekan, M.G., Carleton, J.H., McKinnon, A.D., Flynn, K., Furnas, M., 2003. What
29 determines the growth of tropical reef fish larvae in the plankton: food or
30 temperature? *Mar. Ecol. Prog. Ser.* 256, 193–204. doi:10.3354/meps256193
- 31 Miller, T.J., Crowder, L.B., Rice, J.A., Marschall, E.A., 1988. Larval size and recruitment
32 mechanisms in fishes: Toward a conceptual framework. *Can. J. Fish. Aquat. Sci.*
33 45(9), 1657–1670. doi:10.1139/f88-197
- 34 Mora, C., Chittaro, P.M., Sale, P.F., Kritzer, J.P., Ludsins, S. a., 2003. Patterns and

1 processes in reef fish diversity. *Nature* 421, 933–936.
2 doi:10.1038/nature01421.1.

3 Nanninga, G.B., Saenz-Agudelo, P., Manica, A., Berumen, M.L., 2014. Environmental
4 gradients predict the genetic population structure of a coral reef fish in the Red
5 Sea. *Mol. Ecol.* 23(3), 591–602. doi:10.1111/mec.12623

6 Ngugi, D.K., Antunes, A., Brune, A., Stingl, U., 2012. Biogeography of pelagic
7 bacterioplankton across an antagonistic temperature-salinity gradient in the
8 Red Sea. *Mol. Ecol.* 21(2), 388–405. doi:10.1111/j.1365-294X.2011.05378.x

9 Racault, M.-F., Raitsos, D.E., Berumen, M.L., Brewin, R.J.W., Platt, T., Sathyendranath,
10 S., Hoteit, I., 2015. Phytoplankton phenology indices in coral reef ecosystems:
11 Application to ocean-color observations in the Red Sea. *Remote Sens. Environ.*
12 160, 222–234. doi:10.1016/j.rse.2015.01.019

13 Raitsos, D.E., Pradhan, Y., Brewin, R.J.W., Stenchikov, G., Hoteit, I., 2013. Remote
14 sensing the phytoplankton seasonal succession of the Red Sea. *PLoS One* 8(6),
15 e64909. doi:10.1371/journal.pone.0064909

16 Rankin, T.L., Sponaugle, S., 2011. Temperature influences selective mortality during
17 the early life stages of a coral reef fish. *PLoS One* 6(5), e16814.
18 doi:10.1371/journal.pone.0016814

19 Reimer J., Herrera M., Gatins R., Roberts M., Berumen M.L., In Review. Latitudinal
20 variation in the symbiotic dinoflagellate *Symbiodinium* of the common reef
21 zoantharian *Palythoa tuberculosa* on the Saudi Arabian coast of the Red Sea. *J.*
22 *Biogeogr.*

23 Roberts, C., Shepherd, A., Ormond, R., 1992. Large-scale variation in assemblage
24 structure of Red Sea butterflyfishes and angelfishes. *J. Biogeogr.* 19(3), 239–
25 250. doi:10.2307/2845449

26 Robitzch, V., Banguera-Hinestroza, E., Sawall, Y., Al-Sofyani, A., Voolstra, C.R., 2015.
27 Absence of genetic differentiation in the coral *Pocillopora verrucosa* along
28 environmental gradients of the Saudi Arabian Red Sea. *Front. Mar. Sci.* 2(5), 1–
29 10. doi:10.3389/fmars.2015.00005

30 Sawall, Y., Al-Sofyani, A., Banguera-hinestroza, E., Voolstra, C.R., 2014a. Spatio-
31 temporal analyses of *Symbiodinium* physiology of the coral *Pocillopora*
32 *verrucosa* along large-scale nutrient and temperature gradients in the Red Sea.
33 *PLoS One*, e103179. doi:10.1371/journal.pone.0103179

34 Sawall, Y., Al-Sofyani, A., Kuerten, B., Al-Aidaros, A.M., Hoang, B.X., Marimuthu, N.,

- 1 Khomayis, H.S., Sommer, U., Gharbawi, W.Y., Wahl, M., 2014b. Coral
2 communities, in contrast to fish communities, maintain a high assembly
3 similarity along the large latitudinal gradient along the Saudi Red Sea coast. J.
4 Ecosyst. Ecography 3(4), 1–7. doi:10.4172/2157-7625.S4-003
- 5 Searcy, S.P., Sponaugle, S., 2000. Variable larval growth in a coral reef fish. Mar. Ecol.
6 Ser. 206, 213–226. doi:10.3354/meps206213
- 7 Sponaugle, S., 2010. Otolith microstructure reveals ecological and oceanographic
8 processes important to ecosystem-based management. Environ. Biol. Fishes
9 89(3), 221–238. doi:10.1007/s10641-010-9676-z
- 10 Sponaugle, S., Boulay, J., Rankin, T., 2011. Growth- and size-selective mortality in
11 pelagic larvae of a common reef fish. Aquat. Biol. 13, 263–273.
12 doi:10.3354/ab00370
- 13 Sponaugle, S., Grorud-Colvert, K., 2006. Environmental variability, early life-history
14 traits, and survival of new coral reef fish recruits. Integr. Comp. Biol. 46(5),
15 623–633. doi:10.1093/icb/icl014
- 16 Sponaugle, S., Grorud-Colvert, K., Pinkard, D., 2006. Temperature-mediated
17 variation in early life history traits and recruitment success of the coral reef fish
18 *Thalassoma bifasciatum* in the Florida Keys. Mar. Ecol. Prog. Ser. 308, 1–15.
19 doi:10.3354/meps308001
- 20 Sponaugle, S.U., 2009. Tropical fish otoliths: Information for assessment,
21 management and ecology, in: Green, B.S., Mapstone, B.D., Carlos, G., Begg, G.A.
22 (Eds.), Methods and Technologies in Fish Biology and Fisheries, Reviews:
23 Methods and Technologies in Fish Biology and Fisheries. Springer Netherlands,
24 Dordrecht, pp. 93–132. doi:10.1007/978-1-4020-5775-5
- 25 Takahashi, M., Watanabe, Y., 2005. Effects of temperature and food availability on
26 growth rate during late larval stage of Japanese anchovy (*Engraulis japonicus*)
27 in the Kuroshio-Oyashio transition region. Fish. Oceanogr. 14(3), 223–235.
28 doi:10.1111/j.1365-2419.2005.00334.x
- 29 Thresher, R.E., Colin, P.L., Bell, L.J., 1989. Planktonic duration, distribution and
30 population structure of western and central pacific damselfishes
31 (Pomacentridae). Copeia 1989(2), 420–434. doi:10.2307/1445439
- 32 Victor, B.C., 1991. Settlement strategies and biogeography of reef fishes, in: Sale, P.F.
33 (Ed.), The Ecology of Fishes on Coral Reefs. Academic Press, Inc., San Diego,
34 California, pp. 231–260. doi:10.1016/b978-0-08-092551-6.50014-3

- 1 Victor, B.C., 1986. Duration of the planktonic larval stage of one hundred species of
2 Pacific and Atlantic wrasses (family Labridae). Mar. Biol. 90(30), 317–326.
3 doi:10.1007/BF00428555
- 4 Victor, B.C., Wellington, G.M., 2000. Endemism and the pelagic larval duration of reef
5 fishes in the eastern Pacific Ocean. Mar. Ecol. Prog. Ser. 205, 241–248.
6 doi:10.1098/rspb.2004.2985
- 7 Wellington, G.M., Victor, B.C., 1992. Regional differences in duration of the
8 planktonic larval stage of reef fishes in the eastern Pacific Ocean. Mar. Biol.
9 113(3), 491–498. doi:10.1007/BF00349176
- 10 Wellington, G.M., Victor, B.C., 1989. Planktonic larval duration of one hundred
11 species of Pacific and Atlantic damselfishes (Pomacentridae).
12 doi:10.1007/BF00541659 Mar. Biol. 101(4), 557–567.
- 13 Wilson, D.T., McCormick, M.I., 1997. Spatial and temporal validation of settlement-
14 marks in the otoliths of tropical reef fishes. Mar. Ecol. Prog. Ser. 153, 259–271.
15 doi:10.3354/meps153259
- 16 Wilson, D.T., Meekan, M.G., 2002. Growth-related advantages for survival to the
17 point of replenishment in the coral reef fish *Stegastes partitus* (Pomacentridae).
18 Mar. Ecol. Prog. Ser. 231, 247–260. doi:10.3354/meps231247
- 19 Zapata, F.A., Herrón, P.A., 2002. Pelagic larval duration and geographic distribution
20 of tropical eastern Pacific snappers (Pisces: Lutjanidae). Mar. Ecol. Prog. Ser.
21 230, 295–300. doi:10.1098/rspb.2004.2985