Exploring the genetic diversity of shallow-water Agariciidae (Cnidaria: Anthozoa) from the Saudi Arabian Red Sea

Tullia I. Terraneo¹ · Roberto Arrigoni¹ · Francesca Benzoni^{2,3} · Matthew D. Tietbohl¹ · Michael L. Berumen¹

Received: 1 October 2016 / Revised: 26 April 2017 / Accepted: 26 April 2017
© Senckenberg Gesellschaft für Naturforschung and Springer-Verlag Berlin Heidelberg 2017

Abstract Scleractinian corals ascribed to the family Agariciidae represent an important component of Red Sea coral reef fauna, though little genetic data are currently available for this group, and existing information shows polyphyly in the examined mesophotic taxa from the Pacific Ocean. In this work, we provide a first genetic survey of Agariciidae from the Saudi Arabian Red Sea, based on a collection of shallow-water material (<30 m) from the Gulf of Aqaba to the Farasan Islands. Two molecular markers were sequenced to infer morphospecies monophyly and relationships, the intergenic region between COI and 16S rRNA from mitochondrial DNA and the ribosomal ITS1 region from nuclear DNA. A total of 20 morphospecies were identified based on classical macromorphological characters. Six, namely *Gardineroseris planulata*, *Pavona maldivensis*, *Pavona clavus*, *Pavona decussata*, *Leptoseris fragilis*, and *Leptoseris yabei*, were resolved with both DNA loci. The molecular boundaries among the remaining 14 species remain unclear. Our results further confirm that the morphology-based taxonomy of most agariciid species is in disagreement with

genetics. In order to disentangle the systematics of these taxa, the inclusion of more sampling locations, additional variable loci, and a micromorphological approach are likely needed. Our genetic data represent a first step towards the comparison of biodiversity and connectivity between the Red Sea and the rest of the Indo-Pacific.

Keywords Coral · Systematics · Biodiversity · IGR · ITS · Phylogenetics · DNA taxonomy

Introduction

The Red Sea (RS) is a north–south orientated and semi-enclosed basin at the northwestern corner of the Indian Ocean (IO). It is connected to the IO only at the narrow (18 km) and shallow (137 m) Strait of Bab al Mandab (Sheppard et al. 1992). Despite its modest surface area (438,000 km²), the total coral reef area of the RS (c. 8890 km²) is similar to that of the Caribbean (c. 10,530 km²) and its reef system length (c. 2000 km) is comparable to that of the Great Barrier Reef (GBR) (c. 2300 km) (Berumen et al. 2013). The RS has long been recognized as a region of high marine biodiversity and endemism (Klunzinger 1870, 1877; Ekman 1953). The unique reef fauna of the RS has multiple evolutionary origins (DiBattista et al. 2016a), with some taxa originating within the RS during the Pleistocene glaciations (DiBattista et al. 2015; Coleman et al. 2016) and others diverging from their IO sister groups long before the last glacial cycles (DiBattista et al. 2013; Hodge et al. 2014).

In the RS, 5.5% of the 364 scleractinian coral species are endemic (DiBattista et al. 2016b), whereas levels of endemism for scleractinians is typically less than 2% in the IO (Obura 2012; Veron et al. 2015). However, our knowledge of the coral ecology and biodiversity of the RS is limited compared to

Tullia I. Terraneo and Roberto Arrigoni contributed equally to this work.

Communicated by M. Sonnewald

✉ Tullia I. Terraneo
tulliaisotta.terraneo@kaust.edu.sa

¹ Red Sea Research Center, Division of Biological and Environmental Science and Engineering, King Abdullah University of Science and Technology, Thuwal 23955-6900, Saudi Arabia

² UMR ENTROPIE (IRD, Université de La Réunion, CNRS), Laboratoire d'Excellence–CORAIL, Centre IRD de Nouméa, New Caledonia, 101 Promenade Roger Laroque, BP A5, 98848 Noumea Cedex, New Caledonia

³ Department of Biotechnologies and Biosciences, University of Milano—Bicocca, Piazza della Scienza 2, 20126 Milan, Italy

other major reef systems, e.g., the GBR and the Caribbean (Berumen et al. 2013). While the Gulf of Aqaba (GofA) has been the source of notable and seminal works in the field of coral reef ecology and taxonomy (Loya et al. 2014), the restricted area of the GofA (<2% of the RS) may preclude extrapolation to the broader RS. RS corals attracted the attention of early taxonomists (e.g., Forskål 1775; Lamarck 1816; Ehrenberg 1834; Klunzinger 1877) and were later the subject of more detailed surveys and larger reference collections (e.g., Scheer and Pillai 1983; Sheppard and Sheppard 1991). These publications, representing milestones for coral taxonomy in the RS, are based on classical morphology-based taxonomy, providing detailed descriptions of coralla and corallite features at the macroscopic scale. Recently, molecular works coupled with analyses of the macro- and micromorphology of the skeletons have allowed the description of new reef corals in the RS (Terraneo et al. 2014; Arrigoni et al. 2015, 2016a, b, 2017) and have showed unexpectedly low genetic diversity for some genera previously thought to have high species richness in this region (Terraneo et al. 2016; Arrigoni et al. 2016c). Moreover, other genetic surveys have revealed several cases of genetic differentiation between Indian and Pacific coral populations previously obscured by traditional taxonomy (Stefani et al. 2011; Ladner and Palumbi 2012; Arrigoni et al. 2012; Kitano et al. 2014; Richards et al. 2016).

To date, the scleractinian coral family Agariciidae Gray, 1847 comprises seven genera (Kitahara et al. 2012) and 49 extant species (WoRMS 2016). The genus *Pachyseris* Milne Edwards & Haime, 1849 was also ascribed to the family (Vaughan and Wells 1943; Wells 1956; Veron and Pichon 1980). Nevertheless, recent molecular works provide evidence that *Pachyseris* is not closely related to the Agariciidae and, instead, is basal to the family Euphylliidae Alloiteau, 1952 (Fukami et al. 2008; Kitahara et al. 2010); the position of the genus remains uncertain. Within the family, the genera *Coeloseris* Vaughan, 1918, *Dactylotrachus* Wells, 1954, *Gardineroseris* Scheer & Pillai, 1974, and *Pavona* Lamarck, 1801 occur in the Indo-Pacific; *Agaricia* Lamarck, 1801 and *Helioseris* Milne Edwards & Haime, 1849 occur in the Western Atlantic; and *Leptoseris* Milne Edwards & Haime, 1849 occurs in both the Indo-Pacific and the Atlantic Ocean. Agariciids are common in tropical shallow-water coral reefs (<30 m) throughout their distribution range (Veron 2000; Waheed and Hoeksema 2014; Waheed et al. 2015). Moreover, representatives of *Leptoseris* and *Agaricia* may be a dominant component of mesophotic coral ecosystems (Fricke et al. 1987; Hinderstein et al. 2010; Rooney et al. 2010; Kahng et al. 2014) occurring in the tropics and subtropics between depths of 30 m to over 150 m (Kahng et al. 2010). For example, *Leptoseris fragilis* Milne Edwards & Haime, 1849 has been found at 145 m in the GofA (Fricke and Knauer 1986) and *Leptoseris hawaiiensis* Vaughan, 1907 at 165 m from Johnston Atoll (Maragos and Jokiel 1986). In the RS, species

of *Gardineroseris* and *Pavona* are commonly found between 1 and 20 m depth, while most species of *Leptoseris* are more abundant in low-light conditions under reef overhangs or below 30 m, where they can be a major component of coral coverage (Scheer and Pillai 1983; Sheppard and Sheppard 1991). Notably, *Leptoseris* has been recorded from over 100 m depth (Fricke et al. 1987). Scheer and Pillai (1983) listed 15 species of Agariciidae from the Saudi Arabian RS, Sheppard and Sheppard (1991) listed 16 species from the Saudi Arabian RS, and Veron (2000) listed 19 species from the entire RS (see Table 1 for the complete list of species reported by these three publications). Subsequently, *Leptoseris amitoriensis* Veron, 1990 was reported for the first time in the RS by Ziegler et al. (2015).

Despite the ecological importance of this family, its taxonomy and systematics are still almost entirely based on traditional studies of skeletal morphology (Dinesen 1980; Veron and Pichon 1980; Veron 2000). The definition of agariciid genera and species boundaries occurred prior to the introduction of molecular techniques that have revolutionized coral systematics at all taxonomic ranks (Kitahara et al. 2016). A detailed morpho-molecular approach by Kitahara et al. (2012) demonstrated that the solitary and deep-water species *Dactylotrachus cervicornis* (Moseley, 1881), previously ascribed to the Caryophylliidae Dana, 1846, is the oldest extant representative of Agariciidae. Phylogenetic investigations of the Hawaiian mesophotic *Leptoseris* and *Pavona* species based on the mitochondrial intergenic region between COI and 16S rRNA (IGR hereafter) showed that the two genera are not monophyletic and revealed several hidden species (Luck et al. 2013; Pochon et al. 2015). Other integrative works focusing on species delimitation among shallow-water species of *Pavona* found genetic distinction between *Pavona chiriquiensis* Glynn, Mate & Stemmann, 2001 and *Pavona varians* Verrill, 1864 from Panama (Maté 2003), between *Pavona cactus* Forskål, 1775 and *Pavona decussata* Dana, 1846 from Mauritius (Moothien Pillay et al. 2006), and between *Pavona maldivensis* (Gardiner, 1905) and *Pavona explanulata* (Lamarck, 1816) from Malaysia (Waheed et al. 2015). However, the data produced in the above studies are only partially comparable because they are based on different molecular techniques or distinct loci with variable resolutions; they also typically represent single cases and do not address the whole spectrum of diversity within the family.

In this work, we present the first survey of the genetic diversity of shallow-water (<30 m) Agariciidae from the Saudi Arabian RS for the morphospecies encountered during different biodiversity surveys organized by the King Abdullah University of Science and Technology (KAUST). The skeletal macromorphology of the collected material was used to identify the specimens based on existing taxonomic references and, when possible, through comparison with type material. Two molecular markers, i.e., mitochondrial IGR and internal

Table 1 List of Agariciidae species recorded from the Red Sea (RS) by Scheer and Pillai (1983), Sheppard and Sheppard (1991), Veron (2000), and this study

	Scheer and Pillai (1983)	Sheppard and Sheppard (1991)	Veron (2000)	This study
<i>Gardineroseris planulata</i>	x	x	x	x
<i>Pavona cactus</i>	x	x	x	x
<i>Pavona clavus</i>			x	x
<i>Pavona danai</i>			x	x
<i>Pavona decussata</i>	x	x	x	x
<i>Pavona diffluens</i>		x	x	x
<i>Pavona divaricata</i>	x			
<i>Pavona duerdeni</i>		x	x	
<i>Pavona explanulata</i>	x	x	x	x
<i>Pavona frondifera</i>			x	x
<i>Pavona maldivensis</i>	x	x	x	x
<i>Pavona minuta</i>		x		
<i>Pavona varians</i>	x	x	x	x
<i>Pavona venosa</i>		x	x	x
<i>Leptoseris explanata</i>	x	x	x	x
<i>Leptoseris foliosa</i>		x	x	
<i>Leptoseris fragilis</i>	x			x
<i>Leptoseris gardineri</i>	x			
<i>Leptoseris glabra</i>				x
<i>Leptoseris hawaiiensis</i>	x	x	x	x*
<i>Leptoseris incrustans</i>			x	x
<i>Leptoseris mycetoseroides</i>		x	x	x
<i>Leptoseris scabra</i>	x	x	x	x
<i>Leptoseris tenuis</i>	x			
<i>Leptoseris yabei</i>	x	x	x	x
Total	14	16	19	18

*Recorded as *Leptoseris* cf. *hawaiiensis*

transcribed spacer 1 from nuclear ribosomal DNA (ITS1), were sequenced to define morphospecies monophyly and relationships. This study represents the first molecular phylogeny of Agariciidae from the RS, and in general from the IO, and will allow the comparison of agariciid genetic diversity and connectivity of the RS to other localities of the Indo-Pacific. The results highlight a disagreement between our identifications based on classical morphologic characters and DNA for most species, suggesting the need to include more locations, variable loci, and the integration of micromorphological data to disentangle the systematics of this group.

Materials and methods

Sampling

Between 2012 and 2016, 91 samples belonging to the family Agariciidae were collected from several localities along the

Saudi Arabian RS coast, from the GofA to the Farasan Islands (Table 1). Prior to collection, living coral colonies were photographed in the field using either a Canon G9 or G11 PowerShot digital camera (Canon Inc., Tokyo, Japan) in an Ikelite underwater housing (Ikelite Underwater Systems, Indianapolis, IN, USA). Specimens were sampled during SCUBA diving between 1 and 30 m depth using hammer and chisel. A small fragment (<1 cm²) of each coral colony was preserved in 96% ethanol for molecular analyses. The rest of the corallum was tagged and bleached in sodium hypochlorite for 48 h in order to remove tissue parts, rinsed with fresh water, and air-dried. Images of cleaned skeletons were taken with a Canon G9 or G11 PowerShot digital camera.

Species identification

Specimens were identified to the species level based on macromorphological features of the coralla and the corallites. For the genus *Leptoseris*, we referred to the revision by

Dinesen (1980), including descriptions and illustrations of type material. For *Leptoseris yabei* (Pillai & Scheer, 1976), originally described in *Pavona*, we referred to the original description and holotype illustration (pl. 16, figs. 1, 2) by Pillai and Scheer (1976). For the other genera, Veron and Pichon (1980), Scheer and Pillai (1983), Sheppard and Sheppard (1991), Nishihira and Veron (1995), and Veron (2000) were referenced. The type material of Agariciidae species deposited at the National Museum of Natural History (MNHN), Paris, France and at the Smithsonian Institution, National Museum of Natural History (formerly known as United States National Museum of Natural History) (USNM), Washington, USA, was also examined by FB and imaged. These images were used as reference for the identification of a number of species, namely *Leptoseris fragilis* Milne Edwards & Haime, 1849 (MNHN 468), *Leptoseris scabra* Vaughan, 1907 (USNM 20885; USNM 20886), *Leptoseris hawaiiensis* Vaughan, 1907 (USNM 20874; USNM 20876), *Leptoseris mycetoseroides* Wells, 1954 (USNM 44805; USNM 44807), *Pavona diffluens* (Lamarck, 816) (MNHN IK-2010-587), *Pavona danai* (USNM 136), *Pavona duerdeni* Vaughan, 1907 (USNM 21630; USNM 21631), and *P. decussata* (USNM 176).

DNA extraction, amplification, and sequence analyses

Genomic DNA was extracted from samples using the commercial DNAeasy® Blood & Tissue Kit (Qiagen Inc., Hilden, Germany), following the manufacturer's protocol. Primers AGAH and AGAL (Terraneo et al. 2014) were used to amplify IGR locus, while 1S and 2SS primers (Wei et al. 2006) were used to amplify ITS1. All of the polymerase chain reaction (PCR) products were purified with Illustra ExoStar (GE Healthcare, Buckinghamshire, UK) at 37° for 60 min, followed by 85° for 15 min, and sent for direct sequencing in both forward and reverse directions using an ABI 3730xl DNA analyzer (Applied Biosystems, Carlsbad, CA, USA). All the sequences generated from this work were deposited in EMBL (Table 1).

Forward and reverse sequences were viewed, edited, and assembled using Sequencher 5.3 (Gene Codes Corp., Ann Arbor, MI, USA). Nuclear sequences were phased using SeqPHASE (Flot 2010; available online at <http://seqphase.mpg.de/seqphase/>) and PHASE was used (Stephens et al. 2001; available online at <http://stephenslab.uchicago.edu/software.html>) when alleles showed the same length; Champuru (Flot 2007; available online at <http://seqphase.mpg.de/champuru/>) was used if the two predominant alleles were of different lengths. In the former case, the two alleles with the highest probability (an order of magnitude greater than the other sequence pairs) were chosen whenever there were multiple possible phases. No obvious or significant differences of genetic diversity and haploweb inference were obtained using alternative phases (results not shown). Phased heterozygotes were represented by both alleles in the further alignments and

population genetic analyses. Final alignments were performed using the E-INS-i option in MAFFT 7.130b (Kato and Standley 2013) under default parameters and manually checked using BioEdit 7.2.5 (Hall 1999). Ambiguously aligned regions were removed from the alignments using Gblocks 0.91b (Castresana 2000) using the “more stringent” selection (alignment data are available from the corresponding author upon request). Invariable, polymorphic, and parsimony-informative sites were detected using DnaSP 5.10.01 (Librado and Rozas 2009).

The two loci were analyzed separately, and phylogenetic relationships between species were assessed using Bayesian inference (BI), maximum likelihood (ML), and maximum parsimony (MP), as implemented in MrBayes 3.1.2 (Ronquist and Huelsenbeck 2003), PhyML 3.0 (Guindon and Gascuel 2003), and PAUP* 4.0b10 (Swofford 2002), respectively. We used the Akaike information criterion (AIC) implemented in MrModeltest 2.3 (Posada and Crandall 1998) in conjunction with PAUP* 4.0b10 to determine the best-fit substitution model of sequence evolution. The AIC selected the General Time Reversible (GTR) + invariable sites + gamma model of nucleotide substitution for both markers ($P_{inv} = 0.28$, $\alpha = 0.51$ for IGR, and $P_{inv} = 0.78$, $\alpha = 0.55$ for ITS1). The BI analysis consisted of four independent Markov chain Monte Carlo (MCMC) runs for 6,000,000 generations for IGR (10,000,000 for ITS1), saving a tree every 1000 generations and discarding the first 25% of the trees as burn-in, on the basis of parameter estimations and convergence examined by Tracer 1.6 (Rambaut and Drummond 2007). Clade support was based on posterior probability. The ML analysis was run using the default parameters, the previously selected model, and 1000 bootstrap replicates to verify the robustness of the internal branches of the tree. For MP, a heuristic search was performed using starting trees obtained by random stepwise addition with 10 replicates and the tree bisection and reconnection (TBR) branch-swapping-algorithm, generating a strict consensus tree. The robustness of the internal branches of the tree was verified using 1000 bootstrap replicates. The two alignments and phylogenetic trees are available in TreeBASE (<http://purl.org/phylo/treebase/phyloids/study/TB2:S20870>).

Results

The 91 Agariciidae colonies collected in the Saudi Arabian RS were identified based on colony and corallite macromorphology, amounting to a total of 20 species ascribed to three genera, namely *Pavona*, *Leptoseris*, and *Gardineroseris* (Table 1) (Figs. 1 and 2). One *Leptoseris* specimen was identified as *Leptoseris* cf. *hawaiiensis*. The uncertainty about the identification of this distinct specimen stems from the substantial morphological differences between the actual type material and

Fig. 1 In vivo colony morphology of the Agariciidae collected for this study in the Saudi Arabian Red Sea (RS): **a** SA0076 *Pavona venosa*; **b** SA0857 *Pavona varians*; **c** SA0368 *Pavona maldivensis*; **d** SA0105 *Leptoseris fragilis*; **e** SA0269 *Pavona decussata*; **f** SA0810 *Gardineroseris planulata*; **g** SA0369 *Pavona clavus*; **h** SA0943 *Leptoseris* sp.; **i** SA0942 *Pavona explanulata*; **j** SA0781 *Leptoseris mycetoseroides*; **k** SA0884 *Leptoseris* cf. *hawaiiensis*; **l** SA2654 *Leptoseris scabra*; **m** SA2608 *Leptoseris incrustans*; **n** SA0097 *Leptoseris explanata*; **o** SA0893 *Pavona diffluens*; **p** SA0088 *Leptoseris glabra*; **q** SA0424 *Leptoseris yabei*; **r** SA0052 *Pavona danae*; **s** SA0205 *Pavona cactus*; **t** SA2653 *Pavona frondifera*

specimens and colonies identified by other authors as *L. hawaiiensis* (Sheppard and Sheppard 1991; Veron 2000).

Phylogenetic analyses

The final mitochondrial alignment consisted of 1081 bp and included 187 polymorphic sites (71 of which were singleton sites) and 116 parsimony-informative sites. Two hundred and fifty-seven mutations were recovered from the dataset when evaluating synonymous and non-synonymous substitutions. The ITS alignment consisted of 468 bp and 54 polymorphic sites (14 of which were 14 singleton sites), 40 parsimony-informative sites, and a total of 49 mutations.

Two sequences of *Siderastrea radians* (Pallas, 1776) were downloaded from GenBank and used as the outgroup for both mitochondrial and nuclear datasets because of the known divergence of this species from Agariciidae (Fukami et al. 2008; Luck et al. 2013; Terraneo et al. 2014; Pochon et al. 2015). Topologies derived from BI, ML, and MP were mostly in agreement, resolving the same well-supported molecular clades without any conflicting partition. In Figs. 3 and 4, we report the Bayesian phylogram with branch support indicated

by Bayesian posterior probability (BB_{PI}), ML bootstrapping support (BT_{ML}), and MP bootstrapping support (BT_{MP}).

The mitochondrial phylogenetic reconstruction demonstrates that genus-level boundaries of *Leptoseris*, *Pavona*, and *Gardineroseris* are not genetically supported (Fig. 3). In fact, *Leptoseris* and *Pavona* as currently defined are intermixed between each other, and the monotypic genus *Gardineroseris* nests together with *Leptoseris* and *Pavona*.

According to the mitochondrial reconstruction, species-level boundaries are also unclear for many of the analyzed taxa. *Gardineroseris planulata* forms a monophyletic group (CLADE V) strongly supported by the three analysis criteria (0.94/100/100). Similarly, *Leptoseris fragilis* (CLADE III), *L. yabei* (CLADE XIII), and *Leptoseris* sp. (CLADE VII) are monophyletic and grouped in highly supported clades (node values 1/100/100, 0.99/–/100, and 1/99/100, respectively). Sequences of *L. mycetoseroides* are split into two distinct clades (CLADE IX and CLADE XI), showing cryptic genetic divergence within the morphospecies. *Leptoseris* cf. *hawaiiensis*, *L. scabra*, *L. incrustans*, *L. explanata*, and *L. glabra* are genetically indistinguishable based on this marker and grouped together in CLADE XII, which also includes all the analyzed samples of *P. diffluens*. *Pavona maldivensis* (CLADE II) forms

Fig. 2 Skeletal macromorphology of the Agariciidae species collected for this study in the Saudi Arabian Red Sea (RS): **a** SA0076 *Pavona venosa*; **b** SA0857 *Pavona varians*; **c** SA0368 *Pavona maldivensis*; **d** SA0105 *Leptoseris fragilis*; **e** SA0269 *Pavona decussata*; **f** SA0810 *Gardineroseris planulata*; **g** SA0369 *Pavona clavus*; **h** SA0943 *Leptoseris* sp.; **i** SA0942 *Pavona explanulata*; **j** SA0781 *Leptoseris mycetoseroides*; **k** SA0884 *Leptoseris* cf. *hawaiiensis*; **l** SA2654 *Leptoseris scabra*; **m** SA2608 *Leptoseris incrustans*; **n** SA0097 *Leptoseris explanata*; **o** SA0893 *Pavona diffluens*; **p** SA0088 *Leptoseris glabra*; **q** SA0424 *Leptoseris yabei*; **r** SA0052 *Pavona danae*; **s** SA0205 *Pavona cactus*; **t** SA2653 *Pavona frondifera*

a monophyletic group strongly supported by the three phylogenetic analyses (0.98/96/84). All representatives of *P. decussata* (CLADE IV) cluster together in a highly supported lineage (1/100/100). The only examined sample of *P. clavus* is highly divergent within the tree (CLADE VI), but one specimen is not sufficient to evaluate its monophyly. Based on mitochondrial sequence data, *P. venosa* and *P. varians* are indistinguishable from a molecular point of view, and are nested together in CLADE I, while *P. explanulata* is comprised of two distinct lineages (CLADE VIII and CLADE X). Finally, the frondose *Pavona danai* Milne Edwards, 1860, *Pavona frondifera* (Lamarck, 1816), and *P. cactus* form a single clade (CLADE XIV) basal to the entire phylogeny (Fig. 3, Table 3).

The phylogenetic tree based on nuclear ITS1 is less resolved than the mitochondrial phylogeny reconstruction, though no contrasting patterns are found between the two analyses, supporting the stability of relationships among species. All agariciids form an unresolved clade due to a large basal polytomy. Similar to the IGR tree topology, *Leptoseris*, *Pavona*, and *Gardineroseris* are closely related and nest together based on ITS1 (Fig. 4), which raises questions regarding the validity of these three genera. Five molecular clades identified by IGR are also resolved in the ITS1 phylogeny, including

P. maldivensis (CLADE II), *G. planulata* (CLADE V), *P. clavus* (CLADE VI), *L. yabei* (CLADE XIII), and *P. danai*, *P. frondifera*, and *P. cactus* (CLADE XIV). *Pavona decussata* (CLADE IV) and *P. varians* and *P. venosa* (CLADE I) group together in a molecular lineage (–/79/–) and, interestingly, representatives of *P. decussata* form a group genetically isolated from the other two species. *Leptoseris fragilis* (CLADE III) and *Leptoseris* sp. (CLADE VII) are unresolved, forming part of the large basal polytomy, but they do not cluster with any other morphospecies. *Leptoseris mycetoseroides* (CLADES IX and XI) is unresolved and representatives of these two clades are mixed together. Similar to the placement of *L. mycetoseroides*, *P. explanulata* (CLADES VIII and X) is also unresolved, though sequences ascribed to these two clades mix together. Lastly, *P. diffluens*, *L. explanata*, *L. glabra*, *L. incrustans*, and *L. scabra* (CLADE XII) are unresolved in the large basal polytomy of the ITS1 phylogenetic tree.

Discussion

In the present study, we provide a preliminary genetic survey of the scleractinian coral family Agariciidae from the Saudi

Fig. 3 Phylogenetic reconstruction of the family Agariciidae from the Saudi Arabian Red Sea (RS) inferred from Bayesian inference (BI) analysis of the mitochondrial intergenic spacer between COI and 16S rRNA (IGR). All the different morphospecies are color coded. Node values are posterior Bayesian probabilities, maximum likelihood (ML) bootstrap

values, and maximum parsimony (MP) bootstrap values. Posterior Bayesian probabilities below 0.7, ML bootstrap values below 50%, and MP bootstrap values below 50% are indicated by a dash (-). *Siderastrea radians* was selected as the outgroup

Arabian RS. After intensive sampling over 1200 km from the GofA in the north to the Farasan Islands in the south, we recorded a total of 20 agariciid morphospecies belonging to the genera

Gardineroseris, *Leptoseris*, and *Pavona* (Table 2). Taxa were investigated at two molecular regions: the mitochondrial IGR and the nuclear ITS1. The resulting phylogenetic reconstructions

Fig. 4 Phylogenetic reconstruction of the family Agariciidae from the Saudi Arabian Red Sea (RS) inferred from Bayesian inference (BI) analysis of the internal transcribed spacer 1 from nuclear ribosomal DNA (ITS1). All the different morphospecies are color coded as in Fig. 3. Node values are posterior Bayesian probabilities, maximum likelihood

(ML) bootstrap values, and maximum parsimony (MP) bootstrap values. Posterior Bayesian probabilities below 0.7, ML bootstrap values below 50%, and MP bootstrap values below 50% are indicated by a dash (-). *Siderastrea radians* was selected as the outgroup

showed agreement in resolving six species, *G. planulata*, *P. clavus*, *P. maldivensis*, *P. decussata*, *L. fragilis*, and *L. yabei*, while the boundaries among the remaining 15 species were unclear, with three main species complexes detected (Figs. 3 and 4, Table 3). The IGR and ITS1 sequences represent the first molecular information of Agariciidae from the RS and only the second one from the entire IO, adding to the examination by Moothien Pillay et al. (2006) of the genetic differences between

P. cactus and *P. decussata* from Mauritius. Additionally, seven species (*P. venosa*, *P. danai*, *P. diffluens*, *L. fragilis*, *L. explanata*, *L. glabra*, and *L. yabei*) were included in a molecular phylogenetic study for the first time in this study.

The morphospecies diversity that we reported from the Saudi Arabian RS is compatible with that reported in previous works from the region (Scheer and Pillai 1983; Sheppard and Sheppard 1991; Veron 2000), though with some distinct

Table 2 List of samples analyzed in the present study

Code	Identification	Collector	Locality	Latitude	Longitude	IGR	ITS1
SA0050	<i>Leptoseris scabra</i>	FB	Dolphen Lagoon, Farasan Banks	19°00.320'	40°08.889'	LT630767	LT630858-9
SA0052	<i>Pavona danae</i>	FB	Dolphen Lagoon, Farasan Banks	19°00.320''	40°08.889'	LT630768	LT630860-1
SA0053	<i>Pavona danae</i>	FB	Dolphen Lagoon, Farasan Banks	19°00.320'	40°08.889'	LT630769	LT630862
SA0076	<i>Pavona venosa</i>	FB	Ablo Reef, Farasan Banks	18°39.571'	40°49.618'	LT630770	LT630863-4
SA0077	<i>Pavona danae</i>	FB	Ablo Reef, Farasan Banks	18°39.571'	40°49.618'	LT630771	LT630865-6
SA0081	<i>Leptoseris incrustans</i>	FB	Ablo Reef, Farasan Banks	18°39.571'	40°49.618'	LT630772	LT630867-8
SA0083	<i>Leptoseris incrustans</i>	FB	Ablo Reef, Farasan Banks	18°39.571'	40°49.618'	LT630773	LT630869
SA0088	<i>Leptoseris glabra</i>	FB	Ablo Reef, Farasan Banks	18°39.571'	40°49.618'	LT630774	LT630870
SA0097	<i>Leptoseris explanata</i>	FB	Ablo Reef, Farasan Banks	18°39.571'	40°49.618'	LT630775	LT630871-2
SA0105	<i>Leptoseris fragilis</i>	FB	Ablo Reef, Farasan Banks	18°39.571'	40°49.618'	LT630776	–
SA0146	<i>Pavona danae</i>	FB	Marka Reef, Farasan Banks	18°13.233'	41°19.463'	LT630777	LT630873-4
SA0202	<i>Leptoseris explanata</i>	FB	Marka Reef, Farasan Banks	18°13.233'	41°19.463'	LT630778	–
SA0205	<i>Pavona cactus</i>	FB	Marka Reef, Farasan Banks	18°13.233'	41°19.463'	LT630779	LT630875-6
SA0210	<i>Pavona diffluens</i>	FB	Marka Reef, Farasan Banks	18°13.233'	41°19.463'	LT630780	LT630877-8
SA0231	<i>Pavona explanulata</i>	FB	Shi'b Rabid, Farasan Banks	18°04.385'	40°53.154'	LT630781	LT630879-80
SA0232	<i>Gardineroseris planulata</i>	FB	Shi'b Rabid, Farasan Banks	18°04.385'	40°53.154'	LT630782	LT630881-2
SA0269	<i>Pavona decussata</i>	FB	Shuma North, Farasan Banks	16°45.159'	41°36.297'	LT630783	LT630883-4
SA0368	<i>Pavona maldivensis</i>	FB	Shi'b Nazar, Thuwal	16°45.159'	41°36.297'	LT630784	LT630885
SA0369	<i>Pavona clavus</i>	FB	Shi'b Nazar, Thuwal	16°45.159'	41°36.297'	LT630785	LT630886
SA0393	<i>Leptoseris yabei</i>	FB	Palace Reef, Thuwal	22°18.256'	38°57.505'	LT630786	LT630887
SA0424	<i>Leptoseris yabei</i>	FB	Qita Al Kirsh, Thuwal	22°25.358'	38°59.461'	LT630787	LT630888
SA0676	<i>Leptoseris mycetoseroides</i>	FB	Al Wajh	25°23.515'	36°41.035'	LT630788	–
SA0726	<i>Pavona danae</i>	FB	Ras Al-Ubayd	26°44.167'	36°02.659'	LT630789	LT630889-90
SA0774	<i>Leptoseris yabei</i>	FB	Jaz'air Silah	27°38.302'	35°18.369'	LT630790	LT630891
SA0781	<i>Leptoseris mycetoseroides</i>	FB	Jaz'air Silah	27°38.302'	35°18.369'	LT630791	–
SA0787	<i>Leptoseris mycetoseroides</i>	FB	Jaz'air Silah	27°38.302'	35°18.369'	LT630792	LT630892
SA0810	<i>Gardineroseris planulata</i>	FB	Jaz'air Silah	27°38.302'	35°18.369'	LT630793	LT630893-4
SA0847	<i>Pavona maldivensis</i>	FB	Jazirat Burcan	27°49.003'	35°06.397'	LT630794	LT630895
SA0857	<i>Pavona varians</i>	FB	Jazirat Burcan	27°49.003'	35°06.397'	LT630795	LT630896-7
SA0884	<i>Leptoseris cf. hawaiiensis</i>	FB	Jazirat Burcan	27°49.003'	35°06.397'	LT630796	–
SA0893	<i>Pavona diffluens</i>	FB	Jazirat Burcan	27°49.003'	35°06.397'	LT630797	–
SA0929	<i>Leptoseris glabra</i>	FB	Magna Coast Guard, Gulf of Aqaba	28°24.232'	34°44.440'	LT630798	LT630898
SA0930	<i>Leptoseris scabra</i>	FB	Magna Coast Guard, Gulf of Aqaba	28°24.232'	34°44.440'	LT630799	–
SA0936	<i>Pavona frondifera</i>	FB	Magna Coast Guard, Gulf of Aqaba	28°24.232'	34°44.440'	LT630800	LT630899-900
SA0939	<i>Leptoseris glabra</i>	FB	Magna Coast Guard, Gulf of Aqaba	28°24.232'	34°44.440'	LT630801	LT630901-2
SA0941	<i>Leptoseris incrustans</i>	FB	Magna Coast Guard, Gulf of Aqaba	28°24.232'	34°44.440'	LT630802	LT630903-4
SA0942	<i>Pavona explanulata</i>	FB	Magna Coast Guard, Gulf of Aqaba	28°24.232'	34°44.440'	LT630803	LT630905
SA0943	<i>Leptoseris sp</i>	FB	Magna Coast Guard, Gulf of Aqaba	28°24.232'	34°44.440'	LT630804	LT630906
SA0967	<i>Leptoseris sp</i>	FB	Magna Coast Guard, Gulf of Aqaba	28°24.232'	34°44.440'	LT630805	–
SA0969	<i>Pavona varians</i>	FB	Magna Coast Guard, Gulf of Aqaba	28°24.232'	34°44.440'	LT630806	LT630907-8
SA1010	<i>Leptoseris fragilis</i>	FB	Magna Coast Guard, Gulf of Aqaba	28°24.232'	34°44.440'	LT630807	LT630909-10
SA1040	<i>Pavona frondifera</i>	FB	Magna Coast Guard, Gulf of Aqaba	28°24.232'	34°44.440'	LT630808	LT630911-2
SA1073	<i>Leptoseris incrustans</i>	FB	Ship Wreck Bay, Gulf of Aqaba	28°11.077'	34°38.287'	LT630809	LT630913-4
SA1076	<i>Leptoseris yabei</i>	FB	Ship Wreck Bay, Gulf of Aqaba	28°11.077'	34°38.287'	LT630810	–
SA1115	<i>Pavona frondifera</i>	FB	Al Wajh	25°21.694'	36°54.754'	LT630811	LT630915-6
SA1183	<i>Pavona frondifera</i>	FB	Shark Reef, Thuwal	22°25.543'	38°59.348'	LT630812	LT630917
SA1184	<i>Pavona diffluens</i>	FB	Shark Reef, Thuwal	22°25.543'	38°59.348'	LT630813	LT630918-9
SA1203	<i>Pavona explanulata</i>	FB	Shark Reef, Thuwal	22°25.543'	38°59.348'	LT630814	LT630920-1
SA1208	<i>Leptoseris mycetoseroides</i>	FB	Shark Reef, Thuwal	22°25.543'	38°59.348'	LT630815	LT630922-3

Table 2 (continued)

Code	Identification	Collector	Locality	Latitude	Longitude	IGR	ITS1
SA1211	<i>Pavona frondifera</i>	FB	Shark Reef, Thuwal	22°25.543'	38°59.348'	LT630816	–
SA1236	<i>Pavona danae</i>	FB	Shark Reef, Thuwal	22°25.543'	38°59.348'	LT630817	LT630924
SA1250	<i>Pavona cactus</i>	FB	Shark Reef, Thuwal	22°25.543'	38°59.348'	LT630818	LT630925
SA1252	<i>Pavona cactus</i>	FB	Shark Reef, Thuwal	22°25.543'	38°59.348'	LT630819	LT630926-7
SA1254	<i>Pavona danae</i>	FB	Shark Reef, Thuwal	22°25.543'	38°59.348'	LT630820	–
SA1283	<i>Pavona frondifera</i>	FB	Shark Reef, Thuwal	22°25.543'	38°59.348'	LT630821	LT630928-9
SA1285	<i>Leptoseris incrustans</i>	FB	Shark Reef, Thuwal	22°25.543'	38°59.348'	LT630822	–
SA1287	<i>Pavona decussata</i>	FB	Shark Reef, Thuwal	22°25.543'	38°59.348'	LT630823	LT630930-1
SA1288	<i>Pavona danae</i>	FB	Shark Reef, Thuwal	22°25.543'	38°59.348'	LT630824	LT630932-3
SA1290	<i>Pavona frondifera</i>	FB	Shark Reef, Thuwal	22°25.543'	38°59.348'	LT630825	LT630934-5
SA1304	<i>Leptoseris incrustans</i>	FB	Shark Reef, Thuwal	22°25.543'	38°59.348'	LT630826	LT630936-7
SA1608	<i>Pavona explanulata</i>	FB	Mamali Kebir, Farasan Islands	17°36.369'	41°40.214'	LT630827	LT630938-9
SA1630	<i>Leptoseris mycetoseroides</i>	TIT	Al Baghlah, Farasan Islands	16°58.734'	41°23.094'	LT630828	LT630940-1
SA1640	<i>Pavona maldivensis</i>	FB	Al Baghlah, Farasan Islands	16°58.734'	41°23.094'	LT630829	LT630942
SA1643	<i>Pavona frondifera</i>	FB	Al Baghlah, Farasan Islands	16°58.734'	41°23.094'	LT630830	LT630943
SA1644	<i>Leptoseris mycetoseroides</i>	FB	Al Baghlah, Farasan Islands	16°58.734'	41°23.094'	LT630831	LT630944
SA1653	<i>Pavona cactus</i>	FB	Al Baghlah, Farasan Islands	16°58.734'	41°23.094'	LT630832	LT630945-6
SA1700	<i>Pavona explanulata</i>	FB	Dhi Dahaya, Farasan Islands	16°52.377'	41°26.409'	LT630833	LT630947-8
SA2608	<i>Leptoseris incrustans</i>	MT	Al-Qunfidhah, Farasan Banks	19°09.483'	40°18.065'	LT630834	LT630949
SA2610	<i>Leptoseris mycetoseroides</i>	MT	Al-Qunfidhah, Farasan Banks	19°09.483'	40°18.065'	LT630835	LT630950-1
SA2611	<i>Leptoseris mycetoseroides</i>	MT	Al-Qunfidhah, Farasan Banks	19°09.483'	40°18.065'	LT630836	LT630952-3
SA2612	<i>Pavona varians</i>	MT	Al-Qunfidhah, Farasan Banks	19°09.483'	40°18.065'	LT630837	LT630954-5
SA2613	<i>Gardineroseris planulata</i>	MT	Al-Qunfidhah, Farasan Banks	19°09.483'	40°18.065'	LT630838	LT630956-7
SA2617	<i>Pavona varians</i>	RA	Al-Qunfidhah, Farasan Banks	19°09.483'	40°18.065'	LT630839	LT630958-9
SA2618	<i>Pavona decussata</i>	RA	Al-Qunfidhah, Farasan Banks	19°09.483'	40°18.065'	LT630840	LT630960
SA2621	<i>Leptoseris mycetoseroides</i>	RA	Al-Qunfidhah, Farasan Banks	19°09.483'	40°18.065'	LT630841	LT630961
SA2622	<i>Pavona maldivensis</i>	RA	Al-Qunfidhah, Farasan Banks	19°09.483'	40°18.065'	LT630842	LT630962
SA2623	<i>Leptoseris incrustans</i>	RA	Al-Qunfidhah, Farasan Banks	19°09.483'	40°18.065'	LT630843	LT630963-4
SA2624	<i>Leptoseris mycetoseroides</i>	RA	Al-Qunfidhah, Farasan Banks	19°09.483'	40°18.065'	LT630844	LT630965-6
SA2625	<i>Leptoseris mycetoseroides</i>	RA	Al-Qunfidhah, Farasan Banks	19°09.483'	40°18.065'	LT630845	LT630967
SA2626	<i>Leptoseris scabra</i>	RA	Al-Qunfidhah, Farasan Banks	19°09.483'	40°18.065'	LT630846	LT630968-9
SA2634	<i>Pavona diffluens</i>	RA	Ablo Reef, Farasan Banks	18°42.340'	40°39.240'	LT630847	LT630970
SA2640	<i>Pavona explanulata</i>	RA	Ablo Reef, Farasan Banks	18°42.340'	40°39.240'	LT630848	LT630971-2
SA2652	<i>Pavona diffluens</i>	RA	Ablo Reef, Farasan Banks	18°42.340'	40°39.240'	LT630849	LT630973-4
SA2653	<i>Pavona frondifera</i>	MT	Al-Qunfidhah, Farasan Banks	19°00.186'	40°16.993'	LT630850	LT630975-6
SA2654	<i>Leptoseris scabra</i>	MT	Al-Qunfidhah, Farasan Banks	19°00.186'	40°16.993'	LT630851	LT630977-8
SA2658	<i>Pavona maldivensis</i>	MT	Al-Qunfidhah, Farasan Banks	19°00.186'	40°16.993'	LT630852	LT630979
SA2665	<i>Leptoseris mycetoseroides</i>	RA	Al-Qunfidhah, Farasan Banks	19°00.186'	40°16.993'	LT630853	LT630980-1
SA2667	<i>Pavona maldivensis</i>	RA	Al-Qunfidhah, Farasan Banks	19°00.186'	40°16.993'	LT630854	LT630982
SA2668	<i>Gardineroseris planulata</i>	RA	Al-Qunfidhah, Farasan Banks	19°00.186'	40°16.993'	LT630855	LT630983-4
SA2675	<i>Pavona varians</i>	RA	Al-Qunfidhah, Farasan Banks	19°00.186'	40°16.993'	LT630856	LT630985-6
SA2698	<i>Leptoseris glabra</i>	RA	Al-Qunfidhah, Farasan Banks	19°00.186'	40°16.993'	LT630857	LT630987-8

For each specimen, the voucher code, identification of the matching specimen, collector, collection locality, latitude (N), longitude (E), and EMBL accession numbers of each molecular marker are provided. FB = F. Benzoni; MT = M. Tietbohl; RA = R. Arrigoni; TIT = T.I. Terraneo

differences (Table 2). Notably, we recorded *L. glabra* from the RS for the first time. However, it is possible that previous authors recorded it as *L. explanata* due to its similar

morphology. According to Dinesen (1980), the material identified as *L. explanata* by Veron and Pichon (1980; figs 71–82) actually belongs to *L. glabra*. Dinesen (1980) also indicated

Table 3 List of Agariciidae morphospecies collected in the Red Sea (RS) and relative genetic assignment from IGR and ITS1 molecular markers

	IGR	ITS1
<i>Gardineroseris planulata</i>	CLADE V	CLADE V
<i>Pavona cactus</i>	CLADE XIV	CLADE XIV
<i>Pavona clavus</i>	CLADE VI	CLADE VI
<i>Pavona danai</i>	CLADE XIV	CLADE XIV
<i>Pavona decussata</i>	CLADE IV	CLADE IV
<i>Pavona diffluens</i>	CLADE XII	Unresolved
<i>Pavona explanulata</i>	CLADES VIII–X	Unresolved
<i>Pavona frondifera</i>	CLADE XIV	CLADE XIV
<i>Pavona maldivensis</i>	CLADE II	CLADE II
<i>Pavona varians</i>	CLADE I	CLADES I–IV
<i>Pavona venosa</i>	CLADE I	CLADES I–IV
<i>Leptoseris explanata</i>	CLADE XII	Unresolved
<i>Leptoseris fragilis</i>	CLADE III	CLADE III
<i>Leptoseris glabra</i>	CLADE XII	Unresolved
<i>Leptoseris cf. hawaiiensis</i>	CLADE XII	Unresolved
<i>Leptoseris incrustans</i>	CLADE XII	Unresolved
<i>Leptoseris mycetoseroides</i>	CLADES IX–XI	Unresolved
<i>Leptoseris scabra</i>	CLADE XII	Unresolved
<i>Leptoseris sp</i>	CLADE VII	CLADE VII
<i>Leptoseris yabei</i>	CLADE XIII	CLADE XIII

that the type material of *L. explanata* shown and described in Yabe and Sugiyama (1941: pl. 63, fig. 3e) is morphologically distinct from *L. glabra*, as she described in her revision of *Leptoseris* (Dinesen 1980). In our identifications, we followed the original descriptions of these two morphospecies and considered them as morphologically distinct following Dinesen (1980). Nevertheless, the recovery of specimens ascribed to these two species in the same lineage with other *Leptoseris* merits for further investigation into the relationship between these species. While the existing confusion in the literature concerning *L. explanata* and *L. glabra* is easily explained based on their undeniable similarities in terms of calice size and arrangement, they are both markedly distinct from the other morphospecies recovered in the same clade having smaller calices and more numerous septa (Fig. 2). Easily recognizable and common shallow-water species such as *G. planulata*, *P. cactus*, *P. decussata*, *P. explanulata*, *P. maldivensis*, *P. varians*, *L. explanata*, *L. hawaiiensis*, *L. mycetoseroides*, *L. scabra*, and *L. yabei* were recorded by Scheer and Pillai (1983), Sheppard and Sheppard (1991), Veron (2000), and this work. Compared to Sheppard and Sheppard (1991) and Veron (2000), we did not record *P. duerdeni* and *L. foliosa* during our collection. The former species is rare in the RS, although it can form conspicuous colonies in low-energy waters (Veron 2000), whereas the latter species is uncommon and preferentially found in deep-water habitats (>30 m) (Sheppard and

Sheppard 1991). This work did not investigate mesophotic coral ecosystems, potentially explaining why these species were likely missed. The family Agariciidae is among the most abundant coral families in deep-water oceanic habitats (Lesser et al. 2009). Therefore, more species are likely to occur in deeper waters of the RS. *Leptoseris* often shows an increasing abundance with depth in the IO (Bouchon 1981) and in Hawaii (Kahng and Kelley 2007), where it dominates hard substrates between 60 and 120 m and is, instead, rare in shallow-water reefs. Luck et al. (2013) also found cryptic lineages of *Leptoseris* across depth gradients in Hawaii, and several studies showed limited larval dispersal, genetic segregation, and low gene flow between shallow and mesophotic agariciid reef corals (Bongaerts et al. 2013, 2015; Pochon et al. 2015). Bongaerts et al. (2013) showed a division of *Agaricia* species and associated *Symbiodinium* into two main genetic lineages that matched a clear bathymetric distribution. For these reasons, we caution others to consider our presented survey as fully representative of only the shallow-water genetic diversity of Agariciidae from the Saudi Arabian RS; we strongly encourage further investigations of mesophotic reefs.

Regarding the molecular information, our work further highlighted the hypothesis that some regions of the mitochondrial DNA of corals can be informative in defining boundaries among closely related taxa. Despite the well-documented slow evolution of Scleractinia mitochondrial DNA (Shearer et al. 2002; Hellberg 2006; Huang et al. 2008), recent works have found that some mitochondrial non-coding regions show rates of interspecific polymorphisms comparable to these of nuclear loci and can be phylogenetically informative at low taxonomic levels (Kitahara et al. 2016). The IGR sequenced in the present work has been previously used for delimiting species boundaries in several scleractinian families and genera, including the families Merulinidae Verrill, 1865 and Agariciidae (Kitahara et al. 2016), the genus *Pachyseris* Milne Edwards & Haime, 1849 (Terraneo et al. 2014), and the genera *Sclerophyllia* Klunzinger, 1879, *Homophyllia* Brüggemann, 1877, and *Micromussa* Veron, 2000 (Arrigoni et al. 2015, 2016b). Ribosomal DNA is widely used for low-level taxonomic comparisons in corals, particularly due to high interspecific divergence between the two internal transcribed spacers regions, ITS1 and ITS2 (Kitahara et al. 2016). In the present study, the phylogeny reconstruction based on ITS1 was poorly resolved, as it was characterized by a large basal polytomy and several species were not differentiated. The absence of genetic differentiation among the analyzed species may be explained by insufficient resolution of the ITS1 locus. Hence, we encourage the investigation of further molecular markers and modern genome-wide sequencing techniques that could provide strong phylogenomic analysis to help better determine species boundaries (Pante et al. 2015).

This work raises several questions regarding the classical definition of genera and species boundaries within the

Agariciidae that will require more in-depth observations of skeletal micromorphology before a formal taxonomic revision of the family is proposed. As previously reported by Luck et al. (2013), our mitochondrial phylogeny reconstruction confirmed that *Pavona* and *Leptoseris* are not monophyletic. Our molecular analyses (based on IGR and ITS1) revealed for the first time that the monotypic *Gardineroseris* is closely related to both *Pavona* and *Leptoseris* and it nests within them. The genetic information does not support the morphology-based distinction among these three genera (Scheer and Pillai 1974; Dinesen 1980) and suggests the presence of a single, species-rich genus. However, any taxonomic reclassification is not recommended until a comprehensive phylogenetic reconstruction of the Agariciidae, including all seven extant genera and sequencing of slow-evolving loci such as the mitochondrial COI and 16S rDNA genes and the nuclear 28S region, is conducted (Fukami et al. 2008; Huang et al. 2008). Indeed, non-coding regions are usually considered to be poorly informative at high taxonomic levels because homoplasy caused by repeated mutations in saturated positions is thought to undermine the phylogenetic signal (Kelchner 2000). At the species level, the IGR and ITS1 phylogeny reconstructions revealed the presence of three species complexes (Fig. 3, Table 3). On one hand, the recovery of the two species complexes composed of the frondose and highly plastic *P. cactus*, *P. danai*, and *P. frondifera* and of the encrusting and highly carinated *P. varians* and *P. venosa* (Veron and Pichon 1980) could be morphologically plausible. Indeed, these species display similar colony morphology, related collines and carinae, comparable corallite dimension and organization, and previous taxonomists have identified these morphological affinities (Vaughan 1918; Crossland 1952; Veron and Pichon 1980; Scheer and Pillai 1983; Veron 2000). The recovery of a molecular clade including *L. explanata*, *Leptoseris* cf. *hawaiiensis*, *L. scabra*, *L. incrustans*, *L. glabra*, and *P. diffluens* was unexpected. Although *P. diffluens* may be mistaken for massive parts of *P. explanulata* (Sheppard and Sheppard 1991), it does not resemble any *Leptoseris* species ascribed to this molecular group. Moreover, as mentioned earlier in this discussion, the *Leptoseris* morphospecies recovered in this one cohesive lineage in the RS do display remarkable differences in terms of colony growth form, calice size and organization, and septocostae thickness, with the notable exception of *L. glabra* and *L. explanata* (Fig. 2). A similar situation has been described in the genus *Stylophora* from the RS (Arrigoni et al. 2016c), where different morphospecies, two of which are endemic to the RS, could not be differentiated based on genetic results. Several hypotheses can be suggested to explain this finding, such as taxonomic misidentification, uninformative morphological characters, non-suitable molecular loci, or incipient speciation.

In conclusion, we investigated the genetic diversity of Agariciidae from the isolated Saudi Arabian RS and provide

a regional genetic database for future molecular comparisons between the RS and other localities. The partial disagreement between morphology-based taxonomy and genetic results highlights that agariciid systematics is poorly understood and that the current family taxonomy does not truly represent the evolutionary relationships at the genus or species levels. A combined morpho-molecular approach with a strong focus on skeletal micromorphology is likely necessary to better clarify the boundaries within the family and revise its taxonomy, as previously demonstrated for the agariciid *Dactylotrachus* (Kitahara et al. 2012) and several other scleractinian families (Kitahara et al. 2016). The molecular phylogeny of Agariciidae is still incomplete because only a few regions and species have been genetically characterized to date (Maté 2003; Moothien Pillay et al. 2006; Benzoni et al. 2012; Luck et al. 2013; Pochon et al. 2015; Waheed et al. 2015). Therefore, a more global sampling effort with a larger number of taxa and more genetic data, in addition to detailed comparative micromorphological analysis, are needed and necessary before a robust molecular hypothesis can be produced and the agariciids fully resolved.

Acknowledgments This research was undertaken in accordance with the policies and procedures of the King Abdullah University of Science and Technology (KAUST). Permissions relevant for KAUST to undertake the research have been obtained from the applicable governmental agencies in the Kingdom of Saudi Arabia. We wish to thank the captain and crew of the MV Dream Master, the KAUST Coastal and Marine Resources Core Lab, A. Gusti (KAUST), and T.H. Sinclair-Taylor (KAUST). T.I.T. and R.A. gratefully acknowledge J.D. DiBattista (CURTIN) and P. Saenz-Agudelo (UACH) for their assistance at KAUST. This project was supported by funding from KAUST (award nos. URF/1/1389-01-01 and FCC/1/1973-07, and baseline research funds to M.L.B.).

References

- Arrigoni R, Stefani F, Pichon M, Galli P, Benzoni F (2012) Molecular phylogeny of the robust clade (Faviidae, Mussidae, Merulinidae, and Pectiniidae): an Indian Ocean perspective. *Mol Phylogenet Evol* 65:183–193
- Arrigoni R, Berumen ML, Terraneo TI, Caragnano A, Bouwmeester J, Benzoni F (2015) Forgotten in the taxonomic literature: resurrection of the scleractinian coral genus *Sclerophyllia* (Scleractinia, Lobophylliidae) from the Arabian peninsula and its phylogenetic relationships. *Syst Biodivers* 13:140–163
- Arrigoni R, Berumen ML, Chen CA, Terraneo TI, Baird AH, Payri C, Benzoni F (2016a) Species delimitation in the reef coral genera *Echinophyllia* and *Oxypora* (Scleractinia, Lobophylliidae) with a description of two new species. *Mol Phylogenet Evol* 105:146–159
- Arrigoni R, Benzoni F, Huang D, Fukami H, Chen CA, Berumen ML, Hoogenboom M, Thomson DP, Hoeksema BW, Budd AF, Zayasu Y, Terraneo TI, Kitano YF, Baird AH (2016b) When forms meet genes: revision of the scleractinian genera *Micromussa* and *Homophyllia* (Lobophylliidae) with a description of two new species and one new genus. *Contrib Zool* 85:387–422
- Arrigoni R, Benzoni F, Terraneo TI, Caragnano A, Berumen ML (2016c) Recent origin and semi-permeable species boundaries in the

- scleractinian coral genus *Stylophora* from the Red Sea. *Sci Rep* 6: 34612
- Arrigoni R, Berumen ML, Huang D, Terraneo TI, Benzoni F (2017) *Cyphastrea* (Cnidaria: Scleractinia: Merulinidae) in the Red Sea: phylogeny and a new reef coral species. *Invertebr Syst* 31:141–156
- Benzoni F, Arrigoni R, Stefani F, Stolarski J (2012) Systematics of the coral genus *Craterastrea* (Cnidaria, Anthozoa, Scleractinia) and description of a new family through combined morphological and molecular analyses. *Syst Biodivers* 10:417–433
- Berumen ML, Hoey AS, Bass WH, Bouwmeester J, Catania D, Cochran JE, Khalil MT, Miyake S, Mughal MR, Spaet JL, Saenz-Agudelo P (2013) The status of coral reef ecology research in the Red Sea. *Coral Reefs* 32:737–748
- Bongaerts P, Frade PR, Ogier JJ, Hay KB, Van Bleijswijk J, Englebert N, Vermeij MJ, Bak RP, Visser PM, Hoegh-Guldberg O (2013) Sharing the slope: depth partitioning of agariciid corals and associated *Symbiodinium* across shallow and mesophotic habitats (2–60 m) on a Caribbean reef. *BMC Evol Biol* 13:205
- Bongaerts P, Frade PR, Hay KB, Englebert N, Latijnhouwers KR, Bak RP, Vermeij MJ, Hoegh-Guldberg O (2015) Deep down on a Caribbean reef: lower mesophotic depths harbor a specialized coral-endosymbiont community. *Sci Rep* 5:7652
- Bouchon C (1981) Quantitative study of the scleractinian coral communities of a fringing reef of Reunion Island (Indian Ocean). *Mar Ecol Prog Ser* 4:273–288
- Castresana J (2000) Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Mol Biol Evol* 17:540–552
- Coleman RR, Eble JA, DiBattista JD, Rocha LA, Randall JE, Berumen ML, Bowen BW (2016) Regal phylogeography: range-wide survey of the marine angelfish *Pygoplites diacanthus* reveals evolutionary partitions between the Red Sea, Indian Ocean, and Pacific Ocean. *Mol Phylogenet Evol* 100:243–253
- Crossland C (1952) Madreporaria, Hydrocorallinae, Heliopora and Tubipora. *Br Mus Nat Hist* 6:85–257
- DiBattista JD, Berumen ML, Gaither MR, Rocha LA, Eble JA, Choat JH, Craig MT, Skillings DJ, Bowen BW (2013) After continents divide: comparative phylogeography of reef fishes from the Red Sea and Indian Ocean. *J Biogeogr* 40:1170–1181
- DiBattista JD, Waldrop E, Rocha LA, Craig MT, Berumen ML, Bowen BW (2015) Blinded by the bright: a lack of congruence between colour morphs, phylogeography and taxonomy for a cosmopolitan Indo-Pacific butterflyfish, *Chaetodon auriga*. *J Biogeogr* 42:1919–1929
- DiBattista JD, Howard Choat J, Gaither MR, Hobbs JP, Lozano-Cortés DF, Myers RF, Paulay G, Rocha LA, Toonen RJ, Westneat MW, Berumen ML (2016a) On the origin of endemic species in the Red Sea. *J Biogeogr* 43:13–30
- DiBattista JD, Roberts MB, Bouwmeester J, Bowen BW, Coker DJ, Lozano-Cortés DF, Howard Choat J, Gaither MR, Hobbs JP, Khalil MT, Kochzius M, Myers RF, Paulay G, Robitzsch VSN, Saenz-Agudelo P, Salas E, Sinclair-Taylor TH, Toonen RJ, Westneat MW, Williams ST, Berumen ML (2016b) A review of contemporary patterns of endemism for shallow water reef fauna in the Red Sea. *J Biogeogr* 43:423–439
- Dinesen ZD (1980) A revision of the coral genus *Leptoseris* (Scleractinia: Fungiina: Agariciidae). *Mem Queensl Mus* 20:181–235
- Ehrenberg CG (1834) Beiträge zur physiologischen Kenntniss der Corallenthiere im Allgemeinen, und besonders des Rothen Meeres, nebst einem Versuche zur physiologischen Systematik derselben. *Abhandlungen der Königlischen Akademie der Wissenschaften*, Berlin
- Ekman S (1953) Zoogeography of the sea. Sidgwick & Jackson, London
- Flot JF (2007) Champuru 1.0: a computer software for unraveling mixtures of two DNA sequences of unequal lengths. *Mol Ecol Resour* 7: 974–977
- Flot JF (2010) SeqPHASE: a web tool for interconverting PHASE input/output files and FASTA sequence alignments. *Mol Ecol Resour* 10: 162–166
- Forskål P (1775) Descriptiones animalium, avium, amphibiorum, piscium, insectorum, vermium; quæ in itinere orientali observavit Petrus Forskål. Hauniæ, Heineck et Faber, Copenhagen
- Fricke HW, Knauer B (1986) Diversity and spatial pattern of coral communities in the Red Sea upper twilight zone. *Oecologia* 71:29–37
- Fricke HW, Vareschi E, Schlichter D (1987) Photoecology of the coral *Leptoseris fragilis* in the Red Sea twilight zone (an experimental study by submersible). *Oecologia* 73:371–381
- Fukami H, Chen CA, Budd AF, Collins A, Wallace C, Chuang YY, Chen C, Dai CF, Iwao K, Sheppard C, Knowlton N (2008) Mitochondrial and nuclear genes suggest that stony corals are monophyletic but most families of stony corals are not (order Scleractinia, class Anthozoa, phylum Cnidaria). *PLoS One* 3:e3222
- Guindon S, Gascuel O (2003) A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Syst Biol* 52: 696–704
- Hall TA (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp Ser* 41:95–98
- Hellberg ME (2006) No variation and low synonymous substitution rates in coral mtDNA despite high nuclear variation. *BMC Evol Biol* 6:24
- Hinderstein LM, Marr JCA, Martinez FA, Dowgiallo MJ, Puglise KA, Pyle RL, Zawada DG, Appeldoorn R (2010) Theme section on “Mesophotic coral ecosystems: characterization, ecology, and management”. *Coral Reefs* 29:247–251
- Hodge JR, van Herwerden L, Bellwood DR (2014) Temporal evolution of coral reef fishes: global patterns and disparity in isolated locations. *J Biogeogr* 41:2115–2127
- Huang D, Meier R, Todd PA, Chou LM (2008) Slow mitochondrial COI sequence evolution at the base of the metazoan tree and its implications for DNA barcoding. *J Mol Evol* 66:167–174
- Kahng SE, Kelley CD (2007) Vertical zonation of megabenthic taxa on a deep photosynthetic reef (50–140 m) in the Au’au Channel, Hawaii. *Coral Reefs* 26:679–687
- Kahng SE, Garcia-Sais JR, Spalding HL, Brokovich E, Wagner D, Weil E, Hinderstein L, Toonen RJ (2010) Community ecology of mesophotic coral reef ecosystems. *Coral Reefs* 29:255–275
- Kahng SE, Copus JM, Wagner D (2014) Recent advances in the ecology of mesophotic coral ecosystems (MCEs). *Curr Opin Environ Sustain* 7:72–81
- Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol* 30:772–780
- Kelchner SA (2000) The evolution of non-coding chloroplast DNA and its application in plant systematics. *Ann Mo Bot Gard* 87:482–498
- Kitahara MV, Cairns SD, Stolarski J, Blair D, Miller DJ (2010) A comprehensive phylogenetic analysis of the Scleractinia (Cnidaria, Anthozoa) based on mitochondrial CO1 sequence data. *PLoS One* 5(7):e11490. doi:10.1371/journal.pone.0011490
- Kitahara MV, Stolarski J, Cairns SD, Benzoni F, Stake JL, Miller DJ (2012) The first modern solitary Agariciidae (Anthozoa, Scleractinia) revealed by molecular and microstructural analysis. *Invertebr Syst* 26:303–315
- Kitahara MV, Fukami H, Benzoni F, Huang D (2016) The new systematics of Scleractinia: integrating molecular and morphological evidence. In: Goffredo S, Dubinsky Z (eds) *The Cnidaria, past, present and future: the world of Medusa and her sisters*. Springer, Dordrecht, pp 41–60
- Kitano YF, Benzoni F, Arrigoni R, Shirayama Y, Wallace CC, Fukami H (2014) A phylogeny of the family Poritidae (Cnidaria, Scleractinia) based on molecular and morphological analyses. *PLoS One* 9: e98406

- Klunzinger CB (1870) Synopsis der fische des Rothen Meeres. C. Ueberreuter'shBuchdruckerei, Wien
- Klunzinger CB (1877) Die Korallthiere des Rothen Meeres. Gutmann'schen, Berlin
- Ladner JT, Palumbi SR (2012) Extensive sympatry, cryptic diversity and introgression throughout the geographic distribution of two coral species complexes. *Mol Ecol* 21:2224–2238
- Lamarck JBP (1816) Histoire naturelle des Animaux sans Vertèbres, présentant les caractères généraux et particuliers de ces animaux, leur distribution, leurs classes, leurs familles, leurs genres, et la citation des principales espèces qui s'y rapportent; précédée d'une Introduction offrant la détermination des caractères essentiels de l'animal, sa distinction du végétal et des autres corps naturels, enfin, l'exposition des principes fondamentaux de la zoologie. Détérville & Verdrière, Paris
- Lesser MP, Slattery M, Leichter JJ (2009) Ecology of mesophotic coral reefs. *J Exp Mar Biol Ecol* 375:1–8
- Librado P, Rozas J (2009) DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics* 25:1451–1452
- Loya Y, Genin A, el-Zibdeh M, Naumann MS, Wild C (2014) Reviewing the status of coral reef ecology of the Red Sea: key topics and relevant research. *Coral Reefs* 33:1179–1180
- Luck DG, Forsman ZH, Toonen RJ, Leicht SJ, Kahng SE (2013) Polyphyly and hidden species among Hawai'i's dominant mesophotic coral genera, *Leptoseris* and *Pavona* (Scleractinia: Agariciidae). *Peer J* 1:e132
- Maragos JE, Jokiel PL (1986) Reef corals of Johnston atoll: one of the world's most isolated reefs. *Coral Reefs* 4:141–150
- Maté J (2003) Ecological, genetic, and morphological differences among three *Pavona* (Cnidaria: Anthozoa) species from the Pacific coast of Panama. *Mar Biol* 142:427–440
- Moothien Pillay KR, Aasahida T, Chen CA, Terashima H, Ida H (2006) ITS ribosomal DNA distinctions and the genetic structures of populations of two sympatric species of *Pavona* (Cnidaria: Scleractinia) from Mauritius. *Zool Stud* 45:132–144
- Nishihira M, Veron JEN (1995) Hermatypic corals of Japan. Kaiyusha, Tokyo
- Obura D (2012) The diversity and biogeography of western Indian Ocean reef-building corals. *PLoS One* 7:e45013
- Pante E, Abdelkrim J, Viricel A, Gey D, France SC, Boisselier MC, Samadi S (2015) Use of RAD sequencing for delimiting species. *Heredity* 114(5):450–459
- Pillai CSG, Scheer G (1976) Report on the stony corals from the Maldive Archipelago. Results of the Xarifa Expedition 1957/58. *Zoologica, Stuttgart* 43(126):1–83
- Pochon X, Forsman ZH, Spalding HL, Padilla-Gamiño JL, Smith CM, Gates RD (2015) Depth specialization in mesophotic corals (*Leptoseris* spp.) and associated algal symbionts in Hawai'i. *R Soc Open Sci* 2:140351
- Posada D, Crandall KA (1998) MODELTEST: testing the model of DNA substitution. *Bioinformatics* 14(9):817–818
- Rambaut A, Drummond AJ (2007) Tracer v1.4. Available from <http://beast.bio.ed.ac.uk/Tracer>
- Richards ZT, Berry O, van Oppen MJ (2016) Cryptic genetic divergence within threatened species of *Acropora* Coral from the Indian and Pacific oceans. *Conserv Genet* 17:577–591
- Ronquist F, Huelsenbeck JP (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19:1572–1574
- Rooney J, Donham E, Montgomery A, Spalding H, Parrish F, Boland R, Fenner D, Gove J, Vetter O (2010) Mesophotic coral ecosystems in the Hawaiian archipelago. *Coral Reefs* 29:361–367
- Scheer G, Pillai CSG (1974) Report on the Scleractinia from the Nicobar Islands. E. Schweizerbart'sche Verlagsbuchhandlung, Stuttgart
- Scheer G, Pillai CSG (1983) Report on the stony corals from the Red Sea. E. Schweizerbart'sche Verlagsbuchhandlung, Stuttgart
- Shearer TL, Van Oppen MJH, Romano SL, Wörheide G (2002) Slow mitochondrial DNA sequence evolution in the Anthozoa (Cnidaria). *Mol Ecol* 11:2475–2487
- Sheppard CRC, Sheppard ALS (1991) Corals and coral communities of Saudi Arabia. *Fauna Saudi Arabia* 12:1–170
- Sheppard CRC, Price A, Roberts C (1992) Marine ecology of the Arabian region. Academic Press, London
- Stefani F, Benzoni F, Yang SY, Pichon M, Galli P, Chen CA (2011) Comparison of morphological and genetic analyses reveals cryptic divergence and morphological plasticity in *Stylophora* (Cnidaria, Scleractinia). *Coral Reefs* 30:1033–1049
- Stephens M, Smith N, Donnelly P (2001) A new statistical method for haplotype reconstruction from population data. *Am J Hum Genet* 68:978–989
- Swofford DL (2002) PAUP*. Phylogenetic Analysis Using Parsimony (*and other methods). Version 4. Sinauer Associates, Sunderland
- Terraneo TI, Berumen ML, Arrigoni R, Waheed Z, Bouwmeester J, Caragnano A, Stefani F, Benzoni F (2014) *Pachyseris inattesa* sp. n. (Cnidaria, Anthozoa, Scleractinia): a new reef coral species from the Red Sea and its phylogenetic relationships. *ZooKeys* 433:1–33
- Terraneo TI, Benzoni F, Arrigoni R, Berumen ML (2016) Species delimitation in the coral genus *Goniopora* (Scleractinia, Poritidae) from the Saudi Arabian Red Sea. *Mol Phylogenet Evol* 102:278–294
- Vaughan TW (1918) Some shoal-water corals from Murray Island (Australia), Cocos-Keeling Islands, and Fanning Island. Carnegie Institution of Washington, Washington
- Vaughan TW, Wells JW (1943) Revision of the suborders, families, and genera of the Scleractinia. *Geol Soc Am Special Paper* 44:1–394
- Veron JEN (2000) Corals of the world. Australian Institute of Marine Science, Townsville
- Veron JEN, Pichon M (1980) Scleractinia of eastern Australia. Part III: families Agariciidae, Siderastreidae, Fungiidae, Oculinidae, Merulinidae, Mussidae, Pectiniidae, Caryophylliidae, Dendrophylliidae. *Aust Inst Mar Sci Monogr* 4:1–422
- Veron J, Stafford-Smith M, DeVantier L, Turak E (2015) Overview of distribution patterns of zooxanthellate Scleractinia. *Front Mar Sci* 1: 81
- Waheed Z, Hoeksema BW (2014) Diversity patterns of scleractinian corals at Kota Kinabalu, Malaysia, in relation to exposure and depth. *Raffles Bull Zool* 62:66–82
- Waheed Z, Benzoni F, van der Meij SE, Terraneo TI, Hoeksema BW (2015) Scleractinian corals (Fungiidae, Agariciidae and Euphylliidae) of Pulau Layang-Layang, Spratly Islands, with a note on *Pavona maldivensis* (Gardiner, 1905). *ZooKeys* 517:1–37
- Wei NW, Wallace CC, Dai CF, Pillay KRM, Chen CA (2006) Analyses of the ribosomal internal transcribed spacers (ITS) and 5.8 S gene indicate that extremely high rDNA heterogeneity is a unique feature in the scleractinian coral genus *Acropora* (Scleractinia; Acroporidae). *Zool Stud* 45:404–418
- Wells JW (1956) Scleractinia. In: Moore RC (ed) Treatise on invertebrate paleontology, part F (Coelenterata). The University of Kansas Press, Lawrence, pp F328–F444
- WoRMS (2016) World register of marine species. Available from <http://www.marinespecies.org> at VLIZ. Accessed 29 Sep 2016
- Yabe H, Sugiyama T (1941) Recent reef-building corals from Japan and the South Sea Islands under the Japanese mandate II. The Science reports of the Tôhoku Imperial University, Sendai. 2nd Series (Geologie) Special Volume 2:67–91
- Ziegler M, Roder CM, Büchel C, Voolstra CR (2015) Mesophotic coral depth acclimatization is a function of host-specific symbiont physiology. *Front Mar Sci* 2:1–4