Global ecological impacts of marine exotic species

Andrea Anton^{1*}, Nathan R. Galdi¹, Catherine E. Lovelock², Eugenia T. Apostolaki³, Scott Bennett⁴, Just Cebrian^{5,6,7}, Dorte Krause-Jensen^{8,9}, Nuria Marbà⁴, Paulina Martinetto¹⁰, John M. Pandolfi¹¹, Julia Santana-Garcon⁴ and Carlos M. Duarte^{1,12}

Exotic species are a growing global ecological threat; however, their overall effects are insufficiently understood. While some exotic species are implicated in many species extinctions, others can provide benefits to the recipient communities. Here, we performed a meta-analysis to quantify and synthesize the ecological effects of 76 exotic marine species (about 6% of the listed exotics) on ten variables in marine communities. These species caused an overall significant, but modest in magnitude (as indicated by a mean effect size of $g < 0.2$), decrease in ecological variables. Marine primary producers and predators were the most disruptive trophic groups of the exotic species. Approximately 10% (that is, 2 out of 19) of the exotic species assessed in at least three independent studies had significant impacts on native species. Separating the innocuous from the disruptive exotic species provides a basis for triage efforts to control the marine exotic species that have the most impact, thereby helping to meet Aichi Biodiversity Target 9 of the Convention on Biological Diversity.

Exotic species rank among the most serious environmental threats of the new millennium^{1–3}, as the relocation of species beyond their native range has intensified over the last five decades⁴. The impacts of some of the worst exotic species on biodiversity are well documented, as they are implicated in the local extinction of hundreds of native species^{5–7}. While the ecological outcomes of the most damaging exotic species are clearly dramatic, the impact of many appears to be modest, with some even providing ecological benefits^{8,9}. These contrasting effects have ignited a contemporary debate about the overall impacts and perceptions of exotic species^{10–14}, and this debate can only be resolved by distinguishing innocuous from disruptive exotic species¹⁵. In the case of marine exotic species, there are no documented global extinctions attributed to them¹⁶ (but see also refs. ^{17–19}), although some can substantially reduce native fauna²⁰.

Based on previous syntheses and meta-analyses, we made predictions about the expected ecological effects of marine exotic species^{21–27} (Supplementary Table 1). Marine ecosystems differ from other environments in a number of characteristics that may mitigate the effects of exotic species on the abundance of native species. The open nature of oceans, with large and continuous marine habitats, and the extensive dispersal potential of many marine species are likely to buffer the effects of exotics through a high capacity to repopulate affected areas. We predict that marine exotic primary producers and predators will exert strong effects on native communities, as reported previously for aquatic and terrestrial systems^{21–23,25}. However, marine communities harbour a high proportion of generalist consumers (herbivores and omnivores²⁸) that could limit the effects of exotic primary producers. In terrestrial communities²⁵, exotic herbivores have nonsignificant effects

on the richness of native species, but marine exotic herbivores can cause reductions in the abundance and richness of native species^{29,30}. We expect, as has been reported for other systems, contrasting effects of marine exotic detritivores and overall innocuous effects of exotic omnivores^{25,27}. Exotic seaweeds exert persistent negative effects on native primary producers, but no effects on herbivores and predators²². Finally, a previous meta-analysis²⁴ of marine exotic ecosystem engineers indicated overall nonsignificant effects at the species and community levels and contrasting effects at the ecosystem level.

However, to date, the global effect of exotic species, including higher-order trophic groups, has not been systematically examined in marine ecosystems. Despite this, exotic marine species directly affect 12% of the marine species registered on the Red List of the International Union for Conservation of Nature (IUCN). This value is substantially higher than the percentage of vulnerable species affected by terrestrial and freshwater exotic species (6% and 2%, respectively)¹⁶. In fact, the signatories of the Convention on Biological Diversity of the United Nations agreed to achieve Aichi Biodiversity Target 9 by 2020, which aims, among other things, to identify and then control or eradicate the worst invasive species³¹. However, this target is not on track, as a systematic assessment of the quantitative ecological effects of exotic species has not yet been produced for the marine environment¹⁵.

Here, we quantified the ecological effects of exotic species on marine communities through an exhaustive meta-analysis of published studies (Supplementary Table 2). We used the effect size Hedges' g as a common metric to determine the change in ecological variables resulting from exotic species. Hedges' g was calculated for 1,111 observations from 159 studies at 151 different sites

¹Red Sea Research Centre, King Abdullah University of Science and Technology (KAUST), Thuwal, Saudi Arabia. ²School of Biological Sciences, The University of Queensland, Brisbane, Queensland, Australia. ³Institute of Oceanography, Hellenic Center for Marine Research, Heraklion, Greece.

⁴Global Change Research Group, Institut Mediterrani d'Estudis Avançats (IMEDEA), CSIC-UIB, Esporles, Spain. ⁵Dauphin Island Sea Laboratory, Marine Environmental Sciences Consortium, Dauphin Island, AL, USA. ⁶Department of Marine Sciences, University of South Alabama, Mobile, AL, USA.

⁷Northern Gulf Institute, Mississippi State University, Stennis Space Center, MS, USA. ⁸Arctic Research Centre, Department of Bioscience, Aarhus University, Aarhus, Denmark. ⁹Department of Bioscience, Aarhus University, Silkeborg, Denmark. ¹⁰Laboratorio de Ecología, Instituto de Investigaciones Marinas y Costeras (IIMyC) CONICET-UNMdP, Mar de Plata, Argentina. ¹¹Australian Research Council Centre of Excellence for Coral Reef Studies, School of Biological Sciences, The University of Queensland, Brisbane, Queensland, Australia. ¹²Computational Bioscience Research Center, King Abdullah University of Science and Technology (KAUST), Thuwal, Saudi Arabia. *e-mail: andrea.antongamazo@kaust.edu.sa

Table 1 | List of the 76 exotic species included in our database by alphabetical order

Exotic species	Taxonomic group	Trophic level	IUCN	Observations	Hedges' <i>g</i>	<i>P</i> value
<i>Ammophila arenaria</i> (56) ⁷³	Plant	P. producer	No	1, 4	0.15 ± 0.83	0.714
<i>Arcuatula senhousia</i> ^a (62) ^{74–77}	Mollusc	Herbivore	No	4, 27	0.21 ± 0.45	0.347
<i>Avicennia marina</i> (38) ⁷⁴	Plant	P. producer	No	1, 3	−0.06 ± 0.98	0.893
<i>Avrainvillea amadelpha</i> (68) ⁷⁸	Algae	P. producer	No	1, 1	0.32 ± 1.24	0.606
<i>Baccharis halimifolia</i> (42) ⁷⁹	Plant	P. producer	No	1, 6	−0.03 ± 0.74	0.926
<i>Balanus glandula</i> (40) ⁸⁰	Crustacean	Omnivore	No	1, 4	−0.05 ± 0.84	0.904
<i>Batillaria attramentaria</i> (50) ⁸¹	Mollusc	Detritivore	No	1, 2	0.06 ± 0.96	0.893
<i>Boccardia proboscidea</i> (1) ⁸²	Annelid	Omnivore	No	1, 12	−1.67 ± 0.91	<0.001
<i>Botrylloides violaceus</i> (51) ⁸³	Tunicate	Herbivore	No	1, 9	0.08 ± 0.82	0.848
<i>Brachidontes pharaonis</i> (9) ⁸⁴	Mollusc	Herbivore	No	1, 6	−0.72 ± 0.73	0.053
<i>Carcinus maenas</i> ^a (10) ^{85–91}	Crustacean	Predator	Yes	7, 32	−0.71 ± 0.38	<0.001
<i>Carex macrocephala</i> (42) ^{92,93}	Plant	P. producer	No	2, 11	−0.04 ± 0.5	0.867
<i>Carpobrotus edulis</i> ^a (31) ^{94–96}	Plant	P. producer	No	3, 12	−0.14 ± 0.49	0.555
<i>Casuarina equisetifolia</i> (45) ⁹⁷	Plant	P. producer	No	1, 1	−0.01 ± 0.87	0.981
<i>Caulacanthus ustulatus</i> (63) ⁹⁸	Algae	P. producer	No	1, 18	0.23 ± 0.74	0.531
<i>Caulerpa cylindracea</i> ^a (12) ^{99–108}	Algae	P. producer	No	10, 38	−0.63 ± 0.31	<0.001
<i>Caulerpa taxifolia</i> ^a (23) ^{109–115}	Algae	P. producer	Yes	7, 40	−0.3 ± 0.34	0.081
<i>Centrostephanus rodgersii</i> (13) ^{116,117}	Echinoderm	Herbivore	No	2, 23	−0.62 ± 0.63	0.051
<i>Chrysanthemoides monilifera</i> (2) ¹¹⁸	Plant	P. producer	No	1, 1	−1.3 ± 1.16	0.028
<i>Ciona savignyi</i> (22) ⁸³	Tunicate	Herbivore	No	1, 8	−0.33 ± 0.84	0.439
<i>Cirolana harfordi</i> (52) ¹¹⁹	Crustacean	Predator	No	1, 8	0.08 ± 0.81	0.846
<i>Codium fragile</i> ^a (37) ^{9,120–123}	Algae	P. producer	No	5, 21	−0.07 ± 0.41	0.716
<i>Corbicula fluminea</i> (65) ^{124,125}	Mollusc	Herbivore	No	2, 17	0.28 ± 0.59	0.347
<i>Corbula gibba</i> (36) ¹²⁶	Mollusc	Herbivore	No	1, 4	−0.08 ± 1.07	0.878
<i>Crepidula fornicata</i> (19) ^{127–129}	Mollusc	Herbivore	No	3, 9	−0.46 ± 0.52	0.082
<i>Didemnum vexillum</i> ^a (57) ^{83,130,131}	Tunicate	Herbivore	No	3, 16	0.17 ± 0.53	0.52
<i>Diplosoma listerianum</i> (72) ¹³²	Tunicate	Herbivore	No	1, 4	0.53 ± 0.85	0.221
<i>Elymus athericus</i> (74) ¹³³	Plant	P. producer	No	1, 4	0.58 ± 0.93	0.216
<i>Ficopomatus enigmaticus</i> ^a (30) ^{134–140}	Annelid	Omnivore	No	7, 54	−0.15 ± 0.32	0.345
<i>Gracilaria salicornia</i> (53) ¹⁴¹	Algae	P. producer	No	1, 2	0.1 ± 0.93	0.825
<i>Gracilaria vermiculophylla</i> ^a (60) ^{142–145}	Algae	P. producer	No	4, 41	0.18 ± 0.41	0.368
<i>Grateloupia turuturu</i> (8) ¹⁴⁶	Algae	P. producer	No	1, 8	−0.75 ± 0.84	0.083
<i>Hemigrapsus sanguineus</i> ^a (4) ^{90,147,148}	Crustacean	Omnivore	No	3, 9	−0.81 ± 0.65	0.015
<i>Holcus lanatus</i> (39) ¹⁴⁹	Plant	P. producer	No	1, 13	−0.05 ± 0.76	0.887
<i>Juncus acutus</i> (18) ¹⁵⁰	Plant	P. producer	No	1, 8	−0.47 ± 0.82	0.263
<i>Kappaphycus alvarezii</i> (7) ¹⁵¹	Algae	P. producer	No	1, 4	−0.76 ± 0.78	0.057
<i>Laguncula pulchella</i> (25) ¹⁵²	Mollusc	Predator	No	1, 9	−0.22 ± 0.74	0.556
<i>Linepithema humile</i> (27) ¹⁵³	Insect	Omnivore	No	1, 3	−0.18 ± 1.01	0.726
<i>Lophocladia lallemandii</i> ^a (54) ^{103,104,154}	Algae	P. producer	No	3, 16	0.12 ± 0.47	0.591
<i>Loxothylacus panopaei</i> (16) ^{155,156}	Crustacean	Predator	No	2, 3	−0.48 ± 0.92	0.304
<i>Magallana gigas</i> ^a (59) ^{59,157–164}	Mollusc	Herbivore	No	9, 143	0.18 ± 0.26	0.175
<i>Maoricolpus roseus</i> (69) ¹⁶⁵	Mollusc	Herbivore	No	1, 4	0.33 ± 0.91	0.467
<i>Marenzelleria arctica</i> (75) ¹⁶⁶	Annelid	Detritivore	No	1, 2	0.77 ± 0.98	0.126
<i>Marenzelleria</i> spp. ^a (70) ^{167–169}	Annelid	Detritivore	No	3, 10	0.51 ± 0.67	0.136
<i>Megabalanus coccopoma</i> (20) ¹⁷⁰	Crustacean	Omnivore	No	1, 2	−0.36 ± 1.35	0.599
<i>Membranipora membranacea</i> (15) ¹⁷¹	Bryozoan	Omnivore	No	1, 1	−0.51 ± 1.44	0.482
<i>Mnemiopsis leidyi</i> (33) ¹⁷²	Ctenophora	Predator	Yes	1, 3	−0.14 ± 0.75	0.713
<i>Mytella charruana</i> (14) ¹⁷⁰	Mollusc	Herbivore	No	1, 2	−0.55 ± 1.36	0.429
<i>Mytilopsis trautwineana</i> (29) ¹⁷³	Mollusc	Herbivore	No	1, 3	−0.16 ± 0.93	0.73
<i>Mytilus galloprovincialis</i> ^a (17) ^{174–177}	Mollusc	Herbivore	Yes	4, 45	−0.48 ± 0.4	0.02

Continued

Exotic species	Taxonomic group	Trophic level	IUCN	Observations	Hedges' <i>g</i>	<i>P</i> value
<i>Perna viridis</i> (11) ¹⁷⁰	Mollusc	Herbivore	No	1, 2	-0.69 ± 1.37	0.322
<i>Petrolisthes armatus</i> (73) ¹⁷⁸	Crustacean	Omnivore	No	1, 4	0.53 ± 0.86	0.223
<i>Phragmites australis</i> (26) ¹⁷⁹	Plant	P. producer	No	1, 3	-0.19 ± 0.75	0.602
<i>Poecilia latipinna</i> (48) ¹⁸⁰	Fish	Omnivore	No	1, 1	0.01 ± 1.45	0.981
<i>Potamocorbula amurensis</i> (55) ¹⁸¹	Mollusc	Herbivore	Yes	1, 2	0.14 ± 1.21	0.815
<i>Potamopyrgus antipodarum</i> (43) ¹⁸²	Mollusc	Herbivore	No	1, 8	-0.02 ± 0.83	0.951
<i>Pterois volitans</i> ^a (35) ^{183–188}	Fish	Predator	No	6, 49	-0.12 ± 0.34	0.488
<i>Rattus norvegicus</i> (66) ⁴³	Mammal	Omnivore	No	1, 11	0.28 ± 0.75	0.46
<i>Rhizophora mangle</i> (76) ¹⁸⁹	Plant	P. producer	No	1, 12	1.15 ± 0.88	0.01
<i>Ruditapes philippinarum</i> (47) ¹⁹⁰	Mollusc	Herbivore	No	1, 3	0 ± 1.16	0.998
<i>Sabella spallanzanii</i> (64) ^{191,192}	Annelid	Omnivore	No	2, 8	0.28 ± 0.65	0.395
<i>Sagartia ornata</i> (67) ¹⁹³	Cnidarian	Predator	No	1, 2	0.3 ± 0.94	0.532
<i>Sargassum muticum</i> ^a (46) ^{194–204}	Algae	P. producer	No	11, 108	0 ± 0.25	0.966
<i>Sonneratia apetala</i> (58) ²⁰⁵	Plant	P. producer	No	1, 2	0.18 ± 1.22	0.773
<i>Spartina alterniflora</i> ^a (34) ^{206–216}	Plant	P. producer	No	11, 75	-0.13 ± 0.27	0.352
<i>Spartina townsendii</i> var. <i>anglica</i> (6) ^{217,218}	Plant	P. producer	Yes	2, 11	-0.77 ± 0.59	0.01
<i>Stenotaphrum secundatum</i> (5) ^{219,220}	Plant	P. producer	No	2, 7	-0.78 ± 0.56	0.006
<i>Styela clava</i> (61) ⁸³	Tunicate	Herbivore	No	1, 1	0.19 ± 1.43	0.792
<i>Symbiodinium trenchii</i> (28) ²²¹	Algae	P. producer	No	1, 6	-0.16 ± 0.9	0.713
<i>Tubastraea coccinea</i> (32) ²²²	Cnidarian	Predator	No	1, 12	-0.14 ± 0.72	0.69
<i>Undaria pinnatifida</i> ^a (49) ^{223–225}	Algae	P. producer	Yes	3, 34	0.05 ± 0.46	0.813
<i>Watersipora subtorquata</i> (71) ¹⁷⁵	Bryozoan	Omnivore	No	1, 4	0.51 ± 0.84	0.231
<i>Womersleyella setacea</i> (24) ^{108,154}	Algae	P. producer	No	2, 5	-0.26 ± 0.78	0.508
<i>Xenostrobus securis</i> (44) ²²⁶	Mollusc	Herbivore	No	1, 7	-0.01 ± 0.79	0.966
<i>Zeacumantus subcarinatus</i> (21) ²²⁷	Mollusc	Herbivore	No	1, 6	-0.33 ± 0.91	0.467
<i>Zostera japonica</i> (3) ⁶⁸	Plant	P. producer	No	1, 2	-0.85 ± 1.16	0.149

^aThe 19 sufficiently assessed marine exotic species (with 3 or more studies represented in Fig. 5). The studies used in this meta-analysis for each exotic species are indicated in the first column, and the rank of the exotic species based on their mean Hedges' *g* value is in parentheses. The 'Observations' column includes the number of studies and the number of observations, separated by a comma.

Values in the 'Hedges' *g*' column indicate the mean $g \pm$ CI. *P* values were obtained from mixed-effects models that included the study ID number as a random factor and using the *rma.mv* function from the *metafor* package for R. P., primary.

(Table 1; Supplementary Figs. 1 and 2; Supplementary Table 2; and <https://doi.pangaea.de/10.1594/PANGAEA.895681>). This meta-analysis allowed us to achieve the following: (1) determine the ecological effects of marine exotic species across trophic levels, taxonomic groups and ecological complexity; (2) elucidate the role of ecological characteristics such as the environment of the exotic species (for example, freshwater versus marine versus terrestrial), their mobility (mobile versus sessile), the geographical attributes of the native habitat (for example, continent versus island) and latitude on the ecological effects of marine exotics; and (3) create a unique ranking of marine exotic species based on their overall mean quantitative ecological effects.

Invasion ecology is a discipline that has complex terminology^{32–34}; therefore, to aid clarity, we provide an explicit definition of the terms used in this study. The term 'exotic' refers to species that have been moved outside their native geographical ranges via human actions. The definition of 'invasive species' varies between and among^{33,35} ecologists, stakeholders and policymakers. For many ecologists^{34,36}, invasive species are established exotic species that are spreading in the new environment (see also ref. ³⁶), while for most environmental agencies, it refers to exotic species that have a demonstrable ecological or economic impact (for example, American Federal Law). We refer to invasive species or 'pests' as exotic species that exert an overall and statistically significant ecological impact^{33,36}. Although this definition relies on human valuations of damage³⁴, we use it because of the following reasons: (1) we lack quantitative evidence

to indicate that the populations of the exotic species in our study are spreading beyond its initial establishment location (or locations); (2) it is followed by many ecologists studying invasive species^{33,35}; and (3) it tunes in with the language used by policymakers and environmental managers, providing effective guidance to meet Aichi Biodiversity Target 9. We use the term 'ecological effect' to refer to bidirectional ecological changes (for example, both increases and decreases) attributed to exotic species with regard to a control. The term 'ecological impact' is reserved only for statistically significant ecological changes (that is, mean changes significantly different from 0). Therefore, negative and positive effect size values are decreases and increases in ecological properties, respectively, without implying beneficial or deleterious effects.

Results

Our findings indicate that exotic species overall significantly reduce the ecological properties of native marine communities (mean estimated Hedges' $g = -0.14 \pm 0.1$ ($P = 0.006$); hereafter, mean effect sizes are reported as $g \pm 95\%$ confidence intervals (CI) and *P* values for significance). Ecological changes on marine ecosystems were quantified for 76 exotic species, which constitute 6% of the 1,260 marine and brackish exotic species listed in the Global Register of Introduced and Invasive Species (GRIIS; <http://www.griis.org/>). The ecological effect of 49 of these 76 exotic species was quantified in only one study each. However, we provide evidence to indicate that our global results are robust against both the number of

Fig. 1 | Effect of marine exotic species depends on the trophic levels of the native species or community and the exotic species. a,b, The mean effect size (Hedges' $g \pm 95\%$ CI) of the effects of marine exotic species depends on the trophic levels of the native species or community (a) and the exotic species (b). The unbroken line denotes zero and the broken line indicates the overall mean effect size. The number of observations is specified in parentheses, and * $P \leq 0.05$. P values were obtained from mixed-effects models that included the study ID and the exotic species name as random factors, using the `rma.mv` function from the `metafor` package for R.

exotic species and the number of independent studies based on a comparison of the mean $\pm 95\%$ CI Hedges' g value for species whose effects were assessed in one or more studies (which included the whole dataset) and for exotic species that included five or more independent studies (nine exotic species in total) (Supplementary Fig. 3). We show that while the effects of individual exotic species not included in this meta-analysis (that is, on our list of 76 species) may vary, the mean $\pm 95\%$ CI Hedges' g value for marine exotics is robust for the number of species included. There were small differences in the mean $g \pm CI$ for the 19 sufficiently assessed exotic species (which uses a subset of the data) and Table 1 (which uses the full dataset) due to differences in the effects of the random factor. The results for the 19 sufficiently assessed exotic species were more robust and given preference when interpreting the results. We found that the research effort on exotic species in marine ecosystems is not balanced; the GRIIS list of marine exotic species includes <10% of primary producers, which were the subject of almost 50% of our observations (517 out of 1,111).

Effects of trophic level and taxonomic group. Overall, exotic species significantly decreased the ecological properties of native primary producers and multitrophic assemblages ($g = -0.19 \pm 0.15$

($P = 0.014$) and $g = -0.17 \pm 0.13$ ($P = 0.016$), respectively) but not of native detritivores ($g = 0.02 \pm 0.27$ ($P = 0.870$)), herbivores ($g = -0.12 \pm 0.15$ ($P = 0.108$)), omnivores ($g = -0.15 \pm 0.16$ ($P = 0.051$)) or predators ($g = -0.14 \pm 0.19$ ($P = 0.143$)) (Fig. 1a). Among the exotic species, marine predators and primary producers were the only trophic groups that exerted significant decreases on marine ecological properties ($g = -0.28 \pm 0.27$ ($P = 0.049$) and -0.17 ± 0.13 ($P = 0.011$), respectively), but not herbivores, detritivores or omnivores ($g = -0.07 \pm 0.18$ ($P = 0.438$), $g = 0.40 \pm 0.58$ ($P = 0.173$) and $g = -0.15 \pm 0.27$ ($P = 0.269$), respectively) (Fig. 1b). Finally, the effect of marine exotic filter-feeder species on native communities was not significant ($g = -0.05 \pm 0.16$ ($P = 0.513$)) (Supplementary Fig. 4).

The network analysis of the ecological effects between exotic and native marine species among trophic and taxonomic groups was dominated by decreases in ecological properties (for example, almost 65% and 54% of the total effects, respectively) (Fig. 2). Exotic primary producers and predators caused ecological decreases in almost all trophic levels (Fig. 2). Exotic algae, crustaceans and molluscs were the taxonomic groups that caused overall decreases in the largest number of native taxa (seven, six and five, respectively). Meanwhile, exotic terrestrial mammals, annelids, molluscs and tunicates had the largest number of increasing effects (six, five, five and five, respectively) (Fig. 2).

Changes at species, community and ecosystem levels. Seven out of the ten ecological categories of response variables were not significantly affected by exotic species. Traits related to the behaviour and abundance of native species were significantly reduced by exotic species ($g = -0.53 \pm 0.28$ ($P < 0.001$) and $g = -0.26 \pm 0.10$ ($P < 0.001$), respectively) (Fig. 3). Conversely, ecological traits related to pollution significantly increased ($g = 0.44 \pm 0.41$ ($P = 0.033$)) (Fig. 3). There was no significant effect of exotic species on the following factors: fitness ($g = 0.25 \pm 0.39$ ($P = 0.203$)), growth ($g = -0.11 \pm 0.35$ ($P = 0.544$)), survival ($g = -0.06 \pm 0.23$ ($P = 0.601$)), richness ($g = -0.06 \pm 0.12$ ($P = 0.344$)), biogeochemical elements ($g = 0.20 \pm 0.24$ ($P = 0.108$)), rate processes ($g = 0.08 \pm 0.25$ ($P = 0.511$)) or sediment changes ($g = 0.26 \pm 0.43$ ($P = 0.231$)) (Fig. 3). In general, the ecological effects of exotics at the community level were significant ($g = -0.19 \pm 0.10$ ($P < 0.001$)), while changes at the species and ecological levels were not ($g = -0.15 \pm 0.16$ ($P = 0.084$) and $g = 0.14 \pm 0.19$ ($P = 0.138$), respectively) (Fig. 3).

Context-dependency of ecological effects. Exotics of marine origin caused significant decreases in the ecological variables of marine systems ($g = -0.14 \pm 0.11$ ($P = 0.010$)), while exotic species of freshwater and terrestrial origin did not ($g = -0.05 \pm 0.37$ ($P = 0.795$) and $g = -0.17 \pm 0.31$ ($P = 0.292$), respectively) (Fig. 4a). Sessile exotic species had significant ecological effects on native marine ecosystems ($g = -0.14 \pm 0.11$ ($P = 0.016$)) and mobile species did not ($g = -0.12 \pm 0.22$ ($P = 0.205$)) (Fig. 4b). We found that the ecological effects of exotic species in marine ecosystems were significant in continental margins ($g = -0.17 \pm 0.11$ ($P = 0.003$)) but not on islands ($g = -0.05 \pm 0.18$ ($P = 0.581$)) (Fig. 4c). There was no indication that the ecological effect of marine exotic species varied over latitude ($P = 0.849$ and $P = 0.536$, for the Northern and Southern hemispheres, respectively) (Fig. 4d).

Ranking and significance of exotic species based on their ecological effects. We identified eight exotic species (*B. proboscidea*, *C. maenas*, *C. cylindracea*, *C. monilifera*, *H. sanguineus*, *M. galloprovincialis*, *S. anglica* and *S. secundatum*) that caused overall significant decreases (mean $g < 0$ and 95% CI did not overlap 0; Table 1). Meanwhile, one species caused significant increases (*R. mangle*), and 67 species did not cause significant ecological changes (Table 1). A subset of 19 exotic species, whose effects were

Fig. 2 | Network diagram of the ecological effects of marine exotics on native species classified by trophic level and taxonomic group. The width of the band represents the mean effect size Hedges' g , where effect sizes that increase or decrease ecological properties are presented in the upper and lower panels, respectively. The insect group includes arachnids. Interactive diagrams are available at <https://kaust-vislab.github.io/Biological-Network/>.

assessed in three or more independent studies, was chosen to produce a robust ranking of marine exotic species based on their ecological effects. In this ranking of sufficiently assessed exotic species, only the European green crab *C. maenas* and the green macroalgae *C. cylindracea* had overall significant impacts (Fig. 5).

Discussion

Our findings indicate that globally, exotic species induce significant changes on many ecological attributes of marine systems ($g = -0.14 \pm 0.1$). This global effect is modest in magnitude compared to other quantified human stressors, such as the overall impact of dual environmental stressors in freshwater ecosystems ($g = -0.64 \pm 0.46$, as calculated from data in ref. ³⁷ after removing the effects of invasive species) or the worldwide effects of fishing pressure on reef fish behaviour ($g = 1.3 \pm 0.13$; values obtained from figure 3 in ref. ³⁸). The global impact of marine exotic species is comparable to the overall effect reported for freshwater exotic species worldwide ($g = -0.27 \pm 0.25$; values calculated from raw data available in ref. ²³). However, it is larger than the nonsignificant ecological effects reported for exotic ecosystem engineers on the

abundance or richness of marine native species ($g = 0.04 \pm 0.19$ and 0.06 ± 0.26 , respectively)²⁴.

We provide a quantitative synthesis of ecological effects across both trophic levels and taxonomic groups, while previous meta-analyses evaluated the effects of exotic species on certain native trophic levels^{22,23,25} or taxonomic groups^{21,24–26}. Overall, exotic species had significant effects on primary producers and multitrophic assemblages, while the other native trophic levels experienced overall contrasting effects. Native primary producers were affected by all trophic levels of exotics, which surpassed our expectations drawn from freshwater systems²³ in that they were also affected by exotic predators and detritivores. Exotic predators and primary producers were the only trophic groups that exerted substantial impacts on marine ecological properties; these trophic groups were therefore identified as the most disruptive of all marine exotics. Exotic predators, arguably the most detrimental group for terrestrial and freshwater native communities^{6,7}, caused overall decreases in the ecological properties of all marine trophic groups except detritivores. Similarly, exotic primary producers caused ecological decreases at most trophic levels and taxonomic groups, which is

Fig. 3 | Mean effect size (Hedges' $g \pm 95\%$ CI) of marine exotic species on response variables classified by the levels of ecological complexity. The unbroken line denotes zero, and the broken line indicates the overall mean effect size. The number of observations is specified in parentheses, and * $P \leq 0.05$, *** $P \leq 0.001$. P values were obtained from mixed-effects models that included the study ID and exotic species name as random factors, using the `rma.mv` function from the `metafor` package for R.

in contrast to previous studies that reported contrasting effects of exotic primary producers in freshwater²³ and marine systems²². Conversely, exotic marine herbivores, detritivores and omnivores, of which previous information is scarce²³, had contrasting effects on other trophic levels of the food web. These contrasting effects were probably context-dependent, with changes compensating for each other by going in different directions.

Exotic molluscs and crustaceans, for which no information from previous meta-analyses was available to make predictions, were the other two taxonomic groups that induced overall ecological decreases in many native marine taxa. The findings for exotic mammals, the taxon that caused increases in the largest number of native taxa, were surprising since they are a very disruptive group in terrestrial ecosystems^{5,6,39,40} as well as in some marine coastal communities^{41,42}. This result is derived from one publication⁴³, where the exotic brown rat (*R. norvegicus*) triggered a trophic cascade by directly preying on birds but indirectly enhancing the abundance of many benthic taxa. Further investigation is required to test the generality of this finding.

Exotic species significantly decreased the abundance of native species but not diversity, as reported in freshwater systems²³. Marine exotics might prompt rapid declines in species abundance, but they may take a long time to generate local extinctions (an 'extinction debt'⁴⁴). Our decoupled results between decreases in

Fig. 4 | Mean effect size (Hedges' $g \pm 95\%$ CI) of marine exotic species based on different variables. a–d. Effects were based on the environment of exotic species (a), mobility of exotic species (b), if the study location was a continent or an island (c) and the latitude of location (d). The unbroken line denotes zero and the broken line indicates overall mean effect size. The number of observations is specified in parentheses, and * $P \leq 0.05$, ** $P \leq 0.01$. P values were obtained from mixed-effects models that included the study ID and exotic species name as random factors, using the `rma.mv` function from the `metafor` package for R.

abundance but not overall changes on ecosystem functioning are comparable to those reported for exotics in freshwater²³ and terrestrial systems²¹, and by other anthropogenic factors elsewhere⁴⁵. This indicates that changes in community composition are often decoupled from those at the ecosystem level. Possible buffering mechanisms conferring ecosystem resistance against exotic species in the marine realm include functional redundancy between exotic and native species or between affected and non-affected native species⁴⁶. Our results differ from those reported for marine exotic ecosystem engineers, which exert contrasting strong effects on native ecosystem functioning but not on native community abundance²⁴. The contrasting results among studies might be due to targeting different suites of marine exotic species and clustering ecological response variables differently. Interestingly, we found a lack of ecological effects on survival, growth and fitness of native species, but profound changes in animal behaviour. Changes in behaviour of native species could affect native species populations in the short term and, where adaptation of behavioural traits is possible, buffer the effects in the long term⁴⁷. We found significant increases in ecological traits related to pollution, mainly through the increased concentration of heavy metals in the tissues of the exotic species compared with native species.

Relating features of exotic species with their ecological effects is of utmost importance if we aim to control thousands of exotic species⁴⁸. Sessile exotics or those with a marine origin exerted significant

Fig. 5 | Ranking of the 19 sufficiently assessed marine exotic species based on the species mean effect size. Effect size indicates an increase ($g > 0$) or decrease ($g < 0$) with respect to a reference control. Bars represent 95% CI, which denote significant effects when they do not overlap zero. The unbroken line represents zero, and the broken line indicates the overall mean effect size. The number of studies (left) and the number of observations (right) are specified in parentheses, and $^{**}P \leq 0.01$, $^{***}P \leq 0.001$. P values were obtained from mixed-effects models that included the study ID as a random factor, using the `rma.mv` function from the `metafor` package for R. Red squares indicate significant ecological effects of exotic species, which would be categorized as invasive species³³ or pests³⁶. Circles indicate nonsignificant effects of exotic species, with black and white filled circles indicating mean $g < 0$ or $g > 0$, respectively. Credit: Allende Bodega Martinez (silhouettes).

changes on marine native communities, while mobile exotics or those with a freshwater or terrestrial origin did not. This result therefore identifies important characteristics of potentially detrimental exotic species. Environmental and geographical attributes of the invaded communities can also influence the establishment and success of exotic species. We found that the ecological effects of exotic species on marine ecosystems were substantial in continental margins but

not on islands, which was unexpected based on reports for exotic avian species⁴⁰ or terrestrial plants²¹. There was no clear indication that the ecological effects of marine exotic species varied over latitude, although we anticipated low effects at low latitudes due to their high biodiversity, low space availability and strong biotic interactions^{49,50}. This finding has two possible explanations. First, although benthic biodiversity is highest at tropical latitudes for many marine

habitats (for example, coral reefs, seagrass meadows or mangrove forests), this does not apply to all marine ecosystems (for example, macroalgae beds and saltmarshes)^{51,52}. Second, there were a few tropical and polar study locations (ten and one, respectively), which might have influenced the results.

About 10% of marine exotic species were qualified as invasive species (as defined in ref. ³³) or pests (as defined in ref. ³⁶) by causing overall significant decreases in ecological variables. This percentage is similar to the proportion of exotic species that had been previously estimated to have sizeable impacts on native communities^{36,48,53,54}. However, these former estimates were based on semiquantitative data, unlike our estimate, which was based on a common metric, thus enabling a rigorous evaluation across species and types of impacts⁵⁵. Four species (*C. maenas*, *C. taxifolia*, *M. galloprovincialis* and *U. pinnatifida*) were included in the world's 100 most invasive species by the IUCN⁵⁶. While the overall ecological impact of *C. maenas* was obvious, those of *C. taxifolia*, *M. galloprovincialis* and *U. pinnatifida* are of concern since the wide error bars around the mean indicate that they can exert large impacts. The IUCN list⁵⁶, the most reputable compilation of the worst exotic species worldwide, was based on expert judgment, and our list is an upgrade for marine exotics⁵⁷. By separating the disruptive from the innocuous species, we provide a basis for prioritizing and triaging eradication and management efforts, thereby offering a solid foundation to help meet Aichi Biodiversity Target 9 of the international Convention on Biological Diversity.

Our ranking of exotic species is based on the magnitude of the mean global effect for each exotic species. However, it is important to consider not only the mean but also the variability of the effects. For instance, exotic species that inflict strong disparate impacts depending on geographical locations or habitat types^{58,59} will have large error bars around the mean, which was the case for some exotic species on our list. Therefore, many of the exotic species on our list could qualify as invasive in some locations but not others, although their global effect might not be significant. The lowest confidence interval (that is, the largest ecological decrease) could therefore be used to rank the species under a precautionary approach. The resulting ranking would, however, be very similar to that resulting when using the mean *g*.

The effects summarized here may be biased towards the most damaging of exotic species because studies with negligible ecological effects are usually underreported in the literature⁶⁰. This is probably because exotic species present at low abundances might go unnoticed or are difficult to study. In addition, quantitative analyses of the effects of exotic species are often funded and initiated when detrimental impacts are suspected, thereby probably targeting the most aggressive exotics. Therefore, our quantified ecological effects of marine exotic species may reflect a 'worst case' scenario, but one that is relevant to inform management actions under a precautionary principle.

Recent assessments indicating that marine exotic species, based on ecosystem engineer species, have overall nonsignificant effects²⁴ are challenged when a broader suite of species is assessed, revealing significant ecological changes of exotic species at a global scale. Hence, concerns about the effects of marine exotics are warranted¹⁰, especially since they will probably intensify as the vectors of introduction of marine exotics are expanding and diversifying^{61,62}. However, our analysis also partially supports tempered views on the effects of marine exotics⁶³ through the finding that the overall changes in magnitude are moderately significant ($g < 0.2$)⁶⁴ and that there is a prevalence of overall nonsignificant effects for the majority of exotic species (~90%).

In conclusion, the results provided here offer an integrated, quantitative assessment of the effects of exotic species, enabling the reconciliation of the two polarized sides of the current debate. We also provide a ranked assessment of marine exotic species based on their

quantitative ecological effects on native communities. Together, these results contribute to efforts to meet Aichi Biodiversity Target 9 of the Convention on Biological Diversity of the United Nations, and reveal previously unknown ecological patterns, such as the trend of exotic marine predators and primary producers to cause the largest ecological impacts.

Methods

Literature search. Quantitative data from the literature were gathered to assess the ecological effects of marine exotic species. We performed an exhaustive and systematic review on 4 May 2016 following the guidelines of TOP⁶⁵ (transparency and openness promotion) and PRISMA⁶⁶ (preferred reporting items for meta-analyses) (Supplementary Table 3; Supplementary Fig. 1). We used the Web of Science (Thomson Reuters) search engine to find studies using the following terms: (invader OR exotic OR invasive OR non-native OR alien) NEAR/5 (impact OR effect OR influence OR consequence*) AND (marine OR coastal OR sea OR estuar* OR ocean) NOT (lake OR stream OR freshwater OR terrestrial). The search resulted in 1,012 publications. Each reference was randomly assigned to authors, who each extracted information from a set of papers.

Study selection criteria and data entering. The following criteria were used to select studies to include in the analysis. First, we only included studies within marine systems, up to and including environments getting sea spray. These comprised exotic species that inhabit brackish-water estuaries (such as the snail *P. antipodarum*) or that cover the coastal zones (such as the succulent terrestrial plant *C. edulis*). We included freshwater and terrestrial exotic species reported to interact with native marine species. Indeed, some of these species have been described as potential threats to coastal marine communities, such as the exotic terrestrial American mink (*Mustela vison*), which can deplete many species of European native fish and crabs in the rocky intertidal⁴². Second, we included species (for example, exotic species or species from the native community) that spend any stage of their life cycle in the marine environment. Third, we only included studies that quantitatively compared control and experimental treatments (for example, exotic species removals versus additions, or exotic species versus native species treatments). Articles lacking appropriate controls were excluded. Fourth, we only included studies reporting the mean values of variables, number of replicates and a measure of variability around the mean. If these data could not be obtained from a paper but did appear to have been recorded, the authors of the original study were contacted to request the relevant information. Fifth, when multiple and distinct habitat types or exotic species were examined separately within the same article, they were entered in the database as separate observations (for example, each observation was a row in our database). Sixth, if an article reported data on studies conducted at several locations, they were included as independent studies. Seventh, if more than two densities of exotic species were compared in a study (for example, low, medium and high abundance of the exotic species), they were entered in the database as separate observations. Eighth, if response variables were measured at multiple time points, we recorded the first and last sampling events except if there was a clear pattern in the data that was missed by recording only the first and last sampling events. In those cases, an additional point in the middle was recorded. Ninth, if multiple sampling techniques were used to estimate impacts within one area, only results from the most efficient sampling method were included. Tenth, for studies reporting multiple native community measures (for example, abundance, biomass, richness and diversity) and data on individual native species, we extracted information on community abundance and richness and collected data for a maximum of two individual native species either from the most abundant species or from the native species that was the focus of the study. Eleventh, for studies that reported results for the same species at several life stages or sizes, they were entered in the database as separate observations.

From an initial selection of 1,012 published studies, 159 studies were found to contain suitable quantitative data from 151 study locations (Supplementary Table 2; Supplementary Figs. 1 and 2). We assessed the effects of exotic species on the following ten types of metrics (Supplementary Table 4): abundance, richness, biogeochemical elements (stocks or fluxes of carbon, nitrogen, phosphorus and silicate), rate processes (production, photosynthesis, respiration and decomposition), survival (including mortality), growth (including percentage growth), fitness, behaviour, pollution (including heavy metal concentration and water clarity), and sediment changes. We then grouped these response variables at the species, community and ecosystem levels (Supplementary Table 4). A dropdown list was provided to co-authors to standardize data entry and to exclude redundant measurements. The trophic level of the species was determined after performing data searches on the diet of each exotic species in the Web of Science database and in the Invasive Species Compendium in the Center for Agriculture and Biosciences International. Filter-feeders were categorized as follows: (1) bivalve molluscs, tunicates and suspension-feeding gastropods were categorized as herbivores because they consume primarily phytoplankton; and (2) barnacles, bryozoans and suspension-feeding annelids and crabs were categorized as omnivores because they are not restricted to microscopic algae for

food. The same criteria were followed to categorize the trophic level of the native species, for which, in addition, sponges were classified as omnivores. For exotic species names, we used the ones most recently listed in the World Register of Marine Species. Data were obtained from tables or extracted from plots using Graph Click (available at <http://www.arizona-software.ch/graphclick/download.html>).

Data analysis. Hedges' g effect size and variance were calculated for each observation (1,111 observations in total; see also <https://doi.pangaea.de/10.1594/PANGAEA.895681>) to estimate the differences in the response variable between control and experimental treatment. Hedges' g weighs cases by their sample size and the inverse of their variance⁶⁷. Hedges' g ranges from $-\infty$ to $+\infty$ and can be interpreted as follows⁶⁴: $|g| \leq 0.2$ considered a small effect; $0.2 \leq |g| \leq 0.5$ a medium effect; $0.5 \leq |g| \leq 0.8$ a large effect; and $|g| \geq 1.0$ a very large effect. The effect size Hedges' g was calculated as follows⁶⁷:

$$g = (X_E - X_C) / J \cdot s.d._{pooled}$$

where X_E and X_C are the mean of the experimental (for example, presence of exotic species) and control groups (for example, absence of exotic species), respectively. J corrects for bias attributed to different sample sizes by differentially weighting studies as follows:

$$J = 1 - (3 / (4 \times (N_E + N_C - 2) - 1))$$

The pooled standard deviation ($s.d._{pooled}$) was calculated as follows^{64,67}:

$$s.d._{pooled} = \sqrt{((N_E - 1) \times (s.d._E)^2 + (N_C - 1) \times (s.d._C)^2) / (N_E + N_C - 2)}$$

where $s.d.$ is the standard deviation of the experimental or control group and N is the sample size. We weighted the effect sizes to account for inequality in study variance by using the inverse of the sampling variance, in which the variance for each effect size (V_g) was calculated as follows⁶⁴:

$$V_g = ((N_E + N_C) / (N_E \times N_C)) + (g^2 / (2 \times (N_E + N_C)))$$

Meta-analyses were completed using the metafor package for R⁶⁹. We ran mixed-effects models that included the study identification (ID) number and exotic species name as random factors to account for the effect of the exotic species within a study and across studies (for example, when several observations for a species were obtained from the same publication or from several publications). For models with categorical fixed factors (for example, trophic level of native species, trophic level of exotic species, level of ecological complexity, type of ecological response variable, origin and mobility of exotic species, continentality and exotic species name), effect sizes were significant if the following criteria were met: (1) their 95% CI did not overlap with zero; and (2) P values were ≤ 0.05 . For the model with a continuous fixed factor (latitude), the predictor was considered to have significant effects at $P \leq 0.05$. In all statistical models, we used the `rma.mv` function, which determines statistical significance based on a Wald-type test.

Filter-feeders are a special guild that comprises several trophic levels and can have important ecological effects. We therefore performed an additional analysis on the effects of the trophic level of exotic species on native species including filter-feeders as a trophic level (27 exotic species were filter-feeders) to quantify their effects.

Publication bias. Publication bias, which is the selective publication of articles finding significant effects over those that find nonsignificant effects, might distort the results in a meta-analysis⁶⁴. In our case, publication bias could lead to an overestimate of the effects of exotic species in marine ecosystems. The functions `regtest` and `trimfill` are not implemented in the metafor package for mixed-effects models⁶⁹. Therefore, potential publication bias was evaluated using Egger's regression test⁷⁰ by running models that included the standard error ($s.e.$) of the effect sizes (included as the square root of the variance) as a moderator⁷¹. Potential publication bias was determined when the intercept of the model was different from zero at $P \leq 0.05$. If potential bias was detected, we examined the data for potential outliers by looking at the effect sizes with standardized residual values exceeding the absolute value of three⁷² using the `rstandard` function in R. Potential outliers were removed to adjust for publication bias. Adjusting for publication bias did not change the outcome of the analyses (by comparing fitted random-effects models with and without the influence of the potential outliers), except for both the effects of exotic species on native omnivores and the biogeochemical elements (Supplementary Table 5). We removed the potential outliers detected in the sensitivity analysis and re-ran the mixed-effects models for the effects of exotic species on the effects of trophic level (represented in Fig. 1) and the ecological categories of the response variables (represented in Fig. 3). Otherwise, our sensitivity analyses showed that our findings are robust against publication bias (Supplementary Table 5).

Standardization of variables. We performed the following calculations to standardize our dataset.

Variation around the mean. The $s.d.$ was used as the measure of variation. When another statistic was reported (for example, $s.e.$ or CI), it was converted to the $s.d.$ using the following calculations (where n is the number of replicates, $d.f.$ the degrees of freedom and α the probability of rejecting the null hypothesis):

$$s.d. = s.e. \times \sqrt{n}$$

$$CI = s.e. \times t_{\alpha, d.f.}$$

Growth data. If data were reported as biomass increase, we used the initial (B_0) and final (B_t) reported in the paper, and calculated the fractional growth (F_g):

$$F_g = [B_t / B_0]$$

The $s.d.$ of F_g ($s.d._{F_g}$) is then $s.d._{F_g} = F_g \times (s.d._{B_t} / B_t)$. When data were reported as the percentage growth (%G) with its $s.d.$ ($s.d._{\%G}$), we converted %G into F_g .

Mortality and survivorship data. If data were reported as initial and final number of individuals in the populations (N_0 and N_t , respectively), we calculated the fraction of survivors (F_s) as follows:

$$F_s = [N_t / N_0]$$

where N_t is the number of the survivors at time t and N_0 is the number of individuals at the beginning of the experiment.

The $s.d.$ of F_s ($s.d._{F_s}$) was calculated as follows:

$$s.d._{F_s} = F_s \times (s.d._{N_t} / N_t)$$

When data were reported as percentage survivorship (%S) with its $s.d.$ ($s.d._{\%S}$), we converted %S to F_s .

When the data were reported as number of dead and when the results were reported as percentage mortality, the calculations were performed as above, but converted from number of percentage dead into number of percentage surviving.

Reporting Summary. Further information on research design is available in the Nature Research Reporting Summary linked to this article.

Data availability

All data underlying the study have been deposited in PANGAEA at <https://doi.pangaea.de/10.1594/PANGAEA.895681>.

Code availability

The R script used in this manuscript will be deposited in the Github community repository upon publication (<https://github.com/ngerald/marine-exotics-global-analysis>).

Received: 2 June 2018; Accepted: 24 February 2019;

Published online: 08 April 2019

References

1. Simberloff, D. et al. Impacts of biological invasions: what's what and the way forward. *Trends Ecol. Evol.* **28**, 58–66 (2013).
2. Hulme, P. E., Pyšek, P., Nentwig, W. & Vilà, M. Will threat of biological invasions unite the European Union? *Science* **324**, 40–41 (2009).
3. Catford, J. A., Bode, M. & Tilman, D. Introduced species that overcome life history tradeoffs can cause native extinctions. *Nat. Commun.* **9**, 2131 (2018).
4. Seebens, H. et al. No saturation in the accumulation of alien species worldwide. *Nat. Commun.* **8**, 14435 (2017).
5. Medina, F. M. et al. A global review of the impacts of invasive cats on island endangered vertebrates. *Glob. Change Biol.* **17**, 3503–3510 (2011).
6. Doherty, T. S., Glen, A. S., Nimmo, D. G., Ritchie, E. G. & Dickman, C. R. Invasive predators and global biodiversity loss. *Proc. Natl Acad. Sci. USA* **113**, 11261–11265 (2016).
7. Witte, F. et al. The destruction of an endemic species flock—quantitative data on the decline of the haplochromine cichlids of Lake Victoria. *Environ. Biol. Fishes* **34**, 1–28 (1992).
8. Ramus, A. P., Silliman, B. R., Thomsen, M. S. & Long, Z. T. An invasive foundation species enhances multifunctionality in a coastal ecosystem. *Proc. Natl Acad. Sci. USA* **114**, 8580–8585 (2017).
9. Gerdali, N. R., Smyth, A. R., Piehler, M. F. & Peterson, C. H. Artificial substrates enhance non-native macroalga and N-2 production. *Biol. Invasions* **16**, 1819–1831 (2014).
10. Russell, J. C. & Blackburn, T. M. The rise of invasive species denialism. *Trends Ecol. Evol.* **32**, 3–6 (2017).
11. Davis, M. et al. Don't judge species on their origins. *Nature* **474**, 153–154 (2011).

12. Clergeau, P. & Nuñez, M. A. The language of fighting invasive species. *Science* **311**, 951–951 (2006).
13. Richardson, D. M. & Ricciardi, A. Misleading criticisms of invasion science: a field guide. *Divers. Distrib.* **19**, 1461–1467 (2013).
14. Valéry, L., Fritz, H. & Lefeuvre, J.-C. Another call for the end of invasion biology. *Oikos* **122**, 1143–1146 (2013).
15. Ricciardi, A. & Ryan, R. The exponential growth of invasive species denialism. *Biol. Invasions* **20**, 549–553 (2018).
16. Gurevitch, J. & Padilla, D. K. Are invasive species a major cause of extinctions? *Trends Ecol. Evol.* **19**, 470–474 (2004).
17. Clavero, M. & Garcia-Berthou, E. Invasive species are a leading cause of animal extinctions. *Trends Ecol. Evol.* **20**, 110–110 (2005).
18. Pyšek, P., Blackburn, T. M., García-Berthou, E., Perglová, I. & Rabitsch, W. in *Impact of Biological Invasions on Ecosystem Services* (eds Vilà, M. & Hulme, P. E.) 157–175 (Springer, 2017).
19. Bellard, C., Cassey, P. & Blackburn, T. M. Alien species as a driver of recent extinctions. *Biol. Lett.* **12**, 20150623 (2016).
20. Green, S. J., Akins, J. L., Maljković, A. & Côté, I. M. Invasive lionfish drive Atlantic coral reef fish declines. *PLoS ONE* **7**, e32596 (2012).
21. Vilà, M. et al. Ecological impacts of invasive alien plants: a meta-analysis of their effects on species, communities and ecosystems. *Ecol. Lett.* **14**, 702–708 (2011).
22. Maggi, E. et al. Ecological impacts of invading seaweeds: a meta-analysis of their effects at different trophic levels. *Divers. Distrib.* **21**, 1–12 (2015).
23. Gallardo, B., Clavero, M., Sánchez, M. I. & Vilà, M. Global ecological impacts of invasive species in aquatic ecosystems. *Glob. Change Biol.* **22**, 151–163 (2016).
24. Guy-Haim, T. et al. Diverse effects of invasive ecosystem engineers on marine biodiversity and ecosystem functions—a global review and meta-analysis. *Glob. Change Biol.* **24**, 906–924 (2018).
25. Molot, G., Pantel, J. H. & Romanuk, T. N. in *Advances in Ecological Research* Vol. 56 (eds Bohan, D. A., Dumbrell, A. J. & Massol, F.) Ch. 2 (Academic, 2017).
26. Thomsen, M. S. et al. Impacts of marine invaders on biodiversity depend on trophic position and functional similarity. *Mar. Ecol. Prog. Ser.* **495**, 39–47 (2014).
27. Cameron, E. K., Vilà, M. & Cabeza, M. Global meta-analysis of the impacts of terrestrial invertebrate invaders on species, communities and ecosystems. *Glob. Ecol. Biogeogr.* **25**, 596–606 (2016).
28. Thompson, R. M., Hemberg, M., Starzomski, B. M. & Shurin, J. B. Trophic levels and trophic tangles: the prevalence of omnivory in real food webs. *Ecology* **88**, 612–617 (2007).
29. Vergés, A. et al. Tropical rabbitfish and the deforestation of a warming temperate sea. *J. Ecol.* **102**, 1518–1527 (2014).
30. Sala, E., Kizilkaya, Z., Yildirim, D. & Ballesteros, E. Alien marine fishes deplete algal biomass in the Eastern Mediterranean. *PLoS ONE* **6**, e17356 (2011).
31. *The Strategic Plan for Biodiversity 2011–2020 and the Aichi Biodiversity Targets* UNEP/CBD/COP/DEC/X/2 2010 (UNEP, CBP, 2010).
32. Jeschke, J. M. et al. Defining the impact of non-native species. *Conserv. Biol.* **28**, 1188–1194 (2014).
33. Lockwood, J. L., Hoopes, M. F. & Marchetti, M. P. *Invasion Ecology* (Wiley-Blackwell, 2013).
34. Blackburn, T. M. et al. A proposed unified framework for biological invasions. *Trends Ecol. Evol.* **26**, 333–339 (2011).
35. Young, A. M. & Larson, B. M. H. Clarifying debates in invasion biology: a survey of invasion biologists. *Environ. Res.* **111**, 893–898 (2011).
36. Richardson, D. M. et al. Naturalization and invasion of alien plants: concepts and definitions. *Divers. Distrib.* **6**, 93–107 (2000).
37. Jackson, M. C., Loewen, C. J. G., Vinebrooke, R. D. & Chimimba, C. T. Net effects of multiple stressors in freshwater ecosystems: a meta-analysis. *Glob. Change Biol.* **22**, 180–189 (2016).
38. Nunes, J. A. C. C. et al. Global trends on reef fishes' ecology of fear: flight initiation distance for conservation. *Mar. Environ. Res.* **136**, 153–157 (2018).
39. Bellard, C., Genovesi, P. & Jeschke, J. M. Global patterns in threats to vertebrates by biological invasions. *Proc. R. Soc. B* **283**, 20152454 (2016).
40. Blackburn, T. M., Cassey, P., Duncan, R. P., Evans, K. L. & Gaston, K. J. Avian extinction and mammalian introductions on oceanic islands. *Science* **305**, 1955–1958 (2004).
41. Graham, N. A. J. et al. Seabirds enhance coral reef productivity and functioning in the absence of invasive rats. *Nature* **559**, 250–253 (2018).
42. Delibes, M., Clavero, M., Prenda, J., Blazquez, M. D. & Ferreras, P. Potential impact of an exotic mammal on rocky intertidal communities of northwestern Spain. *Biol. Invasions* **6**, 213–219 (2004).
43. Kurl, C. M., Croll, D. A. & Tershy, B. R. Introduced rats indirectly change marine rocky intertidal communities from algae- to invertebrate-dominated. *Proc. Natl Acad. Sci. USA* **105**, 3800–3804 (2008).
44. Tilman, D., May, R. M., Lehman, C. L. & Nowak, M. A. Habitat destruction and the extinction debt. *Nature* **371**, 65–66 (1994).
45. Anton, A. et al. Decoupled effects (positive to negative) of nutrient enrichment on ecosystem services. *Ecol. Appl.* **21**, 991–1009 (2011).
46. Garcia, D., Martinez, D., Stouffer, D. B. & Tylianakis, J. M. Exotic birds increase generalization and compensate for native bird decline in plant–frugivore assemblages. *J. Anim. Ecol.* **83**, 1441–1450 (2014).
47. Strauss, S. Y., Lau, J. A. & Carroll, S. P. Evolutionary responses of natives to introduced species: what do introductions tell us about natural communities? *Ecol. Lett.* **9**, 354–371 (2006).
48. Strayer, D. L. Eight questions about invasions and ecosystem functioning. *Ecol. Lett.* **15**, 1199–1210 (2012).
49. Freestone, A. L., Ruiz, G. M. & Torchin, M. E. Stronger biotic resistance in tropics relative to temperate zone: effects of predation on marine invasion dynamics. *Ecology* **94**, 1370–1377 (2013).
50. Hillebrand, H. On the generality of the latitudinal diversity gradient. *Am. Nat.* **163**, 192–211 (2004).
51. Saintilan, N. Biogeography of Australian saltmarsh plants. *Austral. Ecol.* **34**, 929–937 (2009).
52. Kerswell, A. P. Global biodiversity patterns of benthic marine algae. *Ecology* **87**, 2479–2488 (2006).
53. Vilà, M. et al. How well do we understand the impacts of alien species on ecosystem services? A pan-European, cross-taxa assessment. *Front. Ecol. Environ.* **8**, 135–144 (2010).
54. Ricciardi, A. & Kipp, R. Predicting the number of ecologically harmful exotic species in an aquatic system. *Divers. Distrib.* **14**, 374–380 (2008).
55. Kumschick, S. et al. Comparing impacts of alien plants and animals in Europe using a standard scoring system. *J. Appl. Ecol.* **52**, 552–561 (2015).
56. Lowe, S., Browne, M. & Boudjelas, S. *100 of the World's Worst Invasive Alien Species* (IUCN/SSC Invasive Species Specialist Group (ISSG), 2000).
57. Pauchard, A. et al. Biodiversity assessments: origin matters. *PLoS Biol.* **16**, e2006686 (2018).
58. South, J., Dick, J. T. A., McCard, M., Barrios-O'Neill, D. & Anton, A. Predicting predatory impact of juvenile invasive lionfish (*Pterois volitans*) on a crustacean prey using functional response analysis: effects of temperature, habitat complexity and light regimes. *Environ. Biol. Fishes* **100**, 1155–1165 (2017).
59. Green, D. S. & Crowe, T. P. Context- and density-dependent effects of introduced oysters on biodiversity. *Biol. Invasions* **16**, 1145–1163 (2014).
60. Thomsen, M. et al. Forty years of experiments on aquatic invasive species: are study biases limiting our understanding of impacts? *NeoBiota* **22**, 1–22 (2014).
61. Ricciardi, A. Are modern biological invasions an unprecedented form of global change? *Conserv. Biol.* **21**, 329–336 (2007).
62. Seebens, H. et al. Global rise in emerging alien species results from increased accessibility of new source pools. *Proc. Natl Acad. Sci. USA* **115**, E2264–E2273 (2018).
63. Davis, M. A. & Chew, M. K. 'The denialists are coming!' Well, not exactly: a response to Russell and Blackburn. *Trends Ecol. Evol.* **32**, 229–230 (2017).
64. Koricheva, J., Gurevitch, J., Mengersen, K. (eds) *Handbook of Meta-analysis in Ecology and Evolution* (Princeton Univ. Press, 2013).
65. Nosek, B. A. et al. Promoting an open research culture. *Science* **348**, 1422–1425 (2015).
66. Moher, D., Liberati, A., Tetzlaff, J., Altman, D. G. & PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med.* **6**, e1000097 (2009).
67. Borenstein, M., Hedges, L. V., Higgins, J. P. T. & Rothstein, H. R. *Introduction to Meta-Analysis* (Wiley-Blackwell, 2009).
68. Hannam, M. P. & Wyllie-Echeverria, S. Microtopography promotes coexistence of an invasive seagrass and its native congener. *Biol. Invasions* **17**, 381–395 (2015).
69. Viechbauer, W. Conducting meta-analyses in R with the metafor package. *J. Stat. Softw.* **36**, 1–48 (2010).
70. Egger, M., Smith, G. D., Schneider, M. & Minder, C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* **315**, 629–634 (1997).
71. Habeck, C. W. & Schultz, A. K. Community-level impacts of white-tailed deer on understory plants in North American forests: a meta-analysis. *Aob Plants* **7**, plv119 (2015).
72. Viechbauer, W. & Cheung, M. W.-L. Outlier and influence diagnostics for meta-analysis. *Res. Synth. Methods* **1**, 112–125 (2010).
73. Zarnetske, P. L., Seabloom, E. W. & Hacker, S. D. Non-target effects of invasive species management: beachgrass, birds, and bulldozers in coastal dunes. *Ecosphere* **1**, art13 (2010).
74. Moseman, S. M., Zhang, R., Qian, P. Y. & Levin, L. A. Diversity and functional responses of nitrogen-fixing microbes to three wetland invasions. *Biol. Invasions* **11**, 225–239 (2009).
75. Castorani, M. C. N. & Hovel, K. A. Invasive prey indirectly increase predation on their native competitors. *Ecology* **96**, 1911–1922 (2015).
76. Crooks, J. A. Assessing invader roles within changing ecosystems: historical and experimental perspectives on an exotic mussel in an urbanized lagoon. *Biol. Invasions* **3**, 23–36 (2001).

77. Munari, C. Effects of the exotic invader *Musculista senhousia* on benthic communities of two Mediterranean lagoons. *Hydrobiologia* **611**, 29–43 (2008).
78. Magalhaes, W. F. & Bailey-Brock, J. H. Polychaete assemblages associated with the invasive green alga *Avrainvillea amadelpha* and surrounding bare sediment patches in Hawaii. *Memoirs Mus. Vic.* **71**, 161–168 (2014).
79. Caño, L., Campos, J. A., García-Magro, D. & Herrera, M. Invasiveness and impact of the non-native shrub *Baccharis halimifolia* in sea rush marshes: fine-scale stress heterogeneity matters. *Biol. Invasions* **16**, 2063–2077 (2014).
80. Mendez, M. M., Schwindt, E. & Bortolus, A. Differential benthic community response to increased habitat complexity mediated by an invasive barnacle. *Aquat. Ecol.* **49**, 441–452 (2015).
81. Wonham, M. J., O'Connor, M. & Harley, C. D. G. Positive effects of a dominant invader on introduced and native mudflat species. *Mar. Ecol. Prog. Ser.* **289**, 109–116 (2005).
82. Elías, R. et al. Effect of the invader *Boccardia proboscidea* (Polychaeta: Spionidae) on richness, diversity and structure of SW Atlantic epilithic intertidal community. *Mar. Pollut. Bull.* **91**, 530–536 (2015).
83. Cordell, J. R., Levy, C. & Toft, J. D. Ecological implications of invasive tunicates associated with artificial structures in Puget Sound, Washington, USA. *Biol. Invasions* **15**, 1303–1318 (2013).
84. Bonnici, L., Evans, J., Borg, J. A. & Schembri, P. J. J. Biological aspects and ecological effects of a bed of the invasive non-indigenous mussel *Brachidontes pharaonis* (Fischer P., 1870) in Malta. *Mediterr. Mar. Sci.* **13**, 153–161 (2012).
85. Malyshev, A. & Quijón, P. A. Disruption of essential habitat by a coastal invader: new evidence of the effects of green crabs on eelgrass beds. *ICES J. Mar. Sci.* **68**, 1852–1856 (2011).
86. Gregory, G. J. & Quijón, P. A. The impact of a coastal invasive predator on infaunal communities: assessing the roles of density and a native counterpart. *J. Sea Res.* **66**, 181–186 (2011).
87. Whitlow, W. L. Changes in survivorship, behavior, and morphology in native soft-shell clams induced by invasive green crab predators. *Mar. Ecol.* **31**, 418–430 (2010).
88. Whitlow, W. L., Rice, N. A. & Sweeney, C. Native species vulnerability to introduced predators: testing an inducible defense and a refuge from predation. *Biol. Invasions* **5**, 23–31 (2003).
89. Large, S. I. & Smee, D. L. Biogeographic variation in behavioral and morphological responses to predation risk. *Oecologia* **171**, 961–969 (2013).
90. Griffen, B. D. & Byers, J. E. Community impacts of two invasive crabs: the interactive roles of density, prey recruitment, and indirect effects. *Biol. Invasions* **11**, 927–940 (2009).
91. Estelle, V. & Grosholz, E. D. Experimental test of the effects of a non-native invasive species on a wintering shorebird. *Conserv. Biol.* **26**, 472–481 (2012).
92. Wootton, L. First report of *Carex macrocephala* in Eastern North America with notes on its co-occurrence with *Carex kobomugi* in New Jersey. *J. Torrey Bot. Soc.* **134**, 126–134 (2007).
93. Wootton, L. S. et al. When invasive species have benefits as well as costs: managing *Carex kobomugi* (Asiatic sand sedge) in New Jersey's coastal dunes. *Biol. Invasions* **7**, 1017–1027 (2005).
94. Novoa, A. & González, L. Impacts of *Carpobrotus edulis* (L.) N.E.Br. on the germination, establishment and survival of native plants: a clue for assessing its competitive strength. *PLoS ONE* **9**, e107557 (2014).
95. Magnoli, S. M., Kleinhesselink, A. R. & Cushman, J. H. Responses to invasion and invader removal differ between native and exotic plant groups in a coastal dune. *Oecologia* **173**, 1521–1530 (2013).
96. Novoa, A., González, L., Moravcová, L. & Pyšek, P. Constraints to native plant species establishment in coastal dune communities invaded by *Carpobrotus edulis*: implications for restoration. *Biol. Conserv.* **164**, 1–9 (2013).
97. Hata, K., Kato, H. & Kachi, N. Seedlings of a native shrub can establish under forests dominated by an alien tree, *Casuarina equisetifolia*, on subtropical oceanic islands. *J. For. Res.* **17**, 208–212 (2012).
98. Smith, J. R., Vogt, S. C., Creedon, F., Lucas, B. J. & Eernisse, D. J. The non-native turf-forming alga *Caulacanthus ustulatus* displaces space-occupants but increases diversity. *Biol. Invasions* **16**, 2195–2208 (2014).
99. Bulleri, F. & Piazza, L. Variations in importance and intensity of competition underpin context dependency in the effects of an invasive seaweed on resident assemblages. *Mar. Biol.* **162**, 485–489 (2015).
100. Tomas, F., Box, A. & Terrados, J. Effects of invasive seaweeds on feeding preference and performance of a keystone Mediterranean herbivore. *Biol. Invasions* **13**, 1559–1570 (2011).
101. Tamburello, L. et al. Variation in the impact of non-native seaweeds along gradients of habitat degradation: a meta-analysis and an experimental test. *Oikos* **124**, 1121–1131 (2015).
102. Klein, J. C. & Verlaque, M. Macrophyte assemblage associated with an invasive species exhibiting temporal variability in its development pattern. *Hydrobiologia* **636**, 369–378 (2009).
103. Kersting, D. K., Ballesteros, E., De Caralt, S. & Linares, C. Invasive macrophytes in a marine reserve (Columbretes Islands, NW Mediterranean): spread dynamics and interactions with the endemic scleractinian coral *Cladocora caespitosa*. *Biol. Invasions* **16**, 1599–1610 (2014).
104. Tejada, S., Deudero, S., Box, A. & Sureda, A. Physiological response of the sea urchin *Paracentrotus lividus* fed with the seagrass *Posidonia oceanica* and the alien algae *Caulerpa racemosa* and *Lophocladia lallemandii*. *Mar. Environ. Res.* **83**, 48–53 (2013).
105. Klein, J. C. & Velarque, M. Temporal trends in invasion impacts in macrophyte assemblages of the Mediterranean Sea. *Cah. Biol. Mar.* **53**, 403–407 (2012).
106. Vázquez-Luis, M., Sanchez-Jerez, P. & Bayle-Sempere, J. T. Effects of *Caulerpa racemosa* var. *cylindracea* on prey availability: an experimental approach to predation of amphipods by *Thalassoma pavo* (Labridae). *Hydrobiologia* **654**, 147–154 (2010).
107. Vázquez-Luis, M., Sanchez-Jerez, P. & Bayle-Sempere, J. T. Changes in amphipod (Crustacea) assemblages associated with shallow-water algal habitats invaded by *Caulerpa racemosa* var. *cylindracea* in the western Mediterranean Sea. *Mar. Environ. Res.* **65**, 416–426 (2008).
108. Cebrian, E., Linares, C., Marschal, C. & Garrabou, J. Exploring the effects of invasive algae on the persistence of gorgonian populations. *Biol. Invasions* **14**, 2647–2656 (2012).
109. Gribben, P. E., Byers, J. E., Wright, J. T. & Glasby, T. M. Positive versus negative effects of an invasive ecosystem engineer on different components of a marine ecosystem. *Oikos* **122**, 816–824 (2013).
110. Gallucci, F., Hutchings, P., Gribben, P. & Fonseca, G. Habitat alteration and community-level effects of an invasive ecosystem engineer: a case study along the coast of NSW, Australia. *Mar. Ecol. Prog. Ser.* **449**, 95–108 (2012).
111. Gribben, P. E. et al. Reduced performance of native infauna following recruitment to a habitat-forming invasive marine alga. *Oecologia* **158**, 733–745 (2009).
112. Wright, J. T. & Gribben, P. E. Predicting the impact of an invasive seaweed on the fitness of native fauna. *J. Appl. Ecol.* **45**, 1540–1549 (2008).
113. Byers, J. E., Wright, J. T. & Gribben, P. E. Variable direct and indirect effects of a habitat-modifying invasive species on mortality of native fauna. *Ecology* **91**, 1787–1798 (2010).
114. Taylor, S. L., Bishop, M. J., Kelaher, B. P. & Glasby, T. M. Impacts of detritus from the invasive alga *Caulerpa taxifolia* on a soft sediment community. *Mar. Ecol. Prog. Ser.* **420**, 73–81 (2010).
115. Gribben, P. E., Wright, J. T., O'Connor, W. A. & Steinberg, P. Larval settlement preference of a native bivalve: the influence of an invasive alga versus native substrata. *Aquat. Biol.* **7**, 217–227 (2009).
116. Strain, E. M. A. & Johnson, C. R. The effects of an invasive habitat modifier on the biotic interactions between two native herbivorous species and benthic habitat in a subtidal rocky reef ecosystem. *Biol. Invasions* **15**, 1391–1405 (2013).
117. Strain, E. M. A. & Johnson, C. R. Competition between an invasive urchin and commercially fished abalone. *Mar. Ecol. Prog. Ser.* **377**, 169–182 (2009).
118. Mason, T. J. & French, K. Management regimes for a plant invader differentially impact resident communities. *Biol. Conserv.* **136**, 246–259 (2007).
119. Bugnot, A. B., Coleman, R. A., Figueira, W. F. & Marzinielli, E. M. Community-level impacts of the invasive isopod *Cirolana harfordi*. *Biol. Invasions* **17**, 1149–1161 (2015).
120. Drouin, A., McKindsey, C. W. & Johnson, L. E. Higher abundance and diversity in faunal assemblages with the invasion of *Codium fragile* ssp. *fragile* in eelgrass meadows. *Mar. Ecol. Prog. Ser.* **424**, 105–117 (2011).
121. Schmidt, A. L. & Scheibling, R. E. Effects of native and invasive macroalgal canopies on composition and abundance of mobile benthic macrofauna and turf-forming algae. *J. Exp. Mar. Biol. Ecol.* **341**, 110–130 (2007).
122. Chavanich, S. & Harris, L. G. Impact of the non-native macroalga *Codium fragile* (Sur.) Hariot ssp. *tomentosoides* (van Goor) Silva on the native snail *Lacuna vincta* (Montagu, 1803) in the Gulf of Maine. *Veliger* **47**, 85–90 (2004).
123. Drouin, A., McKindsey, C. W. & Johnson, L. E. Detecting the impacts of notorious invaders: experiments versus observations in the invasion of eelgrass meadows by the green seaweed *Codium fragile*. *Oecologia* **168**, 491–502 (2012).
124. Novais, A., Souza, A. T., Ilarri, M., Pascoal, C. & Sousa, R. Facilitation in the low intertidal: effects of an invasive species on the structure of an estuarine macrozoobenthic assemblage. *Mar. Ecol. Prog. Ser.* **522**, 157–167 (2015).
125. Ilarri, M. I., Souza, A. T., Antunes, C., Guilhermino, L. & Sousa, R. Influence of the invasive Asian clam *Corbicula fluminea* (Bivalvia: Corbiculidae) on estuarine epibenthic assemblages. *Estuar. Coast. Shelf Sci.* **143**, 12–19 (2014).
126. Talman, S. & Keough, M. Impact of an exotic clam, *Corbula gibba*, on the commercial scallop *Pecten fumatus* in Port Phillip Bay, south-east Australia: evidence of resource-restricted growth in a subtropical environment. *Mar. Ecol. Prog. Ser.* **221**, 135–143 (2001).

127. Ragueneau, O. et al. Biodeposition by an invasive suspension feeder impacts the biogeochemical cycle of Si in a coastal ecosystem (Bay of Brest, France). *Biogeochemistry* **75**, 19–41 (2005).
128. Le Pape, O., Guerault, D. & Desanay, Y. Effect of an invasive mollusc, American slipper limpet *Crepidula fornicata*, on habitat suitability for juvenile common sole *Solea solea* in the Bay of Biscay. *Mar. Ecol. Prog. Ser.* **277**, 107–115 (2004).
129. Martin, S. et al. Benthic community respiration in areas impacted by the invasive mollusk *Crepidula fornicata*. *Mar. Ecol. Prog. Ser.* **347**, 51–60 (2007).
130. Smith, B. E., Collie, J. S. & Lengyel, N. L. Fish trophic engineering: ecological effects of the invasive ascidian *Didemnum vexillum* (Georges Bank, northwestern Atlantic). *J. Exp. Mar. Biol. Ecol.* **461**, 489–498 (2014).
131. Long, H. A. & Grosholz, E. D. Overgrowth of eelgrass by the invasive colonial tunicate *Didemnum vexillum*: consequences for tunicate and eelgrass growth and epifauna abundance. *J. Exp. Mar. Biol. Ecol.* **473**, 188–194 (2015).
132. Dijkstra, J. A., Lambert, W. J. & Harris, L. G. Introduced species provide a novel temporal resource that facilitates native predator population growth. *Biol. Invasions* **15**, 911–919 (2013).
133. Petillon, J., Frederic, Y., Alain, C. & Lefeuvre, J.-C. Impact of an invasive plant (*Elymus athericus*) on the conservation value of tidal salt marshes in western France and implications for management: responses of spider populations. *Biol. Conserv.* **126**, 103–117 (2005).
134. Bruschetti, M., Bazterrica, C., Luppi, T. & Iribarne, O. An invasive intertidal reef-forming polychaete affect habitat use and feeding behavior of migratory and locals birds in a SW Atlantic coastal lagoon. *J. Exp. Mar. Biol. Ecol.* **375**, 76–83 (2009).
135. Bazterrica, M. C., Botto, F. & Iribarne, O. Effects of an invasive reef-building polychaete on the biomass and composition of estuarine macroalgal assemblages. *Biol. Invasions* **14**, 765–777 (2012).
136. Bruschetti, M., Luppi, T., Fanjul, E., Rosenthal, A. & Iribarne, O. Grazing effect of the invasive reef-forming polychaete *Ficopomatus enigmaticus* (Fauvel) on phytoplankton biomass in a SW Atlantic coastal lagoon. *J. Exp. Mar. Biol. Ecol.* **354**, 212–219 (2008).
137. Schwindt, E., Iribarne, O. O. & Isla, F. I. Physical effects of an invading reef-building polychaete on an Argentinean estuarine environment. *Estuar. Coast. Shelf Sci.* **59**, 109–120 (2004).
138. Bazterrica, M. C. et al. Factors controlling macroalgae assemblages in a Southwest Atlantic coastal lagoon modified by an invading reef forming polychaete. *J. Exp. Mar. Biol. Ecol.* **443**, 169–177 (2013).
139. Heiman, K. W. & Micheli, F. Non-native ecosystem engineer alters estuarine communities. *Integr. Comp. Biol.* **50**, 226–236 (2010).
140. Bruschetti, M., Bazterrica, C., Fanjul, E., Luppi, T. & Iribarne, O. Effect of biodeposition of an invasive polychaete on organic matter content and productivity of the sediment in a coastal lagoon. *J. Sea Res.* **66**, 20–28 (2011).
141. Fukunaga, A., Peyton, K. A. & Thomas, F. I. M. Epifaunal community structure and ammonium uptake compared for the invasive algae, *Gracilaria salicornia* and *Acanthophora speciosa*, and the native alga, *Padina thuyi*. *J. Exp. Mar. Biol. Ecol.* **456**, 78–86 (2014).
142. Thomsen, M., Stæhr, P. A., Nejrup, L. B. & Schiel, D. R. Effects of the invasive macroalgae *Gracilaria vermiculophylla* on two co-occurring foundation species and associated invertebrates. *Aquat. Invasions* **8**, 133–145 (2013).
143. Höfle, H., Thomsen, M. S. & Holmer, M. High mortality of *Zostera marina* under high temperature regimes but minor effects of the invasive macroalgae *Gracilaria vermiculophylla*. *Estuar. Coast. Shelf Sci.* **92**, 35–46 (2011).
144. Thomsen, M. S. Experimental evidence for positive effects of invasive seaweed on native invertebrates via habitat-formation in a seagrass bed. *Aquat. Invasions* **5**, 341–346 (2010).
145. Martínez-Lüscher, J. & Holmer, M. Potential effects of the invasive species *Gracilaria vermiculophylla* on *Zostera marina* metabolism and survival. *Mar. Environ. Res.* **69**, 345–349 (2010).
146. Janiak, D. S. & Whitlatch, R. B. Epifaunal and algal assemblages associated with the native *Chondrus crispus* (Stackhouse) and the non-native *Grateloupia turuturu* (Yamada) in eastern Long Island Sound. *J. Exp. Mar. Biol. Ecol.* **413**, 38–44 (2012).
147. Brousseau, D. & Goldberg, R. Effect of predation by the invasive crab *Hemigrapsus sanguineus* on recruiting barnacles *Semibalanus balanoides* in western Long Island Sound, USA. *Mar. Ecol. Prog. Ser.* **339**, 221–228 (2007).
148. Brousseau, D. J., Goldberg, R. & Garza, C. Impact of predation by the invasive crab *Hemigrapsus sanguineus* on survival of juvenile blue mussels in Western Long Island Sound. *Northeast. Nat.* **21**, 119–133 (2014).
149. Bennett, A. E., Thomsen, M. & Strauss, S. Y. Multiple mechanisms enable invasive species to suppress native species. *Am. J. Bot.* **98**, 1086–1094 (2011).
150. Harvey, K. J., Britton, D. R. & Minchinton, T. E. Detecting impacts of non-native species on associated invertebrate assemblages depends on microhabitat. *Austral. Ecol.* **39**, 511–521 (2014).
151. Kamalakannan, B., Jeevamani, J. J. J., Nagendran, N. A., Pandiaraja, D. & Chandrasekaran, S. Impact of removal of invasive species *Kappaphycus alvarezii* from coral reef ecosystem in Gulf of Mannar, India. *Curr. Sci.* **106**, 1401–1408 (2014).
152. Sato, S., Chiba, T. & Hasegawa, H. Long-term fluctuations in mollusk populations before and after the appearance of the alien predator *Euspira fortunei* on the Tona coast, Miyagi Prefecture, northern Japan. *Fish. Sci.* **78**, 589–595 (2012).
153. Rowles, A. D. & O'dowd, D. J. Impacts of the invasive Argentine ant on native ants and other invertebrates in coastal scrub in south-eastern Australia. *Austral. Ecol.* **34**, 239–248 (2009).
154. Bedini, R., Bedini, M., Bonechi, L. & Piazzini, L. Effects of non-native turf-forming Rhodophyta on mobile macro-invertebrate assemblages in the north-western Mediterranean Sea. *Mar. Biol. Res.* **11**, 430–437 (2015).
155. Eash-Loucks, W. E., Kimball, M. E. & Petrinc, K. M. Long-term changes in an estuarine mud crab community: evaluating the impact of non-native species. *J. Crustac. Biol.* **34**, 731–738 (2014).
156. O'Shaughnessy, K. A. & Harding, J. M. & Burge, E. J. Ecological effects of the invasive parasite *Loxothylacus panopaei* on the flatback mud crab *Eurypanopeus depressus* with implications for estuarine communities. *Bull. Mar. Sci.* **90**, 611–621 (2014).
157. Wilkie, E. M., Bishop, M. J. & O'Connor, W. A. The density and spatial arrangement of the invasive oyster *Crassostrea gigas* determines its impact on settlement of native oyster larvae. *Ecol. Evol.* **3**, 4851–4860 (2013).
158. Krassoi, F. R., Brown, K. R., Bishop, M. J., Kelaher, B. P. & Summerhayes, S. Condition-specific competition allows coexistence of competitively superior exotic oysters with native oysters. *J. Anim. Ecol.* **77**, 5–15 (2008).
159. Bray, D. J., Green, I., Golicher, D. & Herbert, R. J. H. Spatial variation of trace metals within intertidal beds of native mussels (*Mytilus edulis*) and non-native Pacific oysters (*Crassostrea gigas*): implications for the food web? *Hydrobiologia* **757**, 235–249 (2015).
160. Green, D. S., Boots, B. & Crowe, T. P. Effects of non-indigenous oysters on microbial diversity and ecosystem functioning. *PLoS ONE* **7**, e48410 (2012).
161. Green, D. S. & Crowe, T. P. Physical and biological effects of introduced oysters on biodiversity in an intertidal boulder field. *Mar. Ecol. Prog. Ser.* **482**, 119–132 (2013).
162. Wegener, A. R. & Buschbaum, C. Alien epibiont (*Crassostrea gigas*) impacts on native periwinkles (*Littorina littorea*). *Aquat. Invasions* **6**, 281–290 (2011).
163. Kochmann, J., Buschbaum, C., Volkenborn, N. & Reise, K. Shift from native mussels to alien oysters: differential effects of ecosystem engineers. *J. Exp. Mar. Biol. Ecol.* **364**, 1–10 (2008).
164. Green, D. S., Rocha, C. & Crowe, T. P. Effects of non-indigenous oysters on ecosystem processes vary with abundance and context. *Ecosystems* **16**, 881–893 (2013).
165. Nicastro, M. A., Bishop, M. J., Kelaher, B. P. & Benedetti-Cecchi, L. Export of non-native gastropod shells to a coastal lagoon: alteration of habitat structure has negligible effects on infauna. *J. Exp. Mar. Biol. Ecol.* **374**, 31–36 (2009).
166. Hietanen, S., Laine, A. O. & Lukkari, K. The complex effects of the invasive polychaetes *Marenzelleria* spp. on benthic nutrient dynamics. *J. Exp. Mar. Biol. Ecol.* **352**, 89–102 (2007).
167. Urban-Malinga, B., Warzocha, J. & Zalewski, M. Effects of the invasive polychaete *Marenzelleria* spp. on benthic processes and meiobenthos of a species-poor brackish system. *J. Sea Res.* **80**, 25–34 (2013).
168. Quintana, C. O., Kristensen, E. & Valdemarsen, T. Impact of the invasive polychaete *Marenzelleria viridis* on the biogeochemistry of sandy marine sediments. *Biogeochemistry* **115**, 95–109 (2013).
169. Kristensen, E., Hansen, T., Delefosse, M., Banta, G. T. & Quintana, C. O. Contrasting effects of the polychaetes *Marenzelleria viridis* and *Nereis diversicolor* on benthic metabolism and solute transport in sandy coastal sediment. *Mar. Ecol. Prog. Ser.* **425**, 125–139 (2011).
170. Yuan, W. S., Hoffman, E. A. & Walters, L. J. Effects of nonnative invertebrates on two life stages of the native eastern oyster *Crassostrea virginica*. *Biol. Invasions* **18**, 689–701 (2016).
171. Krumhansl, K. A., Lee, J. M. & Scheibling, R. E. Grazing damage and encrustation by an invasive bryozoan reduce the ability of kelps to withstand breakage by waves. *J. Exp. Mar. Biol. Ecol.* **407**, 12–18 (2011).
172. Roohi, A. et al. Impact of a new invasive ctenophore (*Mnemiopsis leidyi*) on the zooplankton community of the Southern Caspian sea. *Mar. Ecol.* **29**, 421–434 (2008).
173. Aldridge, D. C., Salazar, M., Serna, A. & Cock, J. Density-dependent effects of a new invasive false mussel, *Mytilopsis trautevineana* (Tryon 1866), on shrimp, *Litopenaeus vannamei* (Boone 1931), aquaculture in Colombia. *Aquaculture* **281**, 34–42 (2008).
174. Branch, G. M., Odendaal, F. & Robinson, T. B. Competition and facilitation between the alien mussel *Mytilus galloprovincialis* and indigenous species: moderation by wave action. *J. Exp. Mar. Biol. Ecol.* **383**, 65–78 (2010).
175. Sellheim, K., Stachowicz, J. J. & Coates, R. C. Effects of a nonnative habitat-forming species on mobile and sessile epifaunal communities. *Mar. Ecol. Prog. Ser.* **398**, 69–80 (2010).

176. Hanekom, N. Invasion of an indigenous *Perna perna* mussel bed on the south coast of South Africa by an alien mussel *Mytilus galloprovincialis* and its effect on the associated fauna. *Biol. Invasions* **10**, 233–244 (2008).
177. Chapman, M. G., People, J. & Blockley, D. Intertidal assemblages associated with natural coralina turf and invasive mussel beds. *Biodivers. Conserv.* **14**, 1761–1776 (2005).
178. Hollebone, A. L. & Hay, M. E. An invasive crab alters interaction webs in a marine community. *Biol. Invasions* **10**, 347–358 (2008).
179. Kimball, M. E. & Able, K. W. Nekton utilization of intertidal salt marsh creeks: tidal influences in natural *Spartina*, invasive *Phragmites*, and marshes treated for *Phragmites* removal. *J. Exp. Mar. Biol. Ecol.* **346**, 87–101 (2007).
180. Torchin, M. E. Native fish grows faster in the presence of a potential introduced competitor. *Aquat. Invasions* **5**, 163–167 (2010).
181. Richman, S. E. & Lovvorn, J. R. Relative foraging value to lesser scaup ducks of native and exotic clams from San Francisco Bay. *Ecol. Appl.* **14**, 1217–1231 (2004).
182. Brenneis, V. E. F., Sih, A. & de Rivera, C. E. Integration of an invasive consumer into an estuarine food web: direct and indirect effects of the New Zealand mud snail. *Oecologia* **167**, 169–179 (2011).
183. Elise, S., Urbina-Barreto, I., Boadas-Gil, H., Galindo-Vivas, M. & Kulbicki, M. No detectable effect of lionfish (*Pterois volitans* and *P. miles*) invasion on a healthy reef fish assemblage in Archipelago Los Roques National Park, Venezuela. *Mar. Biol.* **162**, 319–330 (2015).
184. Black, A. N., Weimann, S. R., Imhoff, V. E., Richter, M. L. & Itzkowitz, M. A differential prey response to invasive lionfish, *Pterois volitans*: prey naivete and risk-sensitive courtship. *J. Exp. Mar. Biol. Ecol.* **460**, 1–7 (2014).
185. Albins, M. A. Effects of invasive Pacific red lionfish *Pterois volitans* versus a native predator on Bahamian coral-reef fish communities. *Biol. Invasions* **15**, 29–43 (2013).
186. Anton, A. et al. Prey naivete to invasive lionfish (*Pterois volitans*) on Caribbean coral reefs. *Mar. Ecol. Prog. Ser.* **544**, 257–269 (2016).
187. Green, S. J. et al. Linking removal targets to the ecological effects of invaders: a predictive model and field test. *Ecol. Appl.* **24**, 1311–1322 (2014).
188. Ingeman, K. E. & Webster, M. S. Native prey mortality increases but remains density-dependent following lionfish invasion. *Mar. Ecol. Prog. Ser.* **531**, 241–252 (2015).
189. Sweetman, A. K. et al. Impacts of exotic mangrove forests and mangrove deforestation on carbon remineralization and ecosystem functioning in marine sediments. *Biogeosciences* **7**, 2129–2145 (2010).
190. Bidegain, G. & Juanes, J. A. Does expansion of the introduced Manila clam *Ruditapes philippinarum* cause competitive displacement of the European native clam *Ruditapes decussatus*? *J. Exp. Mar. Biol. Ecol.* **445**, 44–52 (2013).
191. Holloway, M. G. & Keough, M. J. An introduced polychaete affects recruitment and larval abundance of sessile invertebrates. *Ecol. Appl.* **12**, 1803–1823 (2002).
192. Holloway, M. G. & Keough, M. J. Effects of an introduced polychaete, *Sabella spallanzanii*, on the development of epifaunal assemblages. *Mar. Ecol. Prog. Ser.* **236**, 137–154 (2002).
193. Robinson, T. B. & Swart, C. Distribution and impact of the alien anemone *Sagartia ornata* in the West Coast National Park. *Koedoe* **57**, a1246 (2015).
194. Salvaterra, T., Green, D. S., Crowe, T. P. & O’Gorman, E. J. Impacts of the invasive alga *Sargassum muticum* on ecosystem functioning and food web structure. *Biol. Invasions* **15**, 2563–2576 (2013).
195. Lang, A. C. & Buschbaum, C. Facilitative effects of introduced Pacific oysters on native macroalgae are limited by a secondary invader, the seaweed *Sargassum muticum*. *J. Sea Res.* **63**, 119–128 (2010).
196. Sánchez, I. & Fernández, C. Impact of the invasive seaweed *Sargassum muticum* (phaeophyta) on an intertidal macroalgal assemblage 1. *J. Phycol.* **41**, 923–930 (2005).
197. Rossi, F., Incera, M., Callier, M. & Olabarria, C. Effects of detrital non-native and native macroalgae on the nitrogen and carbon cycling in intertidal sediments. *Mar. Biol.* **158**, 2705–2715 (2011).
198. Britton-Simmons, K. H., Pister, B., Sánchez, I. & Okamoto, D. Response of a native, herbivorous snail to the introduced seaweed *Sargassum muticum*. *Hydrobiologia* **661**, 187–196 (2011).
199. Vye, S. R., Emmerson, M. C., Arenas, F., Dick, J. T. A. & O’Connor, N. E. Stressor intensity determines antagonistic interactions between species invasion and multiple stressor effects on ecosystem functioning. *Oikos* **124**, 1005–1012 (2015).
200. Rodil, I. F., Olabarria, C., Lastra, M. & López, J. Differential effects of native and invasive algal wrack on macrofaunal assemblages inhabiting exposed sandy beaches. *J. Exp. Mar. Biol. Ecol.* **358**, 1–13 (2008).
201. Vaz-Pinto, F., Olabarria, C. & Arenas, F. Ecosystem functioning impacts of the invasive seaweed *Sargassum muticum* (Fucales, Phaeophyceae). *J. Phycol.* **50**, 108–116 (2014).
202. Buschbaum, C., Chapman, A. S. & Saier, B. How an introduced seaweed can affect epibiota diversity in different coastal systems. *Mar. Biol.* **148**, 743–754 (2006).
203. White, L. F. & Shurin, J. B. Density dependent effects of an exotic marine macroalga on native community diversity. *J. Exp. Mar. Biol. Ecol.* **405**, 111–119 (2011).
204. Viejo, R. M. Mobile epifauna inhabiting the invasive *Sargassum muticum* and two local seaweeds in northern Spain. *Aquat. Bot.* **64**, 131–149 (1999).
205. Tang, Y. et al. Ecological influence of exotic plants of *Sonneratia apetala* on understory macrofauna. *Acta Oceanol. Sin.* **31**, 115–125 (2012).
206. Ma, Z., Gan, X., Cai, Y., Chen, J. & Li, B. Effects of exotic *Spartina alterniflora* Loisel on the habitat patch associations of breeding saltmarsh birds at Chongming Dongtan in the Yangtze River estuary, China. *Biol. Invasions* **13**, 1673–1686 (2011).
207. Gan, X. et al. Potential impacts of invasive *Spartina alterniflora* on spring bird communities at Chongming Dongtan, a Chinese wetland of international importance. *Estuar. Coast. Shelf Sci.* **83**, 211–218 (2009).
208. Zhou, H.-X., Liu, J.-E., Zhou, J. & Qin, P. Effect of an alien species *Spartina alterniflora* Loisel on biogeochemical processes of intertidal ecosystem in the Jiangsu coastal region, China. *Pedosphere* **18**, 77–85 (2008).
209. Neira, C., Levin, L. A., Grosholz, E. D. & Mendoza, G. Influence of invasive *Spartina* growth stages on associated macrofaunal communities. *Biol. Invasions* **9**, 975–993 (2007).
210. Chen, H., Zhang, P., Li, B. & Wu, J. Invasive cordgrass facilitates epifaunal communities in a Chinese marsh. *Biol. Invasions* **17**, 205–217 (2015).
211. Ma, Z., Gan, X., Choi, C.-Y. & Li, B. Effects of invasive cordgrass on presence of Marsh Grassbird in an area where it is not native. *Conserv. Biol. J. Soc. Conserv. Biol.* **28**, 150–158 (2014).
212. Tong, C. et al. Contrasting nutrient stocks and litter decomposition in stands of native and invasive species in a sub-tropical estuarine marsh. *Environ. Res.* **111**, 909–916 (2011).
213. Qin, H. et al. Effects of invasive cordgrass on crab distributions and diets in a Chinese salt marsh. *Mar. Ecol. Prog. Ser.* **415**, 177–187 (2010).
214. Wu, Y.-T. et al. Effects of saltmarsh invasion by *Spartina alterniflora* on arthropod community structure and diets. *Biol. Invasions* **11**, 635–649 (2009).
215. Yuan, J. et al. Shifts in methanogen community structure and function across a coastal marsh transect: effects of exotic *Spartina alterniflora* invasion. *Sci. Rep.* **6**, 18777 (2016).
216. Zhou, H.-X., Liu, J. & Qin, P. Impacts of an alien species (*Spartina alterniflora*) on the macrobenthos community of Jiangsu coastal inter-tidal ecosystem. *Ecol. Eng.* **35**, 521–528 (2009).
217. Cutajar, J., Shimeta, J. & Nuggeoda, D. Impacts of the invasive grass *Spartina anglica* on benthic macrofaunal assemblages in a temperate Australian saltmarsh. *Mar. Ecol. Prog. Ser.* **464**, 107–120 (2012).
218. Hacker, S. D. & Dethier, M. N. Community modification by a grass invader has differing impacts for marine habitats. *Oikos* **113**, 279–286 (2006).
219. Gooden, B. & French, K. Impacts of alien plant invasion on native plant communities are mediated by functional identity of resident species, not resource availability. *Oikos* **124**, 298–306 (2015).
220. Gooden, B. & French, K. Non-interactive effects of plant invasion and landscape modification on native communities. *Divers. Distrib.* **20**, 626–639 (2014).
221. Pettay, D. T., Wham, D. C., Smith, R. T., Iglesias-Prieto, R. & LaJeunesse, T. C. Microbial invasion of the Caribbean by an Indo-Pacific coral zooxanthella. *Proc. Natl Acad. Sci. USA* **112**, 7513–7518 (2015).
222. Miranda, R. J., Cruz, I. C. S. & Barros, F. Effects of the alien coral *Tubastraea tagusensis* on native coral assemblages in a southwestern Atlantic coral reef. *Mar. Biol.* **163**, 45 (2016).
223. Irigoyen, A. J., Trobbiani, G., Sgarlatta, M. P. & Raffo, M. P. Effects of the alien algae *Undaria pinnatifida* (Phaeophyceae, Laminariales) on the diversity and abundance of benthic macrofauna in Golfo Nuevo (Patagonia, Argentina): potential implications for local food webs. *Biol. Invasions* **13**, 1521–1532 (2011).
224. Forrest, B. M. & Taylor, M. D. Assessing invasion impact: survey design considerations and implications for management of an invasive marine plant. *Biol. Invasions* **4**, 375–386 (2002).
225. South, P. M. et al. Transient effects of an invasive kelp on the community structure and primary productivity of an intertidal assemblage. *Mar. Freshw. Res.* **67**, 103–112 (2016).
226. Gestoso, I., Arenas, F. & Olabarria, C. Biotic resistance and facilitation of a non-indigenous mussel vary with environmental context. *Mar. Ecol. Prog. Ser.* **506**, 163–173 (2014).
227. Hendrickx, J. P., Creese, R. G. & Gribben, P. E. Impacts of a non-native gastropod with a limited distribution; less conspicuous invaders matter too. *Mar. Ecol. Prog. Ser.* **537**, 151–162 (2015).

Acknowledgements

The authors are thankful to the following individuals: S. Ghani for conducting the network diagrams for Fig. 2, using resources and services at the Visualization Core Lab at KAUST; I. Ferri for advice on the design of Fig. 5; and C. Nelson for her assistance organizing the Invasive Species Workshop at KAUST. This research was supported by King Abdullah University of Science and Technology (KAUST) through baseline funding

to C.M.D., by Horizon 2020 Framework Programme (EU Framework Programme for Research and Innovation H2020) (grant no. 659246) to S.B., by the Ministry of Economy and Competitiveness, Agencia Estatal de Investigación (Spanish Agencia Estatal de Investigación) (grant no. FJCI – 2016 – 30728) to S.B., by the Ministry of Economy and Competitiveness, Agencia Estatal de Investigación (Spanish Agencia Estatal de Investigación) (grant no. CGL 2015 – 71809 – P) to N.M., J.S.-G. and S.B., and by the ARC Centre of Excellence for Coral Reef Studies (grant no. CE 140100020) to J.M.P.

Author contributions

C.M.D., A.A., C.E.L. and N.R.G. conceived and designed the study. A.A., N.R.G., C.E.L., E.T.A., S.B., J.C., D.K.-J., N.M., P.M., J.M.P. and J.S.-G. constructed the dataset. A.A. and N.R.G. performed the data analyses with contributions from all co-authors. All authors contributed to writing and improving the manuscript and approved the submission.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information is available for this paper at <https://doi.org/10.1038/s41559-019-0851-0>.

Reprints and permissions information is available at www.nature.com/reprints.

Correspondence and requests for materials should be addressed to A.A.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

© The Author(s), under exclusive licence to Springer Nature Limited 2019

Reporting Summary

Nature Research wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Research policies, see [Authors & Referees](#) and the [Editorial Policy Checklist](#).

Statistics

For all statistical analyses, confirm that the following items are present in the figure legend, table legend, main text, or Methods section.

n/a Confirmed

- ☐ ☒ The exact sample size (n) for each experimental group/condition, given as a discrete number and unit of measurement
- ☐ ☒ A statement on whether measurements were taken from distinct samples or whether the same sample was measured repeatedly
- ☐ ☒ The statistical test(s) used AND whether they are one- or two-sided
Only common tests should be described solely by name; describe more complex techniques in the Methods section.
- ☐ ☒ A description of all covariates tested
- ☐ ☒ A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons
- ☐ ☒ A full description of the statistical parameters including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient) AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals)
- ☐ ☒ For null hypothesis testing, the test statistic (e.g. F , t , r) with confidence intervals, effect sizes, degrees of freedom and P value noted
Give P values as exact values whenever suitable.
- ☒ ☐ For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings
- ☐ ☒ For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes
- ☐ ☒ Estimates of effect sizes (e.g. Cohen's d , Pearson's r), indicating how they were calculated

Our web collection on [statistics for biologists](#) contains articles on many of the points above.

Software and code

Policy information about [availability of computer code](#)

Data collection	Data from figures and tables were extracted using Graph Click (available at http://www.arizona-software.ch/graphclick/download.html) following a set of study criteria and a protocol for data extraction.
Data analysis	All statistical analysis were conducting in RStudio version 1.1.423 using the Metafor package to obtain the mean effect size Hedges' g and the 95% confidence intervals from mixed-effects models using the <code>rma.mv</code> function. The calculations to obtain effect size Hedges' g and the variance for each effect size are specified in the methods. The R script used in this manuscript will be deposited in the Github community repository (https://github.com/ngerald/marine-exotics-global-analysis).

For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors/reviewers. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Research [guidelines for submitting code & software](#) for further information.

Data

Policy information about [availability of data](#)

All manuscripts must include a [data availability statement](#). This statement should provide the following information, where applicable:

- Accession codes, unique identifiers, or web links for publicly available datasets
- A list of figures that have associated raw data
- A description of any restrictions on data availability

All data underlying the study is deposited in PANGAEA at <https://doi.pangaea.de/10.1594/PANGAEA.895681> and will be public after publication. In addition, Supplementary material Table 2 lists the 159 references reviewed on this study.

Field-specific reporting

Please select the one below that is the best fit for your research. If you are not sure, read the appropriate sections before making your selection.

☐ Life sciences ☐ Behavioural & social sciences ☒ Ecological, evolutionary & environmental sciences

For a reference copy of the document with all sections, see [nature.com/documents/nr-reporting-summary-flat.pdf](https://www.nature.com/documents/nr-reporting-summary-flat.pdf)

Ecological, evolutionary & environmental sciences study design

All studies must disclose on these points even when the disclosure is negative.

Study description	In this study we assessed the global ecological effects of exotic species on native communities. We extracted data from the literature and gathered 1111 observations. We ran mixed-effects models, using the rma.mv function from the metafor package for R, including study ID and exotic species name as independent random factors to account for the effects of study and species ID when several observations were obtained from the same publication or exotic species. The number of observations per treatment is indicated in parenthesis in all the graphs and at every treatment level.
Research sample	An exhaustive literature search was performed in ISI Web of Knowledge using the following terms: (invader OR exotic OR invasive OR non-native OR alien) NEAR/5 (impact OR effect OR influence OR consequence*) AND (marine OR coastal OR sea OR estuar* OR ocean) NOT (lake OR stream OR freshwater OR terrestrial). A search performed on May 4 2016 resulted in 1012 publications.
Sampling strategy	In order to extract data from the published literature, we created a set of study selection criteria and a protocol for data extraction from the literature
Data collection	All authors (11 in total), except CMD, extracted data and information from published literature. Each of the 1012 references was randomly ordered and given an original reference number, and authors extracted information from a set of papers (all authors extracted information from 60 papers except AA and NRG that extracted information from 200+ papers). Agreement between authors on the application of criteria was achieved after discussion and clarification of study selection criteria against particular studies. An exhaustive protocol was then created to extract data from publications. Agreement on data extraction between authors was attained by extracting information using a common subset of references (n=2), followed by an amendment of the protocol for data extraction.
Timing and spatial scale	Data from publications was extracted from November 2016 to April 2017.
Data exclusions	Only studies that met our study criteria (see methods) were included in the analysis
Reproducibility	We did not run experiments on this study
Randomization	We did not run experiments on this study
Blinding	A set of study criteria were followed to collect the data from the published literature
Did the study involve field work?	☐ Yes ☒ No

Reporting for specific materials, systems and methods

We require information from authors about some types of materials, experimental systems and methods used in many studies. Here, indicate whether each material, system or method listed is relevant to your study. If you are not sure if a list item applies to your research, read the appropriate section before selecting a response.

Materials & experimental systems

n/a	Involved in the study
☒	☐ Antibodies
☒	☐ Eukaryotic cell lines
☒	☐ Palaeontology
☒	☐ Animals and other organisms
☒	☐ Human research participants
☒	☐ Clinical data

Methods

n/a	Involved in the study
☒	☐ ChIP-seq
☒	☐ Flow cytometry
☒	☐ MRI-based neuroimaging

Nature Research, brought to you courtesy of Springer Nature Limited (“Nature Research”)

Terms and Conditions

Nature Research supports a reasonable amount of sharing of content by authors, subscribers and authorised or authenticated users (“Users”), for small-scale personal, non-commercial use provided that you respect and maintain all copyright, trade and service marks and other proprietary notices. By accessing, viewing or using the nature content you agree to these terms of use (“Terms”). For these purposes, Nature Research considers academic use (by researchers and students) to be non-commercial.

These Terms are supplementary and will apply in addition to any applicable website terms and conditions, a relevant site licence or a personal subscription. These Terms will prevail over any conflict or ambiguity with regards to the terms, a site licence or a personal subscription (to the extent of the conflict or ambiguity only). By sharing, or receiving the content from a shared source, Users agree to be bound by these Terms.

We collect and use personal data to provide access to the nature content. ResearchGate may also use these personal data internally within ResearchGate and share it with Nature Research, in an anonymised way, for purposes of tracking, analysis and reporting. Nature Research will not otherwise disclose your personal data unless we have your permission as detailed in the Privacy Policy.

Users and the recipients of the nature content may not:

1. use the nature content for the purpose of providing other users with access to content on a regular or large scale basis or as a means to circumvent access control;
2. use the nature content where to do so would be considered a criminal or statutory offence in any jurisdiction, or gives rise to civil liability, or is otherwise unlawful;
3. falsely or misleadingly imply or suggest endorsement, approval, sponsorship, or association unless explicitly agreed to by either Nature Research or ResearchGate in writing;
4. use bots or other automated methods to access the nature content or redirect messages; or
5. override any security feature or exclusionary protocol.

These terms of use are reviewed regularly and may be amended at any time. We are not obligated to publish any information or content and may remove it or features or functionality at our sole discretion, at any time with or without notice. We may revoke this licence to you at any time and remove access to any copies of the shared content which have been saved.

Sharing of the nature content may not be done in order to create substitute for our own products or services or a systematic database of our content. Furthermore, we do not allow the creation of a product or service that creates revenue, royalties, rent or income from our content or its inclusion as part of a paid for service or for other commercial gain. Nature content cannot be used for inter-library loans and librarians may not upload nature content on a large scale into their, or any other, institutional repository.

To the fullest extent permitted by law Nature Research makes no warranties, representations or guarantees to Users, either express or implied with respect to the nature content and all parties disclaim and waive any implied warranties or warranties imposed by law, including merchantability or fitness for any particular purpose.

Please note that these rights do not automatically extend to content, data or other material published by Nature Research that we license from third parties.

If you intend to distribute our content to a wider audience on a regular basis or in any other manner not expressly permitted by these Terms please contact us at

onlineservice@springernature.com

The Nature trademark is a registered trademark of Springer Nature Limited.