Title:

Machine Learning Reveals Spatiotemporal Genome Evolution in Asian Rice Domestication

Authors:

Hajime Ohyanagi, Kosuke Goto, S3nia Negr3o, Rod A. Wing, Mark A. Tester, Kenneth L. McNally, Vladimir B. Bajic, Katsuhiko Mineta, Takashi Gojobori*

Authors' Affiliations:

King Abdullah University of Science and Technology (KAUST), Computational Bioscience Research Center (CBRC), Biological and Environmental Sciences & Engineering Division (BESE), Thuwal, 23955-6900, Saudi Arabia

Hajime Ohyanagi, Kosuke Goto & Takashi Gojobori

School of Biology and Environmental Science, University College Dublin, Belfield, Ireland

S3nia Negr3o

King Abdullah University of Science and Technology (KAUST), Biological and Environmental Sciences & Engineering Division (BESE), Thuwal, 23955-6900, Saudi Arabia

Rod A. Wing & Mark A. Tester

King Abdullah University of Science and Technology (KAUST), Computational Bioscience Research Center (CBRC), Computer, Electrical and Mathematical Sciences & Engineering Division (CEMSE), Thuwal, 23955-6900, Saudi Arabia

Vladimir B. Bajic & Katsuhiko Mineta

International Rice Research Institute, Manila, Philippines

Kenneth L. McNally

***Corresponding Author:**

Takashi Gojobori, Distinguished Professor,

King Abdullah University of Science and Technology (KAUST), Thuwal 23955-6900, Saudi Arabia

E-mail: takashi.gojobori@kaust.edu.sa, Phone: +966-12-808-2893

Domestication is anthropogenic evolution that fulfills mankind's critical food demand. As such, elucidating the molecular mechanisms behind this process promotes the development of future new crops. With the aim of understanding the

whole domestication process of Asian rice and by employing the *Oryza sativa* subspecies (*indica* and *japonica*) as an Asian rice domestication model, we scrutinized genomic introgressions between them as traces of domestication. Here we show the genome-wide introgressive region (IR) map of Asian rice, by utilizing 4,587 accession genotypes with a stable outgroup species, particularly at the finest resolution through a machine learning-aided method. The IR map revealed that 14.2% of the rice genome consists of IRs, including both wide IRs (recent) and narrow IRs (ancient). This introgressive landscape with their time calibration indicates that introgression events happened in multiple genomic regions over multiple periods. From the correspondence between our wide IRs and so-called Selective Sweep Regions, we provide a definitive answer to a long-standing controversy in plant science: Asian rice phylogeny appears to depend on which regions and time frames are examined.

Rice is one of the most essential crops to humankind, playing a critical role in food security¹. Since it has been domesticated to fit it to humanity's needs, its genome holds the secrets to ancient and modern agricultural practices, which can serve as an informative reference for future breeding practices. Rice domestication history can be divided into three independent episodes: *Oryza nivara* (also known as annual *O. rufipogon* or Or-I) and *O. rufipogon* in Asia that led to Asian rice (*O. sativa* L.)², *O. barthii* that was domesticated by early African farmers around 3,000 years ago and led to African rice (*O. glaberrima* Steud.)³, and a New World rice domestication process by Amazon farmers around 4,000 years ago that occurred in South America⁴. In particular,

the Asian domesticated rice (*O. sativa*) is the most prominent species in the genus *Oryza*, which has served as the major staple crop in most Asian countries for millennia.

Among these three domesticated rice species, Asian rice (*O. sativa*) and its origins have been the most intensively studied and continue to be debated in both archeological and genetic research areas⁵⁻²⁰. In short, two conflicting domestication hypotheses have been proposed: 1) a single domestication process where a single subspecies (either *indica* or *japonica*) was first domesticated from a wild rice, while the other arose from a hybridization with another wild rice species; and 2) independent domestication processes where different species of *O. nivara* and *O. rufipogon* with distinct Asian origins gave rise to different domesticated subspecies.

A comprehensive SNP-based genomic phylogeny (*i.e.*, a genomic phylogeny as a whole) clearly shows that at least two origins of *O. sativa* subspecies exist¹⁴, *i.e.*, *O. sativa* ssp. *indica* and *O. nivara* cluster with each other, while *O. sativa* ssp. *japonica* and *O. rufipogon* make another cluster. However, this is just a subspecies phylogeny, which does not reflect the domestication history. To trace back the history, plant scientists have been focusing on their own self-defining genomic entities, *e.g.*, domestication-associated gene regions (with flanking upstream/downstream regions), Selective Sweep Regions¹⁴, Co-located Low-Density Genomic Regions¹⁰, transposable elements⁶, microsatellites¹², and so forth. In other words, there have been multiple definitions for domestication-derived regions. Meanwhile, phylogenies inferred by plant scientists do not always agree with one another, either supporting theories 1) or 2). In fact, the domesticated Asian rice accessions have supposedly introduced agronomically advantageous traits from one subspecies to another during the domestication process^{7,9,20-22}. Therefore, their genomes

are presumed to be mosaics since they have been exchanging alleles over introgression events throughout history. In this sense, the controversy over the origins of rice domestication arose from the disagreed domestication-derived regions. Moreover, the window size studied is a critical factor in the controversy. In our study, the phylogenetic analysis of a domestication-associated gene with variable lengths of upstream/downstream flanking regions, as also shown in the result of Choi & Purugganan⁸ showed that the gene window size profoundly affects the resultant gene phylogenies (shown in **Fig. 2e, f, g, h, and i**, details will be described in **Consequence of Analysis Window Size**).

Given that introgression events are representative of human intervention (*i.e.*, the domestication process), our simple and robust rationale is not to focus on particular genomic regions, but rather to exhaustively detect any introgressive regions (IRs) between subspecies as traceable signs of domestication, employing windows with as fine a resolution as possible. In keeping with this notion, we present not only gene-by-gene introgressive states but also a genome-wide IR map between *O. sativa* ssp. *indica* and ssp. *japonica* at the finest resolution using an efficient machine learning model, with the aim of revealing the whole domestication process of Asian rice.

Results

Invention of *Distance Difference (DD)* to Detect Introgressions

To capture the entire introgressive landscape of domesticated Asian rice genomes using a large-scale genotype set (**Fig. 1a and b**), we needed to overcome three major difficulties described in the **Methods**. In short, i) the low density of rice genotypes, ii) over-diversity

within each subspecies (**Fig. 1c**), and iii) the instability of an outgroup. To overcome these challenges, we employed 14x coverage genotypes supplied by the 3,000 Rice Genomes Project²²⁻²⁵, a median 10th subset extraction from the comprehensive dataset, and a reproductively isolated accession of *O. punctata* (BB diploid, 2n=24, with African geographical origin)²⁶ as an outgroup species. For more details, see **Methods**.

Each domesticated subpopulation has its own particular evolutionary rate²⁷. Therefore, each of *indica* and *japonica* subpopulations should show, to some extent, different genetic distances to an outgroup (a wild rice accession), since they have been separated from each other for a length of time (**Fig. 2a**) with the assumption that any inter-subspecies cross (*i.e.*, an introgression) has not occurred. On the other hand, they will show more similar genetic distances to the outgroup when an inter-subspecies cross has occurred (**Fig. 2b**). In particular, subspecies in domesticated plants have been artificially forced to make inter-subspecies crossings in order to introduce agronomically important traits, thereby particular regions of their genomes must be strongly affected by the decrease in difference of genetic distance (distance difference).

Even though this decrease may disturb an accurate inference of genetic phylogeny of rice subspecies and wild relatives, it can be paradoxically utilized as an index of introgression, *i.e.*, once a decrease is observed, it is a possible sign of an introgression event. To distinguish IRs from non-IRs (**Fig. 2a** and **b**), we conceptually defined *DD* (*Distance Difference* to the outgroup: A unit is number of substitutions per nucleotide site) as:

$$DD = |F84(\text{outgroup to } indica) - F84(\text{outgroup to } japonica)|$$

(*) F84 = Felsenstein84 nucleotide genetic distance²⁸

Here, the regions with smaller *DDs* represent IRs, while the regions with larger *DDs* represent non-IRs. For more details, see **Methods**. Note that because IRs at the very early stage of domestication will not show enough decrease in *DDs*, IRs of very ancient origin are out of scope of this method.

Incoherent Introgressive States of Domestication-associated Genes (D-genes)

Based on the logic above, we firstly aimed to determine *DDs* of 25 manually curated domestication-associated genes (D-genes, **Fig. 2c**) as indices of their introgressive states. To achieve the best accuracy in this limited scale analysis, we constructed 25 gene-by-gene phylogenetic trees without any flanking upstream/downstream regions, and we visually inspected their *DDs* thoroughly, to determine whether *indica* and *japonica* show a similar genetic distance to the outgroup, or different genetic distances to the outgroup. Our results show that incoherent introgressive states of D-gene regions, *i.e.* nine D-genes (*Bh4*, *C1*, *GAD1*, *LABA1*, *LG1*, *Progl*, *qSW5*, *Rc*, and *sh4*) out of 25, are introgressive (regardless of the direction), whereas 14 D-genes (*BADH2*, *Bph14*, *DPL2*, *Ehd1*, *Ghd7*, *Gn1a*, *GS3*, *GW2*, *Phr1*, *qSH1*, *Rd*, *sdl*, *tb1*, and *waxy*) are not (**Fig. 2c and d**, yellow = non-introgressive, red = introgressive, full size phylogenetic tree pictures with detailed color system are shown in **Supplementary Fig. 1**). *Hd1* and *S5* have status-undetermined. Through a statistical analysis (**Supplementary Table 2**), we found significant enrichment in the introgressed proportion of D-genes to that of the control (all genes) by a G-test of Goodness-of-Fit (P -value < 0.000121). However, the use of this approach with the D-genes did not yield a coherent introgressive state, thus providing little insight into the history of Asian rice at the present stage, emphasizing the need for a more systematic

approach to decipher the genome-wide status of Asian rice. For a further interpretation of these results, see **Discussion**.

Consequence of Analysis Window Size

Because the introgressive states of D-genes did not give clear answer to the history of Asian rice, we consequently explored the genome-wide introgressive states in a manner involving significantly more computational resource costs and time.

Our phylogenetic analysis for one of the D-genes (*LGI*) with variable lengths of flanking upstream/downstream regions (**Fig. 2e** : CDS only, **f** : +5kb-upstream/+5kb-downstream, **g** : +10kb-upstream/+10kb-downstream, **h** : +20kb-upstream/+20kb-downstream, and **i** : +100kb-upstream/+100kb-downstream, respectively) clearly shows that region size heavily affects the resultant phylogeny. More precisely, a narrow region (CDS only) showed a monophyletic topology of *LGI* between *indica* and *japonica*, suggesting that it is introgressive (**Fig. 2e**), while wider region analyses resulted in a polyphyletic relationship resembling non-introgressive state (**Fig. 2g, h, and i**). Full-size tree pictures with a detailed color system are shown in **Supplementary Fig. 2**. Therefore, we emphasize that window size is important; the window size setup in genome-wide analysis is significant when we are dealing with phylogenies of domesticated Asian rice at the loci-level.

The genome of domesticated Asian rice is polyphyletic as a whole, yet not always so at the loci-level^{7,9,14,20-22}. This is in line with our inconsistent result (**Fig. 2e, f, g, h, and i**), indicating that a narrower window setup leads to a more accurate inference of phylogeny at the loci-level. Moreover, adopting a wider window size is inaccurate

because it does not deal with phylogenies at the loci-level^{7,9,21,22}, but rather with a whole-genome phylogeny. Furthermore, our preliminary analyses with imputed 4,587 accession genotypes unsuccessfully resulted in similar inconsistent phylogenetic relationships, indicating that methods based on the haplotype linkages in certain-sized regions (*e.g.*, wider window size; imputation) are not suitable for exploring the phylogenies at the loci-level.

Genome-wide Introgressive States Occur in Blocks

We developed a machine learning classification model to distinguish the non-introgressed windows (**Fig. 2a**) from introgressed windows (**Fig. 2b**) computationally. This is to streamline a time-consuming visual inspection (*e.g.*, if we set 1kb windows all along the rice genome (~373Mb), we would need to handle ~373,000 windows). Another merit for adopting a machine learning-aided method is that it is free from null hypotheses and *P*-value-dependent approach²⁹. As shown in **Methods**, we achieved 96.1% accuracy for the binary classifier by the Breiman & Cutler's Random Forest Algorithm³⁰, and thus we adopted it for further analyses.

Initially, we scanned the rice genome and developed an *indica* - *japonica* IR map at 100kb-resolution using a random forest classification model (for details, see **Methods**), but it was blocky and the introgressive landscape was still veiled, shown in **Fig. 3a** showing chromosome 1. We then increased the resolution to 20kb- (**Fig. 3b**), 10kb- (**Fig. 3c**), 5kb- (**Fig. 3d**), and finally to 1kb (**Fig. 3e**). The 1kb-resolution IR map produced a sharp image that discriminate introgressive states at the gene loci-level along the entire genome (IR maps for chromosome 2 to chromosome 12 are shown in **Supplementary**

Fig. 3). We identified large amounts of IR bands all along the genome (**Fig. 3e** and **Supplementary Fig. 3**). Surprisingly, we determined that 14.2% of genomic contents are introgressive (**Fig. 4a**). In addition, the IRs are not uniformly distributed, but rather unevenly located in blocks (**Fig. 3e** and **Supplementary Fig. 3**). To be precise, there are several major wide IRs in each chromosome, while thousands of narrow IRs are scattered all over the genome (**Fig. 3e** and **Supplementary Fig. 3**), suggesting that there are multiple genetic backgrounds behind the introgressions.

Non-uniform Ages of Introgressions

Now that we have established that a substantial amount (14.2%) of the genetic contents has been exchanged between *indica* and *japonica* subpopulations, we aimed to uncover what the biased introgressive pattern (**Fig. 3e** and **Supplementary Fig. 3**) means. By plotting the window proportions of particular *DD*s, we observed apparent non-uniform genetic backgrounds (**Fig. 4b**). We propose that these multiple genetic backgrounds correspond to multiple classes of IRs, and that wide IRs and narrow IRs have different *DD* values. To test our proposal, we operationally and precisely defined two IR classes according to the dimensional continuity of IR windows, with wide IRs ($\geq 40\text{kb}$) and narrow IRs ($=1\text{kb}$), and explored their *DD*s. The genomic positions of the wide IRs are shown in **Supplementary Table 3**. The results show that wide IRs have a small *DD* of 5.89×10^{-6} substitutions/site, on average for all chromosomes, and narrow IRs have roughly 100 times larger *DD* than wide IRs (5.84×10^{-4} substitutions/site). Non-IRs show a much larger *DD* (1.71×10^{-3} substitutions/site) (**Fig. 4a** shows the average for all chromosomes; results for each chromosome are shown in **Supplementary Table 4**). This

similar trend of *DD* can also be observed in the continuous-valued histogram (continuity of IR windows; from one-IR to 15-IRs) shown in **Supplementary Fig. 4**.

When we roughly extrapolate the *indica-japonica* divergence time to 500,000 years ago^{7,26} (**Fig. 5**, non-IRs), we can then estimate that the wide IRs are approximately 1,700 years old, whereas the narrow IRs are approximately 170,000 years old (**Fig. 5**). Hence, we concluded that the wide IRs are relatively recently formed, while the narrow IRs have existed for considerably longer time.

Correspondence between Wide IRs and Selective Sweep Regions

To gain insight into the history of the domestication of Asian rice and to address the controversy on the origins of this domestication, we compare the genomic locations of our IRs with those of previously reported domestication-associated genomic entities, namely; Selective Sweep Regions (SSRs)¹⁴ and Co-located Low-Density Genomic Regions (CLDGRs)¹⁰. We re-computed these previously described SSRs and CLDGRs^{10,14} with our 4,587 rice accessions dataset (**Fig. 1a**) onto the Os-Nipponbare-Reference-IRGSP-1.0 reference genome (see **Methods** for more details), as shown in parallel with our IRs in **Fig. 3e, f, and g** and **Supplementary Fig. 3** (red lines: SSRs, blue lines: CLDGRs). Interestingly, our results show that the SSRs correspond well with our IRs, in particular with wide IRs (*i.e.*, young IRs), suggesting that the SSRs capture recently happened events of introgression. However, in contrast, we observed less correspondence between the CLDGRs and our wide IRs (**Fig. 3e, f and g** and **Supplementary Fig. 3**), suggesting that CLDGRs do not deal with such events of introgression. We discuss these patterns of correspondence further in **Discussion**.

Discussion

The genetic structure of domesticated Asian rice includes five major subpopulations³¹. A recently study shows that it can be subdivided into nine detailed subpopulations²². Ancient Chinese literature reported as early as the Han dynasty in China (100 AD) the existence of two ecogeographical rice groups called ‘*Xian* (or *Hsien*)’ and ‘*Geng* (or *Keng*)’, which correspond to *indica* and *japonica* subpopulations, respectively^{32,33}. This indicates that *indica* and *japonica* subpopulations have been cultivated for at least around 2000 years, being exposed to human intervention for a long time. For this reason, we chose these two subspecies as the best model for studying the domestication of Asian rice. In addition, we considered these subspecies because of the availability of high quality sequenced genomes³⁴, curated genome annotations³⁵, more than 3,000 re-sequenced closely-related accessions²²⁻²⁵, and additional quality reference genomes (IR8 for *indica* and N22 for *aus*), together with eight wild *Oryza* species²⁶.

Archeological evidence indicates that Asian rice was first domesticated in the early Holocene period ca. 9000^{5,36}, but Asian rice domestication and its origin is still a matter of ongoing debate in both archeological and genetic research areas⁵⁻²⁰. Plant scientists have expected that the availability of whole-genome sequences of domesticated Asian rice, its wild relatives, and ancient rice³⁷, would provide a resolution to this long-standing debate, yet the controversy is ongoing, because the genetic structure of rice genomes turned out to be more complex than expected. In the two research studies of evolutionary origins of domesticated Asian rice^{10,14}, they analyzed a single dataset, which included 1,529 genotypes of wild and domesticated rice^{14,38}, leading to opposite

domestication scenarios. More recently, the same dataset was re-evaluated by the third team, who suggested that rice originated from multiple populations of *O. rufipogon* (and/or *O. nivara*): *De novo* domestication only occurred once where domestication alleles were introgressed predominantly from *japonica* into *indica* subpopulations^{7,8}.

In this study, we explore possible events of introgression between subspecies, considering them as traceable signs of domestication (**Fig. 2a and b**). We capture the genome-wide IR map between *O. sativa* ssp. *indica* and *japonica*, with the aim of encapsulating the entire history of Asian rice domestication. We exhaustively scan and reveal the genome-wide introgressive landscape between *indica* and *japonica* at the finest resolution using a machine learning classification model (**Fig. 3e and Supplementary Fig. 3**). Our results show that a surprisingly large proportion of the rice genome (14.2%) consists of wide and narrow traces of introgression between *indica* and *japonica* (**Fig. 4a**). This suggests that even after the initial diversification of Asian rice roughly 500,000 years ago^{7,26}, *indica* and *japonica* subpopulations have been exchanging alleles between each other.

In addition, we explore the introgressive state of 25 D-gene regions. We detected a significantly large number of D-genes upon IRs, though not all of D-genes (**Supplementary Table 2**), which shows that introgression was a major but non-exclusive molecular mechanism for D-gene propagation. In other words, some D-genes moved along the introgressive flows (regardless of the direction). Note that not all D-genes were mobilized via introgression events.

We also observed that, in terms of *DD*, the wide IRs have emerged recently, whereas the narrow IRs have existed for a much longer time (**Fig. 4a and Supplementary Table**

4). This mosaic introgressive landscape in terms of time (**Fig. 5**) clearly indicates that multiple introgression events between subpopulations have taken place multiple times throughout history (**Fig. 6**). In each of these events, the brand-new wide IRs would comprise some beneficial alleles and many non-beneficial alleles. The beneficial alleles would have been selected for and fixed in recipient subpopulations, while the non-beneficial alleles would not have been fixed in the subpopulation. Thus, the genomic regions with less advantageous alleles would have been replaced, eventually disappearing following subsequent multiple backcrosses within the recipient subpopulation (**Fig. 6**). Such genome dynamics can look like “sequentially built sandcastles” on a beach, whereby newly built castles are still intact, while the older castles are already beginning to crumble (**Fig. 6**). From the standpoint of our Sandcastles Model, the vast majority of detected IRs correspond to non-beneficial alleles, which are mostly derived by hitchhiking effects (**Fig. 6**). Extrapolating the *indica-japonica* divergence time (500,000 years ago corresponds to 1.71×10^{-3} substitutions/site in terms of *DD*)^{7,26}, we can estimate that the narrow and wide IRs are approximately 170,000 and 1,700 years old, respectively (**Fig. 5**). This is consistent with the Asian rice domestication timeline: It was initially domesticated in the early Holocene period^{5,36} and has been maintained for at least about 2,000 years^{32,33}.

The history, particularly the first origins of Asian rice domestication has long been a subject of active discussion in plant biology⁵⁻²⁰. Studies have focused specifically on the domestication-associated regions that presumably reflect the domestication process in rice genomes. Those regions are typically defined by D-gene loci with flanking upstream/downstream regions, SSRs, and CLDGRs. As an inevitable consequence in

those studies^{10,14}, the definition of domestication-associated regions heavily affected the reconstructed genetic phylogenies and the conclusions.

In this study, by employing highly dense SNP information and a machine learning modeling approach, we elucidated a 1kb-resolution IR map and found that the young IRs were well co-localized with SSRs¹⁴, but not with CLDGRs¹⁰. In terms of population genetics, each of the IRs and SSRs were derived from a different population statistic, *i.e.*, IRs were detected by a decrease in genetic distance difference to the wild relative (*DD*), while SSRs were inferred by a decrease in nucleotide diversity (Π) compared to that of the wild relatives. However, since gene introgressions will act in the direction of decreasing Π in the domesticated population, $\Pi(\text{wild}) / \Pi(\text{domesticated})$ will have a higher value, and thus the correspondence between SSRs and young IRs makes sense. In terms of molecular phylogeny, the young IRs show a quite higher genetic identity between *indica* and *japonica*, which could lead to monophyly (**Fig. 5**, bottom right panel). On the other hand, the old IRs and non-IRs tend to represent more genetic divergence, which seems to be polyphyletic (**Fig. 5**, bottom left panel and top panel). Hence the discrepancy in results from the two previous studies^{10,11,14,15} can be reasonably explained by our Sandcastles Model (**Fig. 6**), *i.e.*, one study focused on the new castles (young IRs)¹⁴, while the other did not¹⁰. We propose that focusing on certain-sized genomic regions is a misleading way to understand the primal origins of domesticated life, because these regions may contain recently built young IR blocks (**Fig. 6**).

If we pursue the very first ancestor of domesticated Asian rice, we need to eliminate carefully the SSR-like entities that overlap with the young IR blocks from the analysis, because they are recent and do not reflect ancient domestication history. We should

instead probe into other SSRs (old SSRs) and/or old IRs in the genome, which are the true traces of ancient domestication history. However, since the extant domesticated subspecies (*e.g.*, *indica* and *japonica*), and closely-related wild relatives (*O. nivara* and *O. rufipogon*) as well, are not yet completely isolated reproductively^{39,40}, the subspecies boundary of initial Asian rice subpopulations should be much more permeable. In that sense, it may not be meaningful to explore whether the initial domesticated rice individual(s) is in a single subpopulation or are in multiple subpopulations, because they were too permeable in terms of conventional taxonomy. Leastwise, be that as the initial domestication might happen multiple times, these were in a single population *sensu lato*.

In summary, we have determined that a surprisingly large proportion (14.2%) of genetic contents has been exchanged between *indica* and *japonica* subpopulations. We have also demonstrated that introgression events have happened in multiple genomic regions over multiple periods throughout the history of domesticated Asian rice, revealing the complex spatiotemporal genome dynamics in Asian rice domestication. Concomitantly, we settle the major controversy in plant science between two hypotheses⁵⁻²⁰ using our Sandcastles Model, *i.e.*, each study was focusing on a different genomic region of a different era. Moreover, because the IRs contribute to the domestication process in a proactive manner, our IR map provides a unique reference for potential target loci in breeding of rice. This gives insight into new breeding designs and practices based on the introgressive genomic map.

Methods summary

The genotypes of domesticated and wild rice accessions were all retrieved from publicly

available databases. The full methods and any associated information are available in the online version of the paper.

Methods

Reference genome. For the reference genome sequences and reference genome annotations, the reference Nipponbare genome Os-Nipponbare-Reference-IRGSP-1.0 (*O. sativa* ssp. *japonica* cv. Nipponbare)⁴¹; hereinafter referred to as Nipponbare RefSeq and CGSNL annotations served in RAP-DB⁴² were employed, respectively.

Domestication-associated genes (D-genes). Based on our literature survey, we manually selected and curated a total of 25 D-genes (**Fig. 2c**) for this study. The selection criteria were based on agronomically beneficial effects of genes selected.

Issues on rice genotypes. In particular, we focused our analyses on two *O. sativa* subspecies, ssp. *indica* and ssp. *japonica*, as an Asian rice domestication model. Despite multiple studies conducted to explore the history of Asian rice introgression and domestication with large-scale accessions datasets including *indica* and *japonica*^{8-10,14,21,22}, their genome-wide scanning procedures have been performed using relatively large window size setups (5kb -100kb). The importance of window size in such analyses are outlined in this study (**Fig. 2e, f, g, h, and i**) and also in Choi & Purugganan⁸, but due to the low SNPs density (56.4% missing data rate) in the dataset^{14,38}, the issue of window size had not yet been overcome. Another problem is that each *indica* and *japonica* subpopulation contains a significant amount of genetic diversity^{14,22,31}, or rather, some subspecies accessions can be intermediate accessions between the two subspecies since these subpopulations are not yet completely reproductively isolated from

each other³⁹. In fact, both *indica* and *japonica* subpopulations show a certain degree of phenotypic diversity, including some intermediate traits (**Fig. 1c**). Consequently, when taking all the *indica* and *japonica* accessions into account, the conclusion may be ambiguous because of the intermediate states of genetic distance. The final issue to be overcome when we trace back the domestication history of Asian rice is to choose which species to use as an outgroup. It is widely believed that *O. nivara* and *O. rufipogon* are the immediate ancestors of ssp. *indica* and ssp. *japonica*, respectively². However, those wild rice species are still able to intermate with *O. sativa*⁴⁰; thus, the genetic distance between those wild rice species and *O. sativa* could be underestimated in introgressive regions. Hence, those wild rice species are not always suitable for outgroup species in phylogenetic analysis. Our preliminary gene-by-gene phylogenetic analyses with the 3,000 Rice Genomes Project²²⁻²⁵, higher coverage wilds^{26,38,43,44} and the *O. punctata*²⁶ datasets (**Fig. 1a**, in total 3,060 accessions) aimed to assess the suitability of *O. nivara*, *O. rufipogon*, *O. glaberrima*, *O. barthii*, *O. glumaepatula* and *O. punctata* as outgroup species for this study (**Supplementary Fig. 5**). Our analyses showed that in some cases (e.g. *Gn1a*, *LGI1*, *Phr1*, and *qSH1*) (**Supplementary Fig. 5i, n, o and q**), a close-relatives (*O. rufipogon* or *O. nivara*) can serve as an outgroup species. However, in most cases, they are not suitable for an outgroup since they are not genetically isolated from domesticated rice (**Supplementary Fig. 5**).

Solutions on rice genotype issues. To develop an accurate high-resolution (up to 1kb window width) map of Asian rice introgression in a reasonable manner, we needed to address the above-mentioned three problems: i) the low density of rice genotypes, ii) over-diversity within each subspecies, and iii) the instability of outgroup. With the aim of

achieving good quality and quantity of rice genotypes, we collected imputation-free ~14x coverage genotypes of 3,024 rice cultivars (**Fig. 1a**) from the 3,000 Rice Genomes Project²²⁻²⁵, in conjunction with other publicly available genotypes (**Fig. 1a**). We appropriately converted their genomic coordinates to that of the Nipponbare RefSeq as described³⁸ when needed. We performed genomic imputation with the Beagle program⁴⁵ in two batches (wild/domesticated) separately and exclusively on the 4,553 accessions only for the purpose of SSRs and CLDGRs re-computation (**Fig. 1a**), but not on any other accession datasets. The core dataset (**Fig. 1a**, 3,025 accessions) contained 1,712 *indica* and 833 *japonica* accessions with a missing genotype rate of 15.0% on average. Then, to overcome the effect of intra-subspecies divergence, we dynamically picked up median 10th accessions from *indica* and *japonica* window by window (see **Introgressive Regions (IRs) detection**). Finally, to adopt an appropriate outgroup species in our study, based on preliminary gene-by-gene phylogenetic analyses (**Supplementary Fig. 5**), we exclusively employed the *O. punctata* (IRGC105690, BB diploid, 2n=24, geographical origin: Africa)²⁶ only, with the assumption that it has been mostly reproductively isolated from *O. sativa* populations. We can ignore the underestimate effect of nucleotide distance due to possible introgression events between *O. sativa* and *O. punctata* (**Supplementary Fig. 5**).

Mapping and SNPs calling. We first quality inspected all short reads by FastQC (<http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>), and then we filtered out and/or trimmed out adaptor sequences and low-quality bases using Trimmomatic⁴⁶. After those preprocessing steps, we mapped the remaining reads onto the Nipponbare RefSeq using ‘bwa mem’ commands in BWA⁴⁷ with default parameters, except for the proper

insert size limitation (-w 500 or -w 800, dictated by the data source). Repeat sequences scattered within the Nipponbare RefSeq were not masked in our mapping process. Next, we called variants using the GATK⁴⁸ with a conventional best practice method (<https://software.broadinstitute.org/gatk/best-practices/>).

Phylogenetic tree construction. For window-base analysis, we generated each 1,000bp multiple alignment. For gene-by-gene analysis, we generated a multiple alignment of actual CDS for each gene (including intron regions, but not including any flanking upstream/downstream regions). All nucleotide genetic distances between domesticated rice windows/genes and outgroup windows/genes were estimated by PHYLIP-dnadist command with default parameters (Felsenstein84 distance)²⁸. We reconstructed all phylogenetic trees using the PHYLIP-neighbor command with default parameters (Neighbor-Joining method)^{28,49}. Trees were drawn by FigTree software GUI (<http://tree.bio.ed.ac.uk/software/figtree/>), rooted by *O. punctata* as the fixed outgroup.

Invention of Distance Difference (DD). Under isolated conditions, each of *indica* and *japonica* subpopulations should show different genetic distances to an outgroup (a wild rice accession) to some extent, since they have been separated for a length of time in each subpopulation (**Fig. 2a**). However, they will show unexpectedly similar genetic distance to an outgroup when an inter-subspecies cross (*i.e.* introgression) has occurred recently (**Fig. 2b**). Together with incomplete lineage sorting and other possible situations^{50,51}, this is one of the reasons why a particular gene phylogeny does not always agree with the (sub)species phylogeny. Here we conceptually define *DD* (genetic *Distance Difference* to the outgroup) as;

$$DD = |\text{F84 (outgroup to } indica) - \text{F84 (outgroup to } japonica)| .$$

(*) F84 = Felsenstein84 nucleotide genetic distance²⁸

Here, smaller *DD*s represent IRs, while larger *DD*s mean that those are non-IRs. Note that IRs happened in the initial period of domestication will not show enough decrease in *DD*, hence such IRs are out of scope of this method. In terms of population genetics, we have multiple *indica* accessions and multiple *japonica* accessions, and each subpopulation includes much genetic diversity (see **Issues on rice genotypes**). To overcome the undesirable effect on intra-subspecies over-diversity in terms of nucleotide distance to the outgroup, we dynamically chose the median 10th accessions from *indica* (172 accessions) window by window (or gene by gene), and median 10th accessions from *japonica* (84 accessions) window by window (or gene by gene), respectively. They are representative subpopulations in each window (or each gene) in the sense that the most mediocre members reflect the profile of population. Therefore, the actual *DD* value is not computed by a single *indica* accession and a single *japonica* accession. Instead, it is computed by the average form of median 10th accessions of *indica*, and by the average form of median 10th accessions of *japonica*. Hence, the actual formula for *DD* is;

$$DD = \left| \frac{\sum_{indica}^{median10th} F84(outgroup\ to\ indica)}{172} - \frac{\sum_{japonica}^{median10th} F84(outgroup\ to\ japonica)}{84} \right|.$$

(*) F84 = Felsenstein84 nucleotide genetic distance²⁸

Introgressive Regions (IRs) detection. For the gene-by-gene analysis, we conducted visual phylogeny inspection (**Fig. 2** and **Supplementary Fig. 1**). For the window-based analysis, although visual inspection of each window phylogeny would give the best accuracy, it is too time consuming. We thus aimed to computationally distinguish the non-introgressed windows (**Fig. 2a**) from the introgressed windows (**Fig. 2b**) by the use of a binary classifier through Breiman & Cutler's Random Forest Algorithm³⁰. The

accuracy of the binary classifier was 96.1%, as determined by a 10-fold cross validation (for more details, see **Optimization of machine learning models**). The 1kb resolution machine learning classification result showed that 14.2% of the rice genome was introgressive, and 50.0% was non-introgressive (was excluded 35.8% from the analysis and marked as status-undetermined, for reasons outlined below) (**Fig. 4a**). In the window-based analysis, we excluded windows that have less alignable length with the outgroup (<5% of the window region, *i.e.* <50bp in the case of the 1kb window setup). We also excluded windows with no genetic difference (*i.e.*, no SNP) from any of the *indica/japonica* accessions to the outgroup at all. Those windows are shown as gray windows (**Fig. 3** and **Supplementary Fig. 3**).

Training of machine learning models. For the training dataset of machine learning classification models, we firstly conducted visual phylogeny inspection for randomly chosen 640 1kb-windows (~0.267% of total phylogeny determined windows, see **Fig. 4a**), and we identified 114 windows as IRs and 526 windows as non-IRs. We then balanced the ratio between positive cases (IRs) and negative cases (non-IRs) in 114 IRs and randomly sub-sampled 114 non-IRs, respectively, and these 228 cases were finally used as the actual training dataset for generating the classification models.

Optimization of machine learning models. For the features used to develop the classification models, we extracted the nucleotide distance matrices for median 10th 257 accessions (172 *indica*, 84 *japonica*, and 1 outgroup). Since the $257^2 = 66,049$ variables were too computationally demanding, we reduced the variables by equal subsampling to 50 accessions, retaining the original variations in each subspecies (50 *indica*, 50 *japonica*, and 1 outgroup). Finally, we adopted $101^2 = 10,201$ variables as the features

for developing the classification models. In order to find the best option for our machine learning analysis, then we conducted a grid search for model parameters with a support vector machine model (with non-linear Gaussian kernel) (with parameters $C = 2, 4, 8, 16, 32, 64, 128, 256, 512, 1024$; $\sigma = 2, 4, 8, 16, 32, 64, 128, 256, 512, 1024$; 100 cases in total), and a random forest model (with parameters $ntree = 16, 32, 64, 128, 256, 512, 1024, 2048, 4096, 8192$; $mtry = 2, 4, 8, 16, 32, 64, 128, 256, 512, 1024$; 100 cases in total). We determined that the random forest model ($ntree = 512$, $mtry = 256$, accuracy = 96.1% by 10-fold cross validation, data not shown) was the best option. We implemented the support vector machine model, random forest model, and cross validation framework by R language and R packages (kernlab, randomForest, and mlr) (<https://www.r-project.org>).

Verification of the machine learning model. To verify the effectiveness of our random forest classifier, we drew an identical conclusion by adopting another statistical classification method as shown below. Assuming that the median 10th subset data are not normally distributed, we tested whether the difference between F84 (outgroup to *indica*) and F84 (outgroup to *japonica*) is statistically significant or not, using the non-parametric statistical test method (Mann-Whitney U test, P -value $< 10^{-7}$), window by window. When the null hypothesis is rejected, the window will be non-introgressive (**Fig. 2a**, significantly different). Otherwise (*i.e.*, not significantly different), it is considered a candidate for introgression (**Fig. 2b**). As noted above, although the P -value threshold is quite conservative (P -value $< 10^{-7}$), 54.8% of the rice genome (similarly to random forest model at 50.0%) was still determined as significant (*i.e.*, non-introgressive). We determined that genomic locations were introgressive similarly to the random forest

model (data not shown), and our conclusion was identical to that of the random forest model. Even if we adopted a more aggressive P -value < 0.05 , the significant (*i.e.*, non-introgressive) window percentages were still quite similar (56.4%), the genomic locations as introgressive were still similar to those of the random forest model (data not shown) and again we reached identical conclusions, thus demonstrating the robustness of our random forest model. Moreover, manual phylogeny curation of 25 gene-by-gene results was well in line with the window-based results of random forest (**Fig. 3** and **Supplementary Fig. 3**), reconfirming the accuracy of our random forest model.

Enrichment test for D-genes on IRs. We tested whether the 25 D-genes (**Fig. 2c**) are statistically significantly enriched (or depleted) on IRs or not. A G-test of Goodness-of-Fit showed statistically significant enrichment on the proportion of introgressive D-genes (9 genes) against non-introgressive D-genes (14 genes) (**Supplementary Table 2**) (2 D-genes (*Hdl* and *S5*) showed undetermined phylogeny). For the control (all genes, *i.e.*, expected proportion), we computationally determined each gene's IRs concordance when the entire gene locus was inclusively contained in any continuous IRs of 1kb resolution (introgressive = 3,498 genes: 9.24%; non-introgressive = 34,350 genes: 90.8%). The G-test was conducted with the following R script:

```
> observed      = c(9,14)
> expected.prop = c(0.0924, 0.908)
> degrees = 1
> expected.count = sum(observed)*expected.prop
> G = 2 * sum(observed * log(observed / expected.count))
> G
[1] 14.78253
> pchisq(G,df=degrees,lower.tail=FALSE)
[1] 0.0001206482
> q()
```

Re-computation of Selective Sweep Regions (SSRs) and Co-located Low-Density

Genomic Regions (CLDGRs). For the already reported domestication-associated

genomic entities (Selective Sweep Regions (SSRs)¹⁴ and Co-located Low-Density Genomic Regions (CLDGRs)¹⁰), we re-computed their SSRs and CLDGRs using our 4,587 accessions dataset (**Fig. 1a**) on the Nipponbare RefSeq, and we conducted independent permutation tests to determine the appropriate $\Pi(\text{wild}) / \Pi(\text{domesticated})$ threshold. In **Fig. 3e** and **Supplementary Fig. 3**, re-computed SSRs and CLDGRs were shown as red lines and blue lines, respectively. The re-computation procedures are summarized in **Supplementary Fig. 6** and **7**.

Data availability. All the intermediate and final analysis results in this study are available from the corresponding author upon request.

D-genes' References (will be imported to Fig. 2c):

*BADH2*⁵²
*Bh4*⁵³
*Bph14*⁵⁴
*C1*⁵⁵
*DPL2*⁵⁶
*Ehd1*⁵⁷
*GAD1*⁵⁸
*Ghd7*⁵⁹
*Gn1a*⁶⁰
*GS3*⁶¹
*GW2*⁶²
*Hd1*⁶³
*LABA1*⁶⁴
*LG1*⁶⁵
*Phr1*⁶⁶
*Prog1*⁶⁷
*qSH1*⁶⁸
*qSW5*⁶⁹
*Rc*⁷⁰
*Rd*⁷¹
*S5*⁷²

*sd1*⁷³
*sh4*⁷⁴
*tb1*⁷⁵
*waxy*⁷⁶

562

563 **References:**

- 564 1. FAO. FAO Statistical Yearbook Part3 : Feeding the world. (2013).
- 565 2. Kumagai, M., Tanaka, T., Ohyanagi, H., Hsing, Y.C. & Itoh, T. Genome Sequence of *Oryza*
- 566 Species. in *Rice Genomics, Genetics and Breeding* (eds. Sasaki, T. & Ashikari, M.) 1-20
- 567 (Springer, 2018).
- 568 3. Wang, M. *et al.* The genome sequence of African rice (*Oryza glaberrima*) and evidence
- 569 for independent domestication. *Nat Genet* **46**, 982-8 (2014).
- 570 4. Hilbert, L. *et al.* Evidence for mid-Holocene rice domestication in the Americas. *Nat Ecol*
- 571 *Evol* **1**, 1693-1698 (2017).
- 572 5. Callaway, E. Domestication: The birth of rice. *Nature* **514**, S58-9 (2014).
- 573 6. Carpentier, M.C. *et al.* Retrotranspositional landscape of Asian rice revealed by 3000
- 574 genomes. *Nat Commun* **10**, 24 (2019).
- 575 7. Choi, J.Y. *et al.* The Rice Paradox: Multiple Origins but Single Domestication in Asian
- 576 Rice. *Mol Biol Evol* **34**, 969-979 (2017).
- 577 8. Choi, J.Y. & Purugganan, M.D. Multiple Origin but Single Domestication Led to *Oryza*
- 578 *sativa*. *G3 (Bethesda)* **8**, 797-803 (2018).
- 579 9. Civan, P. & Brown, T.A. Role of genetic introgression during the evolution of cultivated
- 580 rice (*Oryza sativa* L.). *BMC Evol Biol* **18**, 57 (2018).
- 581 10. Civan, P., Craig, H., Cox, C.J. & Brown, T.A. Three geographically separate domestications
- 582 of Asian rice. *Nat Plants* **1**, 15164 (2015).
- 583 11. Civan, P., Craig, H., Cox, C.J. & Brown, T.A. Multiple domestications of Asian rice. *Nat*
- 584 *Plants* **2**, 16037 (2016).
- 585 12. Gao, L.Z. & Innan, H. Nonindependent domestication of the two rice subspecies, *Oryza*
- 586 *sativa* ssp. *indica* and ssp. *japonica*, demonstrated by multilocus microsatellites.
- 587 *Genetics* **179**, 965-76 (2008).
- 588 13. Gross, B.L. & Zhao, Z.J. Archaeological and genetic insights into the origins of
- 589 domesticated rice. *Proceedings of the National Academy of Sciences of the United States*
- 590 *of America* **111**, 6190-6197 (2014).
- 591 14. Huang, X. *et al.* A map of rice genome variation reveals the origin of cultivated rice.
- 592 *Nature* **490**, 497-501 (2012).
- 593 15. Huang, X.H. & Han, B. Rice domestication occurred through single origin and multiple
- 594 introgressions. *Nature Plants* **2**(2016).
- 595 16. Londo, J.P., Chiang, Y.C., Hung, K.H., Chiang, T.Y. & Schaal, B.A. Phylogeography of Asian
- 596 wild rice, *Oryza rufipogon*, reveals multiple independent domestications of cultivated
- 597 rice, *Oryza sativa*. *Proc Natl Acad Sci U S A* **103**, 9578-83 (2006).
- 598 17. Molina, J. *et al.* Molecular evidence for a single evolutionary origin of domesticated rice.
- 599 *Proc Natl Acad Sci U S A* **108**, 8351-6 (2011).
- 600 18. Sang, T. & Ge, S. Genetics and phylogenetics of rice domestication. *Curr Opin Genet Dev*
- 601 **17**, 533-8 (2007).

- 602 19. Vitte, C., Ishii, T., Lamy, F., Brar, D. & Panaud, O. Genomic paleontology provides
603 evidence for two distinct origins of Asian rice (*Oryza sativa* L.). *Mol Genet Genomics* **272**,
604 504-11 (2004).
- 605 20. Yang, C.C. *et al.* Independent domestication of Asian rice followed by gene flow from
606 japonica to indica. *Mol Biol Evol* **29**, 1471-9 (2012).
- 607 21. Santos, J.D. *et al.* Fine scale genomic signals of admixture and alien introgression among
608 Asian rice landraces. *Genome Biol Evol* (2019).
- 609 22. Wang, W. *et al.* Genomic variation in 3,010 diverse accessions of Asian cultivated rice.
610 *Nature* **557**, 43-49 (2018).
- 611 23. Alexandrov, N. *et al.* SNP-Seek database of SNPs derived from 3000 rice genomes.
612 *Nucleic Acids Res* **43**, D1023-7 (2015).
- 613 24. Li, J.Y., Wang, J. & Zeigler, R.S. The 3,000 rice genomes project: new opportunities and
614 challenges for future rice research. *Gigascience* **3**, 8 (2014).
- 615 25. Mansueto, L. *et al.* Rice SNP-seek database update: new SNPs, indels, and queries.
616 *Nucleic Acids Res* **45**, D1075-D1081 (2017).
- 617 26. Stein, J.C. *et al.* Genomes of 13 domesticated and wild rice relatives highlight genetic
618 conservation, turnover and innovation across the genus *Oryza*. *Nat Genet* **50**, 285-296
619 (2018).
- 620 27. Sun, X., Jia, Q., Guo, Y., Zheng, X. & Liang, K. Whole-genome analysis revealed the
621 positively selected genes during the differentiation of indica and temperate japonica
622 rice. *PLoS One* **10**, e0119239 (2015).
- 623 28. Felsenstein, J. PHYLIP - Phylogeny Inference Package (Version 3.2). *Cladistics* **5**, 164-6
624 (1989).
- 625 29. Johnson, D.H. The insignificance of statistical significance testing. *Journal of Wildlife*
626 *Management* **63**, 763-772 (1999).
- 627 30. Breiman, L. Random forests. *Machine Learning* **45**, 5-32 (2001).
- 628 31. Garris, A.J., Tai, T.H., Coburn, J., Kresovich, S. & McCouch, S. Genetic structure and
629 diversity in *Oryza sativa* L. *Genetics* **169**, 1631-8 (2005).
- 630 32. Oka, H.I. *Origin of Cultivated Rice*, (Elsevier Science, Tokyo, 1988).
- 631 33. Ting, Y. Chronological studies of the cultivation and the distribution of rice varieties,
632 Keng and Sen. *Sun Yatsen University Agronomy Bulletin* **6**, 1-32 (1949).
- 633 34. International Rice Genome Sequencing, P. The map-based sequence of the rice genome.
634 *Nature* **436**, 793-800 (2005).
- 635 35. Ohyanagi, H. *et al.* The Rice Annotation Project Database (RAP-DB): hub for *Oryza sativa*
636 ssp. japonica genome information. *Nucleic Acids Res* **34**, D741-4 (2006).
- 637 36. Zuo, X.X. *et al.* Dating rice remains through phytolith carbon-14 study reveals
638 domestication at the beginning of the Holocene. *Proceedings of the National Academy*
639 *of Sciences of the United States of America* **114**, 6486-6491 (2017).
- 640 37. Kumagai, M. *et al.* Rice Varieties in Archaic East Asia: Reduction of Its Diversity from Past
641 to Present Times. *Mol Biol Evol* **33**, 2496-505 (2016).
- 642 38. Ohyanagi, H. *et al.* OryzaGenome: Genome Diversity Database of Wild *Oryza* Species.
643 *Plant Cell Physiol* **57**, e1 (2016).
- 644 39. Guo, J. *et al.* Overcoming inter-subspecific hybrid sterility in rice by developing indica-
645 compatible japonica lines. *Sci Rep* **6**, 26878 (2016).
- 646 40. Wang, H.R., Vieira, F.G., Crawford, J.E., Chu, C.C. & Nielsen, R. Asian wild rice is a hybrid
647 swarm with extensive gene flow and feralization from domesticated rice. *Genome*
648 *Research* **27**, 1029-1038 (2017).

649 41. Kawahara, Y. *et al.* Improvement of the *Oryza sativa* Nipponbare reference genome
650 using next generation sequence and optical map data. *Rice (N Y)* **6**, 4 (2013).
651 42. Sakai, H. *et al.* Rice Annotation Project Database (RAP-DB): an integrative and interactive
652 database for rice genomics. *Plant Cell Physiol* **54**, e6 (2013).
653 43. Xu, X. *et al.* Resequencing 50 accessions of cultivated and wild rice yields markers for
654 identifying agronomically important genes. *Nat Biotechnol* **30**, 105-11 (2011).
655 44. Zhao, Q. *et al.* Pan-genome analysis highlights the extent of genomic variation in
656 cultivated and wild rice. *Nat Genet* **50**, 278-284 (2018).
657 45. Browning, S.R. & Browning, B.L. Rapid and accurate haplotype phasing and missing-data
658 inference for whole-genome association studies by use of localized haplotype clustering.
659 *American Journal of Human Genetics* **81**, 1084-1097 (2007).
660 46. Bolger, A.M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina
661 sequence data. *Bioinformatics* **30**, 2114-20 (2014).
662 47. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler
663 transform. *Bioinformatics* **25**, 1754-60 (2009).
664 48. McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for analyzing
665 next-generation DNA sequencing data. *Genome Res* **20**, 1297-303 (2010).
666 49. Saitou, N. & Nei, M. The neighbor-joining method: a new method for reconstructing
667 phylogenetic trees. *Mol Biol Evol* **4**, 406-25 (1987).
668 50. Pamilo, P. & Nei, M. Relationships between Gene Trees and Species Trees. *Molecular*
669 *Biology and Evolution* **5**, 568-583 (1988).
670 51. Yang, C.C., Sakai, H., Numa, H. & Itoh, T. Gene tree discordance of wild and cultivated
671 Asian rice deciphered by genome-wide sequence comparison. *Gene* **477**, 53-60 (2011).
672 52. Kovach, M.J., Calingacion, M.N., Fitzgerald, M.A. & McCouch, S.R. The origin and
673 evolution of fragrance in rice (*Oryza sativa* L.). *Proc Natl Acad Sci U S A* **106**, 14444-9
674 (2009).
675 53. Zhu, B.F. *et al.* Genetic control of a transition from black to straw-white seed hull in rice
676 domestication. *Plant Physiol* **155**, 1301-11 (2011).
677 54. Du, B. *et al.* Identification and characterization of Bph14, a gene conferring resistance to
678 brown planthopper in rice. *Proc Natl Acad Sci U S A* **106**, 22163-8 (2009).
679 55. Saitoh, K., Onishi, K., Mikami, I., Thidar, K. & Sano, Y. Allelic diversification at the C
680 (Osc1) locus of wild and cultivated rice: Nucleotide changes associated with
681 phenotypes. *Genetics* **168**, 997-1007 (2004).
682 56. Mizuta, Y., Harushima, Y. & Kurata, N. Rice pollen hybrid incompatibility caused by
683 reciprocal gene loss of duplicated genes. *Proc Natl Acad Sci U S A* **107**, 20417-22 (2010).
684 57. Doi, K. *et al.* Ehd1, a B-type response regulator in rice, confers short-day promotion of
685 flowering and controls FT-like gene expression independently of Hd1. *Genes Dev* **18**,
686 926-36 (2004).
687 58. Jin, J. *et al.* GAD1 Encodes a Secreted Peptide That Regulates Grain Number, Grain
688 Length, and Awn Development in Rice Domestication. *Plant Cell* **28**, 2453-2463 (2016).
689 59. Xue, W.Y. *et al.* Natural variation in Ghd7 is an important regulator of heading date and
690 yield potential in rice. *Nature Genetics* **40**, 761-767 (2008).
691 60. Ashikari, M. *et al.* Cytokinin oxidase regulates rice grain production. *Science* **309**, 741-5
692 (2005).
693 61. Fan, C. *et al.* GS3, a major QTL for grain length and weight and minor QTL for grain width
694 and thickness in rice, encodes a putative transmembrane protein. *Theor Appl Genet* **112**,
695 1164-71 (2006).

- 696 62. Song, X.J., Huang, W., Shi, M., Zhu, M.Z. & Lin, H.X. A QTL for rice grain width and weight
697 encodes a previously unknown RING-type E3 ubiquitin ligase. *Nat Genet* **39**, 623-30
698 (2007).
- 699 63. Yano, M. *et al.* Hd1, a major photoperiod sensitivity quantitative trait locus in rice, is
700 closely related to the arabidopsis flowering time gene CONSTANS. *Plant Cell* **12**, 2473-
701 2483 (2000).
- 702 64. Hua, L. *et al.* LABA1, a Domestication Gene Associated with Long, Barbed Awns in Wild
703 Rice. *Plant Cell* **27**, 1875-1888 (2015).
- 704 65. Zhu, Z.F. *et al.* Genetic control of inflorescence architecture during rice domestication.
705 *Nature Communications* **4**(2013).
- 706 66. Yu, Y.C. *et al.* Independent Losses of Function in a Polyphenol Oxidase in Rice:
707 Differentiation in Grain Discoloration between Subspecies and the Role of Positive
708 Selection under Domestication. *Plant Cell* **20**, 2946-2959 (2008).
- 709 67. Tan, L. *et al.* Control of a key transition from prostrate to erect growth in rice
710 domestication. *Nat Genet* **40**, 1360-4 (2008).
- 711 68. Konishi, S. *et al.* An SNP caused loss of seed shattering during rice domestication.
712 *Science* **312**, 1392-6 (2006).
- 713 69. Shomura, A. *et al.* Deletion in a gene associated with grain size increased yields during
714 rice domestication. *Nat Genet* **40**, 1023-8 (2008).
- 715 70. Sweeney, M.T., Thomson, M.J., Pfeil, B.E. & McCouch, S. Caught red-handed: Rc encodes
716 a basic helix-loop-helix protein conditioning red pericarp in rice. *Plant Cell* **18**, 283-94
717 (2006).
- 718 71. Furukawa, T. *et al.* The Rc and Rd genes are involved in proanthocyanidin synthesis in
719 rice pericarp. *Plant J* **49**, 91-102 (2007).
- 720 72. Du, H., Ouyang, Y., Zhang, C. & Zhang, Q. Complex evolution of S5, a major reproductive
721 barrier regulator, in the cultivated rice *Oryza sativa* and its wild relatives. *New Phytol*
722 **191**, 275-87 (2011).
- 723 73. Asano, K. *et al.* Artificial selection for a green revolution gene during japonica rice
724 domestication. *Proc Natl Acad Sci U S A* **108**, 11034-9 (2011).
- 725 74. Li, C., Zhou, A. & Sang, T. Rice domestication by reducing shattering. *Science* **311**, 1936-9
726 (2006).
- 727 75. Takeda, T. *et al.* The OsTB1 gene negatively regulates lateral branching in rice. *Plant J* **33**,
728 513-20 (2003).
- 729 76. Olsen, K.M. *et al.* Selection under domestication: Evidence for a sweep in the rice Waxy
730 genomic region. *Genetics* **173**, 975-983 (2006).

731

732 **Acknowledgements**

733 The research reported in this publication was supported through funding from King
734 Abdullah University of Science and Technology (KAUST), under award numbers
735 BAS/1/1059-01-01 (to T.G.), BAS/1/1606-01-01 (to V.B.B.), FCC/1/1976-03-01 (to T.G.)
736 and FCC/1/1976-20-01 (to T.G.).

737

738 **Author contributions**

739 H.O. designed the study, performed the bioinformatics and statistical analysis, and wrote
 740 the manuscript. K.G. performed the bioinformatics analysis. S.N. wrote the manuscript
 741 and contributed to insightful discussions. R.A.W., M.A.T., K.M. and V.B.B. edited the
 742 manuscript and contributed to insightful discussions. K.L.M. provided easy access to the
 743 genotypes and phenotypes of 3,000 Rice Genomes Project and contributed to insightful
 744 discussions. T.G. designed the study and wrote the manuscript. All the authors discussed
 745 the results and commented on the manuscript.

746

747 **Competing interests**

748 The authors declare no competing interests.

749

750 **Corresponding author**

751 Correspondence to Takashi Gojobori: takashi.gojobori@kaust.edu.sa

752

753 **Figure Legends**

Fig. 1. Passport data of domesticated and wild Asian rice accessions in this study (**a**, in total 4,587 accessions. for more details in higher coverage wilds, see **Supplementary Table 1**), and geographical origin of accessions in 3,000 Rice Genomes Project (**b**, 3,024 accessions). A typical phenotypic diversity within subspecies (**c**, grain length over grain width in *O. sativa* ssp. *indica* (n=1269, green) and *japonica* (n=533, blue)).

Fig. 2. Schematic view of underestimate on genetic *Distance Difference* (**a** and **b**), and phylogenetic analysis of manually curated D-genes (25 genes) and their determined introgressive states (**c** and **d**). Under isolated conditions, each of *indica* and *japonica* subpopulation shall show different genetic distance to the outgroup (a wild rice accession) to some extent, since they have been isolated from each other for a length of time (**a**), whereas they will show unexpectedly similar genetic distance to the outgroup when they made an inter-subspecies crossing (*i.e.* introgression) recently (**b**). Manually curated D-genes (25 genes) and their determined introgressive state (**c**). Reconstructed phylogenetic trees of 25 D-genes (**d**), green nodes : *indica*, blue nodes : *japonica*. Non-introgressive genes were shown in yellow background. Introgressive genes were shown in red background. Genes of undetermined phylogeny were shown in gray background. Phylogenetic trees for one of the D-genes (*LGI*) with variable length of flanking upstream/downstream regions (**e** : CDS only, **f** : +5kb-upstream/+5kb-downstream, **g** : +10kb-upstream/+10kb-downstream, **h** : +20kb-upstream/+20kb-downstream, and **i** : +100kb-upstream/+100kb-downstream, respectively). Full size tree pictures with detailed color system are shown in **Supplementary Fig. 1** and **Supplementary Fig. 2**.

Fig. 3. 100kb- (**a**), 20kb- (**b**), 10kb- (**c**), 5kb- (**d**), and 1kb-resolution (**e**) IR maps (showing chromosome 1 only). The chromosome coordinate was shown in bp on the left

side of horizontal chromosomal rectangles, lined in every 2,500,000 bp. Introgressive windows were shown in red. Non-introgressive windows were shown in yellow. Windows of undetermined phylogeny were shown in gray. Each green rectangle stands for a D-gene region. The 1kb-resolution windows (e) were shown in parallel with SSRs (red lines) and CDRGs (blue lines). Magnified views for two regions in chr01 (f) and chr04 (g) were exemplified as well.

Fig. 4. Numerical distribution of *DD* (Distance Difference). The *DD* statistics according to dimensional continuity of all 1kb windows (a, average of all 12 chromosomes) and the window proportion histogram of particular *DD*s (b, x-axis : *DD* in logarithmic scale, y-axis : frequency of windows). *DD* is defined as below:

$$DD = | F84 (\text{outgroup to } indica) - F84 (\text{outgroup to } japonica) |$$

(*) F84 = Felsenstein 84 nucleotide genetic distance

For more details of the formula, see **Methods**.

Fig. 5. Conceptual diagram of estimated introgression ages. The magnitudes of *DD*s (Distance Differences, red scales) were overdrawn.

Fig. 6. The Sandcastles Model in domestication, a case scenario with three independent introgression events. Each * (asterisk) stands for an agronomically beneficial allele.

a

	3000 Rice Genomes Project		RiceHap3	OryzaGenome	Rice3000+RiceHap3+OryzaGenome	Higer coverage wilds (AA)	Oryza punctata (BB, diploid)	Grand Total
reference	The 3000 rice genomes project 2014		Huang et al. 2012	Ohyanagi et al. 2016	(This study)	Xu et al., 2012	Stein et al. 2018	
	Alexandrov et al. 2015					Ohyanagi et al. 2016		
	Mansueto et al. 2017					Zhao et al. 2018		
	Wang et al. 2018					Stein et al. 2018		
reference genome	Os-Nipponbare-Reference-IRGSP-1.0		IRGSP-build4.0	Os-Nipponbare-Reference-IRGSP-1.0	Os-Nipponbare-Reference-IRGSP-1.0	Os-Nipponbare-Reference-IRGSP-1.0 (This study)	Os-Nipponbare-Reference-IRGSP-1.0 (This study)	
# of accessions		3,024	1,529	446	4,553	35	1	4,587
cultivated		3,024	1,083	-	4,107	-	-	
B#		246 (3KRice 2014 TableS1B)	-	-	246 (3KRice 2014 TableS1B)	-	-	
CX#		312 (3KRice 2014 TableS1B)	-	-	312 (3KRice 2014 TableS1B)	-	-	
IRIS_313-#		2466 (3KRice 2014 TableS1A)	-	-	2466 (3KRice 2014 TableS1A)	-	-	
HP#		-	621 (Huang et al. 2012 TableS7)	-	621 (Huang et al. 2012 TableS7)	-	-	
GP#		-	462 (Huang et al. 2012 TableS7)	-	462 (Huang et al. 2012 TableS7)	-	-	
close-wild (<i>nivara</i> & <i>rufipogon</i>)		-	446 (Huang et al. 2012 TableS2)	446 (Ohyanagi et al. 2016 sup.data)	446 (Ohyanagi et al. 2016 sup.data)	32	-	
distant-wild		-	-	-	-	3	1	
Coverage (against Nipponbare)	High (14x in average)	Low (1x or 2x)	Low (2x)	High + Low (imputed)	High (12x each, at least)	High (140x)		
Is employed in preliminary outgroup assesment?	Yes	No	No	(No)	Yes	Yes		3,060 (Outgroup assesment)
Is employed in main analysis (introgression detection)?	Yes	No	No	(No)	No	Yes		3,025 (Main analysis)
Is employed in SSRs & CLDGRs recomputation?	(Yes)	(Yes)	(Yes)	Yes	No	No		4,553 (SSRs & CLDGRs recomputation)

bioRxiv preprint doi: <https://doi.org/10.1101/829168>; this version posted November 1, 2018. The copyright holder for this preprint (which was not peer-reviewed) is the author/funder. All rights reserved. No reuse allowed without permission.

b

Origin of country	Number of accessions
China	481
India	435
Philippines	229
Bangladesh	186
Thailand	147
Laos	126
Myanmar	75
Malaysia	75
Madagascar	66
Cambodia	59
(Other countries)	374
(Origin unknown)	771

(In total 89 countries)

c

Fig. 1. Passport data of domesticated and wild Asian rice accessions in this study (**a**, in total 4,587 accessions. for more details in higher coverage wilds, see **Supplementary Table 1**), and geographical origin of accessions in 3,000 Rice Genomes Project (**b**, 3,024 accessions). A typical phenotypic diversity within subspecies (**c**, grain length over grain width in *O. sativa* ssp. *indica* (n=1269, green) and *japonica* (n=533, blue)) .

c

GeneSymbol	Description	Reference	LocusID	Location	Introgressive state (by visual inspection)
<i>BADH2</i>	Fragrance	52	Os08g0424500	chr08:20379823..20385975 (+ strand)	Non-introgressive
<i>Bh4</i>	Change hull color	53	Os04g0460200	chr04:22969845..22971859 (+ strand)	Introgressive
<i>Bph14</i>	Brown planthopper resistance	54	Os03g0848700	chr03:35693286..35699010 (- strand)	Non-introgressive
<i>C1</i>	Leaf sheath color and apiculus color	55	Os06g0205100	chr06:5315163..5316640 (+ strand)	Introgressive
<i>DPL2</i>	Hybrid incompatibility	56	Os06g0184100	chr06:4201250..4202851 (- strand)	Non-introgressive
<i>Ehd1</i>	Early heading date	57	Os10g0463400	chr10:17076098..17081344 (- strand)	Non-introgressive
<i>GAD1</i>	Grain number, length and awn development	58	Os08g0485500	chr08:23998787..24000176 (+ strand)	Introgressive
<i>Ghd7</i>	Heading date and yield potential	59	Os07g0261200	chr07:9152377..9155030 (- strand)	Non-introgressive
<i>Gn1a</i>	Grain number	60	Os01g0197700	chr01:5270449..5275585 (- strand)	Non-introgressive
<i>GS3</i>	Increase grain length	61	Os03g0407400	chr03:16729501..16735109 (- strand)	Non-introgressive
<i>GW2</i>	Grain width and weight	62	Os02g0244100	chr02:8115223..8121651 (+ strand)	Non-introgressive
<i>Hd1</i>	Heading date	63	Os06g0275000	chr06:9336376..9338569 (+ strand)	(undetermined)
<i>LABA1</i>	Long and barned awns	64	Os04g0518800	chr04:25959399..25963504 (+ strand)	Introgressive
<i>LG1</i>	Inflorescence architecture	65	Os04g0656500	chr04:33488722..33492700 (+ strand)	Introgressive
<i>Phr1</i>	Change hull color	66	Os03g0329900	chr03:12126320..12131242 (+ strand)	Non-introgressive
<i>Prog1</i>	Tiller erectness	67	Os07g0153600	chr07:2839194..2840089 (- strand)	Introgressive
<i>qSH1</i>	Seed shattering	68	Os01g0848400	chr01:36445456..36449951 (- strand)	Non-introgressive
<i>qSW5</i>	Increase grain width	69	Os05g0187500	chr05:5365122..5366701 (+ strand)	Introgressive
<i>Rc</i>	Change pericarp color	70	Os07g0211500	chr07:6062889..6069304 (+ strand)	Introgressive
<i>Rd</i>	Change pericarp color	71	Os01g0633500	chr01:25382714..25384678 (+ strand)	Non-introgressive
<i>S5</i>	Hybrid sterility	72	Os06g0213100	chr06:5759685..5761518 (+ strand)	(undetermined)
<i>sd1</i>	Reduce tiller length	73	Os01g0883800	chr01:38382385..38385469 (+ strand)	Non-introgressive
<i>sh4</i>	Seed shattering	74	Os04g0670900	chr04:34231186..34233221 (- strand)	Introgressive
<i>tb1</i>	Teosinte branched	75	Os03g0706500	chr03:28428504..28430438 (+ strand)	Non-introgressive
<i>waxy</i>	Amylose content	76	Os06g0133000	chr06:1765622..1770653 (+ strand)	Non-introgressive

Fig. 2. Schematic view of underestimate on genetic *Distance Difference* (**a** and **b**), and phylogenetic analysis of manually curated D-genes (25 genes) and their determined introgressive states (**c** and **d**). Under isolated conditions, each of *indica* and *japonica* subpopulation shall show different genetic distance to the outgroup (a wild rice accession) to some extent, since they have been isolated from each other for a length of time (**a**), whereas they will show unexpectedly similar genetic distance to the outgroup when they made an inter-subspecies crossing (*i.e.* introgression) recently (**b**). Manually curated D-genes (25 genes) and their determined introgressive state (**c**). Reconstructed phylogenetic trees of 25 D-genes (**d**), green nodes : *indica*, blue nodes : *japonica*. Non-introgressive genes were shown in yellow background. Introgressive genes were shown in red background. Genes of undetermined phylogeny were shown in gray background. Phylogenetic trees for one of the D-genes (*LG1*) with variable length of flanking upstream/downstream regions (**e** : CDS only, **f** : +5kb-upstream/+5kb-downstream, **g** : +10kb-upstream/+10kb-downstream, **h** : +20kb-upstream/+20kb-downstream, and **i** : +100kb-upstream/+100kb-downstream, respectively). Full size tree pictures with detailed color system are shown in **Supplementary Fig. 1** and **Supplementary Fig. 2**.

a

all chromosomes

	counts	counts (%)	outgroup to <i>indica</i> (F84 distance)	outgroup to <i>japonica</i> (F84 distance)	<i>DD</i>
overall windows	373,204	100			
phylogeny N.D. windows	133,623	35.8			
phylogeny determined windows	239,581	64.2	0.055106967	0.053881707	1.23E-03
non-introgressive windows	186,567	50.0	0.055653817	0.053942041	1.71E-03
introgressive windows (all)	53,014	14.2	0.05318249	0.05366938	4.87E-04
introgressive windows (narrow = 1)	18,814	5.04	0.052480345	0.053064024	5.84E-04
introgressive windows (wide >= 40)	334	0.0895	0.055056613	0.055050718	5.89E-06

bioRxiv preprint doi: <https://doi.org/10.1101/829168>; this version posted May 1, 2023. The copyright holder for this preprint (which was not peer-reviewed) is the author/funder. All rights reserved. No reuse allowed without permission.

b

Fig. 4. Numerical distribution of *DD* (*Distance Difference*). The *DD* statistics according to dimensional continuity of all 1kb windows (**a**, average of all 12 chromosomes) and the window proportion histogram of particular *DD*s (**b**, x-axis : *DD* in logarithmic scale, y-axis : frequency of windows). *DD* is defined as below:

$$DD = | \text{F84 (outgroup to } indica) - \text{F84 (outgroup to } japonica) |$$

(*) F84 = Felsenstein 84 nucleotide genetic distance

For more details of the formula, see **Methods**.

Non-IRs

bioRxiv preprint doi: <https://doi.org/10.1101/829168>; this version posted May 1, 2018. The copyright holder for this preprint (which was not peer-reviewed) is the author/funder. All rights reserved. No reuse allowed without permission.

Narrow IRs

Wide IRs

Fig. 5. Conceptual diagram of estimated introgression ages. The magnitudes of *DDs* (Distance Differences, red scales) were overdrawn.

past

(Introgression #1)

bioRxiv preprint doi: <https://doi.org/10.1101/829168>; this version posted August 1, 2018. The copyright holder for this preprint (which was not peer-reviewed) is the author/funder. All rights reserved. No reuse allowed without permission.

(Introgression #2)

(Introgression #3)

present

Fig. 6. The Sandcastles Model in domestication, a case scenario with three independent introgression events. Each * (asterisk) stands for an agronomically beneficial allele.