Neotype designation and re-description of Forsskal's reticulate whipray *Himantura uarnak*

Authors: Philippe Borsa, Collin T. Williams, Ashlie J. McIvor, Michael L. Berumen

ORCID: Philippe Borsa: <https://orcid.org/0000-0001-9469-8304>; Collin Taylor Williams: <https://orcid.org/0000-0003-1595-1223>; Ashlie Jane McIvor: <https://orcid.org/0000-0002-2693-216X>; Michael L. Berumen: <https://orcid.org/0000-0003-2463-2742>

P. Borsa

Institut de recherche pour le développement, IRD-UMR 250 "Ecologie marine tropicale des océans Pacifique et Indien", 34394 Montpellier, France

C.T. Williams, A.J. McIvor, M.L. Berumen

Red Sea Research Center, Division of Biological and Environmental Science and Engineering, King Abdullah University of Science and Technology (KAUST), Thuwal 23955-6900, Saudi Arabia

Email : philippe.borsa@ird.fr

Abstract A serious impediment to the taxonomy of the reticulate whipray *Himantura* spp. species complex has been the absence of a type specimen for P. Forsskal's *H. uarnak*. Here, reticulate whipray specimens were sampled from the Jeddah region, the assumed type locality of *H. uarnak*, and characterized genetically at the cytochrome-oxidase subunit 1 (*CO1*) locus. One of these specimens now in the fish collection of the California Academy of Sciences in San Francisco was designated as neotype. A maximum-likelihood phylogeny of all available *CO1* gene sequences from the genus *Himantura* had the following topology: ((*H. leoparda*, *H. uarnak*), (*H. undulata*, (*Himantura* sp. 2, (*H. australis* + *Himantura* sp. 1))), *H. tutul*), where *H. uarnak* haplotypes formed a distinct sub-clade sister to *H. leoparda*. Based on these *CO1* gene sequences, the geographic distribution of *H. uarnak* includes the eastern Mediterranean, the Red Sea, the East African coast, and the Arabian Sea. Two lineages in the reticulate whipray species complex remain to be named.

Keywords Taxonomy · Dasyatidae · Neotype · Redescription · Nucleotide sequence

Notice The present article in portable document (.pdf) format is a published work in the sense of the International Code of Zoological Nomenclature [International Commission on Zoological Nomenclature (ICZN)1999]. It has been registered in ZooBank (<http://zoobank.org/>), the online registration system for the ICZN. The ZooBank life science identifier for this publication is urn:lsid:zoobank.org:pub:B2113697-5EBF-4364-B50C-63019A1A076A. The online version of this work is archived and available from the bioRxiv (<https://biorxiv.org/>) repository.

Introduction

Among the marine fishes listed from P. Forsskål's expedition to the Red Sea were three stingrays, namely the cowtail stingray *Pastinachus sephen* (Fabricius, 1775), the reticulate whipray *Himantura uarnak* (Gmelin, 1789) and the blue-spotted ribbontail ray *Taeniura lymma* (Fabricius, 1775). All three species have long been believed to have a wide Indo-Pacific distribution (e.g., Last & Compagno 1999) but this view has recently been challenged by molecular phylogenetics (Naylor et al. 2012; Arlyza et al. 2013). The type locality for these species is assumed to be Jeddah, the place where P. Forsskål has described most of its fish material from the Red Sea (Fricke 2008). *H. uarnak* has long been understood as the whipray species having densely and regularly spaced, solid, round or oblong dark spots all over the surface of the dorsal side (Rüppell 1835; Duméril 1865; Randall 1983, 1995; Last and Compagno 1999; Manjaji 2004; Arlyza et al. 2013; Manjaji-Matsumoto et al. 2016). *H. uarnak* belongs to a species complex now found to comprise at least six distinct lineages also including *Himantura undulata* Bleeker, 1852, *Himantura leoparda* Manjaji-Matsumoto and Last, 2008, *Himantura australis* Last, White and Naylor, 2016, *Himantura tutul* Borsa, Durand, Shen, Arlyza, Solihin and Berrebi, 2013, previously under *H. leoparda*, and one lineage, '*uarnak* 4' that still has to be named formally (Naylor et al. 2012; Last et al. 2016; Borsa 2017). To date, no material from the Red Sea has been examined genetically. Reticulate whiprays have recently been reported from the eastern Mediterranean Sea (Lessepsian migrants) and identified as either *H. uarnak* based on spot patterns (Ali et al. 2010) or *H. leoparda* based on nucleotide sequences (Yucel et al. 2017). In the absence of type material, it is unclear which of these lineages – or eventually another lineage – is P. Forsskål's *H. uarnak*, emphasizing the need for genetically examining reticulate whipray material from the Red Sea (Naylor et al. 2012). Efforts to stabilize the group's taxonomy by designating a neotype for *H. uarnak* is indeed a task that authors who previously dealt with the taxonomy of species in the genus *Himantura* have neglected (Manjaji 2004; Manjaji-Matsumoto and Last 2008; Borsa et al. 2013; Last et al. 2016; Borsa 2017). The purpose of the present note is to characterize reticulate whipray material collected recently from the Red Sea and designate a neotype for *H. uarnak* in an attempt to clarify the intricate taxonomy of the reticulate whipray species complex.

Materials and methods

Thirteen reticulate whipray individuals were captured on shallow-water sand flats of the eastern shore of the central Red Sea between 2016 and 2019. Eight individuals were captured near the coastal town of Thuwal, Saudi Arabia (22.313°N 39.090°E); five individuals were captured approximately 15 km offshore on Sirrayn Island (19.616°N 40.647°E) in the Farasan Banks region of Saudi Arabia (Fig. 1). Specimen identification numbers and sampling details are provided in Supplementary Table S1. All live specimens were photographed, measured, and a small section of the pelvic fin was excised as genetic material. One individual [no. KAUST-RSRC-H006 (H006)] was retained whole as voucher specimen to be deposited in a museum collection; it was euthanized by immersion in ice-refrigerated seawater; so was another specimen (H009) whose jaws and tissue sample were preserved. The other individuals were released at their

capture site. An additional tissue sample was obtained from an individual on sale at the Jeddah (Saudi Arabia) fish market. The total length, disc width (DW), and length from snout to origin of cloaca were measured on each live specimen. Additional measurements were made on the voucher specimen using the indications of Manjaji (2004). Pigmentation was described using the following parameters: (i) number of spots counted in a rectangular band drawn between spiracles; (ii) number of spots crossed along a line running from mid-scapular point to extremity of pectoral fin, as shown in figure 1 of Borsa et al. (2013); (iii) dorsal-spot diameter; (iv) thickness of paler outer disc margin at extremity of pectoral fin on the dorsal side; and (v) thickness of pigmented outer disc margin at extremity of pectoral fin on the ventral side. Tissue samples ($N = 14$) were placed in ~95% ethanol and stored at -20°C until further processing.

Genomic DNA was extracted using the DNeasy Blood and Tissue Kit (Qiagen, Valencia, CA) following the manufacturer's recommendations. The *FishF1*/*FishR1* primer pair (Ward et al. 2005) was used to target a 655-bp portion of the *COI* gene located between homologous nucleotide sites no. 5571 and no. 6225 of the mitochondrial DNA in *H. leoparda* (NC_028325; Shen et al. 2016) for amplification by polymerase chain reaction (PCR). PCR was run in individual wells loaded with 12 μL reaction mixture composed of 2 μL DNA extract, 5.4 μL Multiplex PCR Master Mix (Qiagen, Valencia, USA), 0.4 μL of each primer (10 μM), and 3.7 μL H_2O . The PCR program included a denaturing step at 95°C for 15 min, followed by 35 cycles of denaturing at 94°C for 30 s, annealing at 50°C for 60 s, and extension at 72°C for 45 s, ending with by a final extension step at 72°C for 10 min. The PCR products were analyzed with the QIAxcel DNA Screening Kit (Qiagen) to confirm successful amplification. Amplified DNA was purified by adding 1.2 μL of ExoProStar (GE Healthcare, Piscataway, USA) to each well before a single thermal cycle of 37°C for 60 min, 85°C for 15 min, and a hold at 4°C until sequencing. PCR samples were then sent for Sanger sequencing at the KAUST Bioscience Core Lab. The sequencing reaction was done in each direction, using either *Fish F1* or *Fish R1* as the sequencing primer. Sanger sequencing of purified target DNA was conducted using the 3730xl DNA Analyzer (Applied Biosystems, Foster City, USA).

Reticulate whipray *COI* gene sequences from the Red Sea were compared to all homologous *Himantura* spp. sequences available including 201 sequences downloaded from the GenBank (www.ncbi.nlm.nih.gov/genbank/; Clark et al. 2016) and BOLD (<http://www.boldsystems.org/>; Ratnasingham and Hebert 2007) public repositories, and two unpublished sequences (Supplementary Table S2). Reticulate whipray, *Himantura* spp. *COI* gene sequences from GenBank and BOLD were labelled "*H. astra*" ($N = 1$), "*H. australis*" ($N = 1$), "*H. fava*" ($N = 1$), "*H. leoparda*" ($N = 85$), "*H. tutul*" ($N = 30$), "*H. uarnak*" ($N = 62$), "*H. uarnak i*" ($N = 2$), "*H. uarnak ii*" ($N = 3$), "*H. uarnak iii*" ($N = 3$), "*H. undulata*" ($N = 11$), "*Himantura* sp." ($N = 1$) and "*Urogymnus asperrimus*" ($N = 1$) (Supplementary Table S2). Partial nucleotide sequences of the *COI* gene were aligned and trimmed to a core length of 613 bp using the BioEdit sequence editing software (Hall 1999). The nucleotide sequences were sorted by species and nucleotide synapomorphies were then assessed visually.

A maximum-likelihood tree of partial *COI* gene sequences was constructed using the MEGA X package (Kumar et al. 2018). All three codon positions were included. Of all models tested under MEGA X, the Hasegawa-Kishino-Yano model (Hasegawa et al. 1985) had the lowest Bayesian information score. The evolutionary history of individual haplotypes was inferred by using the Maximum-Likelihood (ML) method (Felsenstein 1981). A discrete Gamma distribution was used to model evolutionary rate differences among sites ($G = 1.19$). Some sites were allowed to be evolutionarily invariable. GenBank sequences nos. EU398838, EU398867, EU398869, JF493649, KF899470, KF899491, KM073002, KM073006 and KP641389 (*Maculabatis* spp.), EU398849, KF899499, KM072993, KM073009 and KM073010 (*Pateobatis* spp.), EU398860, FJ384709, KF899521, KF899528, KM072995 and KR003772 (*Brevitrygon* spp.) and KF965292

were included as an outgroup to root the tree. Mean nucleotide distances between and within lineages were estimated using the Maximum Composite Likelihood model under MEGA X.

Results

Disc width ranged between 284 mm and 748 mm. The claspers of the one of the five male specimens (HC256) were calcified. Reticulate whipray specimen no. H006 is presented in Fig. 2a-d. The colour of the dorsal side of all individuals sampled was beige and the ventral side was creamy white. The dorsal side was covered with numerous brown spots, which generally were evenly distributed and had a circular, or slightly elongated, or dumbbell shape. In a proportion of individuals, circular spots were often associated as pairs. Colour tones and spot patterns on the dorsal side were similar to those encountered in the reticulate whipray's mangrove habitat in the central Red Sea (Fig. 2e). The ventral side was either immaculate or with few dark-grey spots, apart from the outer margin of the posterior half of the disk which had a pale greyish hue with densely distributed small, dark-grey spots. A summary of the measurements used to describe spot patterns on all twelve specimens is reported in Table 1. Dorsal-spot diameter was correlated with disk width (Pearson's $r = 0.66$; $P < 0.05$) and so was dorsal-spot density estimated from transects ($r = 0.82$, $r = 0.56$; Fisher's combined $P < 0.01$), indicating that both the size and number of dorsal spots increase with age.

The ML tree of *Himantura* spp. sequences (Fig. 3) showed four main clades (clades I-IV; Arlyza et al. 2013) with strong statistical support. Clade I comprised haplotypes sampled from off eastern Africa, including the holotype of *H. tutul*, to the Sulu Sea. Clade-I haplotypes were previously identified as either "*H. leoparda*" (Lim et al. 2015; Bineesh et al. 2017; Mohd-Arshaad & Jamaludin 2018), "*H. tutul*" (Arlyza et al. 2013; Borsa et al. 2013), "*H. uarnak*", "*H. uarnak i*", "*H. uarnak ii*", "*H. uarnak iii*" (Bineesh et al. 2017; Gouws 2020; Hanner 2020), "*Himantura* sp." (Hanner 2020), or "*Urogymnus asperrimus*" (Priyanga et al. 2013). Clade II comprised haplotypes sampled from the Arabian Sea to the Sulu Sea, all identified as either "*H. fava*" (Ward et al. 2005) or "*H. undulata*" (Arlyza et al. 2013; Lim et al. 2015; Bineesh et al. 2017; Mohd-Arshaad & Jamaludin 2018; Segura-Garcia & Yain Tun 2018; Ahmed et al. 2019; present work). Three subclades or haplogroups were distinguished within clade III. Subclade III₁ included two sequences sampled from the Sulu Sea so far labelled "*H. uarnak*" and "*H. uarnak ii*", respectively (Lim et al. 2015; Hanner 2020) while subclade III₂ included haplotypes sampled from tropical Australia exclusively and so far referred to as either "*H. astra*" (Cerutti-Pereyra et al. 2012), "*H. australis*" (Appleyard 2020), "*H. leoparda*" (Cerutti-Pereyra et al. 2012), or "*H. uarnak*" (Cerutti-Pereyra et al. 2012 ; Appleyard 2020 ; McGrouther 2020). Haplogroup III₃, which in the ML tree was paraphyletic with III₂, exclusively comprised haplotypes sampled from the Coral Triangle and labelled either "*H. uarnak*" (Arlyza et al. 2013; Santos et al. 2014; Lim et al. 2015; Mohd Arshaad & Jamaludin 2018; Appleyard 2020; Hanner 2020; Mohd Arshaad 2020) or "*H. uarnak i*" (Hanner 2020). All *Himantura* spp. CO1 gene sequences sampled from the Red Sea clustered into a distinctive lineage within clade IV, hereafter designated subclade IV₂, also including haplotypes from the eastern Mediterranean Sea, Natal (South Africa) and the Arabian Sea. These were initially assigned to either "*H. leoparda*" (Bineesh et al. 2017; Yucel et al. 2017) or "*H. uarnak*" (Steinke et al. 2011). Subclade IV₁ which was sister to IV₂ comprised haplotypes from India, Sunda, Western Australia and Taiwan. These were initially identified as either "*H. leoparda*" (Arlyza et al. 2013; Shen et al. 2016; Appleyard 2017; Ravi et al. 2019) or "*H. undulata*" (Appleyard 2020). Sub-clade IV₂ specific to the Red Sea and adjacent localities was separated from sub-clade IV₁ by 1.8% nucleotide distance at the CO1 locus. The mean nucleotide distance was 0.3% within sub-clade IV₁ and 0.4% within sub-clade IV₂.

Discussion

The taxonomic value of mitochondrial DNA sequences has been demonstrated in morphologically intractable species complexes in Elasmobranchs such as those of the long-tailed butterfly ray *Gymnura poecilura* (Naylor et al. 2012 ; Muktha et al. 2018), the whitespotted whiptail *Maculabatis gerrardi* (Ward et al. 2008; Naylor et al. 2012) or the blue-spotted maskray *Neotrygon kuhlii* (Naylor et al. 2012; Puckridge et al. 2013; Borsa et al. 2018; Pavan-Kumar et al. 2018). The reticulate whiptail *Himantura* spp. species complex is another example where morphological overlap between species has led to considerable confusion (Naylor et al. 2012; Borsa 2017), as confirmed by the plethora of names assigned to mitochondrial DNA haplotypes that cluster within the same clade or subclade (present survey).

In total, seven distinct lineages or haplogroups were identified in the phylogenetic tree of the reticulate whiptail species complex (Fig. 3). Subclade *IV*₂ included the *CO1* gene sequence of a specimen here chosen as *H. uarnak*'s neotype (see next section). As a consequence, sub-clade *IV*₂ is here designated as P. Forsskål's *H. uarnak*. The *H. uarnak* subclade was distinct from the subclade *IV*₁ clustering all leopard whiptail *H. leoparda* specimens from the Coral Triangle. *Himantura leoparda*, whose type locality is the Gulf of Carpentaria in the Coral Triangle region, thus is the sibling Pacific species of *H. uarnak*. The other lineages were identified as *H. australis* (sub-clade *III*₂), *H. tutul* (clade *I*) and *H. undulata* (clade *II*) based on previous genetic work employing the *CO1* marker (Arlyza et al. 2013; Borsa et al. 2013; Borsa 2017; Appleyard 2020) and on geographic consistency with the type locality of *H. australis* (southern New Guinea; Last et al. 2016). Two sub-clades or haplogroups here labelled *Himantura* sp. 1 (haplogroup *III*₂) and *Himantura* sp. 2 (subclade *III*₁) remain to be described as new species or, possibly re-described as a resurrected species. Thus, the fixation of a neotype for *H. uarnak* clarifies the identity of over 200 specimens whose *CO1* gene sequences have been deposited in public databases and paves the way to the description or re-description of two other yet-unnamed species in the genus *Himantura*.

Despite the brevity of its description, P. Forsskål's *H. uarnak* has been universally treated as a single, valid taxon in the subsequent literature (Gmelin 1789 ; Rüppell 1835; Bleeker 1852 ; Duméril 1865; Last and Compagno 1999; Manjaji 2004). However, no type is known; and multiple lineages that qualify as distinct species have been assigned to *H. uarnak* (Naylor et al. 2012; Last et al. 2016; Borsa 2017) demonstrating taxonomic confusion. For clarifying past and future research dealing with the reticulate whiptail species complex we elected to designate a neotype that unambiguously identifies *H. uarnak*.

Neotype designation and re-description of *Himantura uarnak*

Genus *Himantura* Müller & Henle, 1837; species *H. uarnak* [Gmelin (ex-Forsskål), 1789], type species of the genus. The authorship of the species has been discussed extensively by Fricke (2008).

As argued above, designating a neotype for *Himantura uarnak* is necessary to clarify its taxonomic status. This act also helps secure the stability of the nomenclature of other species in the reticulate whiptail species complex. Forsskål's (1775) original description mentioned a whiptail ("cauda, quae apterygia") with spots all over ("tota maculata"). Reticulate whiptail specimen KAUST-RSRC-H006 (male, 369 mm DW, collected on April 25, 2019 by AJM and CTW) has the foregoing attributes and it was captured off the Arabian shore of the central Red Sea ca. 80 km north of Jeddah, the assumed type locality (Fricke 2008). This specimen, which was allocated collection no. CAS-ICH 247241 in the fish collections of the California Academy of Sciences, San Francisco is here designated as *Himantura uarnak*'s neotype. This designation thus satisfies the

conditions expressed in the International Code of Zoological Nomenclature (International Commission on Zoological Nomenclature 1999: Article no. 75).

Neotype locality is Thuwal, Saudi Arabia (22°18'40"N 39°05'26"E); habitat is shallow sand flat ~50 cm deep adjacent to mangroves dominated by *Avicennia marina*.

Morphological description: disc rhomboidal, length 94.6% DW; snout with distinct apical lobe; anterior margins of disc convex, lateral apices narrowly rounded; posterior margin convex, free rear tip rounded. Pelvic fins moderately elongate, length 17.4% DW; width across base 14.5% DW. Tail whip-like, tapering gently toward sting, length 230.8% DW; see also Table 1, and Fig. 2a-d. Additional morphological measurements taken on neotype after it was fixed in formalin were the following (in mm): disc width 351, total length 1089, snout to pectoral-fin insertion 299, end of orbit to pectoral insertion 214, snout to maximum width 135, snout to origin of cloaca 284, cloaca origin to sting 129, pectoral insertion to sting origin 138, disc thickness 48, snout to preorbital (direct) 77, snout to preorbital (horizontal) 68, orbit diameter 16, eye diameter 19, spiracle length 14, orbit+spiracle length 36, interorbital width 51, inter-eye width 77, distance between spiracles 82, head length (direct) 165, preoral length (to lower jaw) 77, snout (prenasal) 58, nostril length 20, nasal curtain length 23, nasal curtain width 41, distance between nostrils 34, mouth width 38, distance between 1st gill slits 74, distance between 5th gill slits 51, width of 1st gill slit 10, width of 3rd gill slit 10, width of 5th gill slit 7, tail width at axil of pelvic fins 21, tail width at base of sting 11, tail height at axil of pelvic fins 17, tail height base of caudal sting 8, sting length 79, cloaca length 2, greatest width across pelvic fins 88 (resting) or 102 (spread), clasper length 31 (post-cloaca) or 13 (from pelvic axil).

Genetic description: *CO1* gene sequences cluster into a distinct lineage in the phylogenetic tree of the genus *Himantura* (Fig. 3). Nucleotide sequence of partial *CO1* gene of neotype, comprised between homologous nucleotide sites no. 69 and no. 699 of the *CO1* gene in *H. leoparda*

(GenBank accession no. NC_028325; Shen et al. 2016) is 5'- C G G T G C G T G A G C A C G G A T A G T G G G T A C T G G C C T T A G C C T G C T T A T T C G G A C A G A G C T A A G C C A A C C A G G C G C A T T A C T G G G T G A T G A T C A A A A A T A T A A T G T A A T T G T T A C C G C C C A T G C C T T C G T A A T A A T C T T T T T C A T G G T A A T A C C T A T T A T A A T T G G G G G C T T T G G T A A T T G A C T C G T C C C C C T A A T A A T C G G T G C T C C A G A T A T A G C C T T T C C T C G A A T A A A C A A C A T G A G T T T T T G A C T T C T T C C A C C A T C C T T T C T A C T A C T T T T G G C C T C T G C T G G A G T A G A G G C T G G C G C T G G A A C A G G C T G A A C A G T C T A T C C C C C A C T A G C T G G T A A T C T A G C A C A T G C A G G G G C T T C A G T A G A C T T A G C A A T C T T T T C C C T A C A C C T G G C C G G T G T A T C T T C T A T C C T A G C C T C T A T T A A T T T T A T C A C C A C A A T C A T T A A C A T A A A A C C A C C A G C A A T T T C G C A G T A T C A A A C A C C C C T C T T T G T C T G A T C A A T C C T T A T C A C A G C C G T A C T C C T C T T G T T A T C T C T T C C T G T C C T A G C A G C A G G T A T T A C G A T A C T T C T A A C A G A T C G T A A C C T C A A T A C A A C C T T C T T T G A T C C T G C A G G A G G A G G T G A C C C A A T T C T T T A T C A A -3'.

Four nucleotides here underlined characterize *H. uarnak*.

Geographic distribution: *H. uarnak* specimens whose identity was here ascertained from their partial *CO1* gene sequences (Supplementary Table S2) were previously reported as "*H. leoparda*" (Arlyza et al. 2013; Borsa et al. 2013; Yucel et al. 2017; Bineesh et al. 2018) or "*H. uarnak*" (Steinke et al. 2011). Conversely, the identity of specimens previously assigned to "*H. uarnak*" are now correctly assigned to *H. australis*, *H. tutul*, and two yet-unnamed *Himantura* spp. (see Supplementary Table S2). Based on these genetically validated records, the distribution of *H. uarnak* outside the Red Sea includes the eastern Mediterranean Sea (Yucel et al. 2017), Natal (Steinke et al. 2011), and the Arabian Sea (Bineesh et al. 2017).

Acknowledgements Foremost, thanks are due to M. Tietbohl for donating tissue samples, assisting in field collections, and providing laboratory assistance.. Additionally, We are grateful to J. Spaet and members of KAUST's Reef Ecology Lab for their support at various stages of this project; to D. Catania and L.A. Rocha for allocating a CAS collection number to the neotype of *H. uarnak*; to X. Chen for sharing reticulate whipray sequences; to M. Bennett-Smith for sharing excellent photographs; and to S. Bogorodsky, H. Debelius and R. Fricke for insightful discussions about the taxonomy and nomenclature of reticulate whiprays.

Funding information Support to this study was provided by King Abdullah University of Science and Technology (KAUST) through baseline funding to MLB and Institut de recherche pour le développement, Marseille to PB..

Conflict of interest The authors declare that they have no conflict of interest.

Ethical approval Reticulate whipray specimens were captured in accordance with KAUST's institutional animal care and use committee (IACUC) under animal study proposal # 18IACUC14.

Data availability All data generated during this study are included in this published article or in the appended Supplementary Tables S1 and S2. Sequences produced through the present study have been deposited in GenBank (<https://www.ncbi.nlm.nih.gov/nucleotide/>) under accession nos. XX000000-XX000000. Photographs of all specimens are available from the authors upon request.

Author contributions AJM and CTW participated in the design of the study, collected and measured specimens, did the molecular analyses and participated in the writing. MLB contributed funding and laboratory facilities and supervised the study. PB designed the study, provided directions, compiled and analyzed the data and led the writing. All authors read, edited and approved the final version of the manuscript.

References

- Ahmed M.S., Zhilik A.A., Chowdhury N.Z., Ahmed S. (2019) DNA barcoding of marine fishes of Bangladesh. <https://www.ncbi.nlm.nih.gov/nuccore/MN013427>
- Ali M, Saad A, Ben Amor MM, Capapé C (2010) First records of the honeycomb stingray, *Himantura uarnak* (Forskål, 1775), off the Syrian coast (eastern Mediterranean) (Chondrichthyes: Dasyatidae). Zool Middle East 49:104-106
- Appleyard SA (2017) http://www.boldsystems.org/index.php/Public_RecordView?processid=FOAO1113-18,FOAO1135-18,FOAO1142-18
- Appleyard SA (2020) http://www.boldsystems.org/index.php/Public_RecordView?processid=FOAG126-07,FOAJ878-09,FOAO1107-18
- Arlyza IS, Shen K-N, Solihin DD, Soedharma D, Berrebi P, Borsa P (2013) Species boundaries in the *Himantura uarnak* species complex (Myliobatiformes: Dasyatidae). Mol Phyl Evol 66:429-435
- Bineesh KK, Gopalakrishnan A, Akhilesh KV, Sajeela KA, Abdussamad EM, Pillai NGK, Basheer VS, Jena JK, Ward RD (2017) DNA barcoding reveals species composition of sharks and rays in the Indian commercial fishery. Mitoch DNA Pt A 28:458-472
- Bleeker P (1852) Bijdrage tot de kennis der Plagiostomen van den Indischen archipel. Verhandel Batav Genoot Kunst Wetensch 24:1-92, pls 1-4
- Borsa P (2017) Comments on “Annotated checklist of the living sharks, batoids and chimaeras (Chondrichthyes) of the world, with a focus on biogeographical diversity” (Weigmann, 2016). J Fish Biol 90:1170-1175

- Borsa P, Durand J-D, Shen K-N, Arlyza IS, Solihin DD, Berrebi P (2013) *Himantura tutul* sp. nov. (Myliobatoidei: Dasyatidae), a new ocellated whipray from the tropical Indo-West Pacific, described from its cytochrome-oxidase I gene sequence. C R Biol 336:82-92
- Cerutti-Pereyra F, Meekan MG, Wei N-WV, O'Shea O, Bradshaw CJA, Austin CM (2012) Identification of rays through DNA barcoding: an application for ecologists. PLoS One 7:e36479
- Clark K, Karsch-Mizrachi I, Lipman DJ, Ostell J, Sayers EW (2016) GenBank. Nucl Acids Res 44:D67–D72
- Duméril A (1865) Histoire naturelle des poissons ou ichthyologie générale. Tome premier Elasmobranches. Librairie encyclopédique du Roret, Paris, 720 p
- Felsenstein J (1981) Evolutionary trees from DNA sequences: A maximum likelihood approach. J Mol Evol 17:368–376
- Felsenstein J (1985) Confidence limits on phylogenies: an approach using the bootstrap. Evolution 39:783–791
- Forsskål P (1775) Descriptiones animalium, avium, amphibiorum, piscium, insectorum, vermium; quae in itinere orientali observavit Petrus Forsskål (post mortem auctoris edidit Carsten Niebuhr). Adjuncta est materia medica kahirina atque tabula maris Rubri geographica. Mölleri, Hauniae, xxxiv+164 p
- Fricke R (2008) Authorship, availability and validity of fish names described by Peter (Pehr) Simon Forsskål and Johann Christian Fabricius in the '*Descriptiones animalium*' by Carsten Niebuhr in 1775 (Pisces). Stuttg Beitr Naturk A Neue Ser 1:1–76
- Gmelin JF (1789) Caroli a Linné [...] Systema Naturae per regna tria naturae, secundum classes, ordines, genera, species; cum characteribus, differentiis, synonymis, locis. Editio decimo tertia, aucta, reformata, vol 1, pars III. G.E. Beer, Lipsiae, pp 1033–1516
- Gouws G (2020) [http://www.boldsystems.org/index.php/Public_RecordView?processid=SAIAB712-08, SAIAB713-08, SAIAB714-08, SAIAB715-08](http://www.boldsystems.org/index.php/Public_RecordView?processid=SAIAB712-08,SAIAB713-08,SAIAB714-08,SAIAB715-08)
- Hall TA (1999) BIOEDIT: A user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. Nucl Acids Symp Ser 41:95-98
- Hanner R (2020) [http://www.boldsystems.org/index.php/Public_RecordView?processid=IRREK171-08, IRREK193-08, IRREK662-08, IRREK678-08, IRREK679-08, IRREK687-08, IRREK741-08, IRREK752-08, IRREK829-08](http://www.boldsystems.org/index.php/Public_RecordView?processid=IRREK171-08,IRREK193-08,IRREK662-08,IRREK678-08,IRREK679-08,IRREK687-08,IRREK741-08,IRREK752-08,IRREK829-08)
- Hasegawa M., Kishino H., Yano T. (1985) Dating the human-ape split by a molecular clock of mitochondrial DNA. J Mol Evol 22:160-174.
- International Commission on Zoological Nomenclature (1999) International code of zoological nomenclature, 4th edn. International Trust for Zoological Nomenclature, London, 306 p
- Kumar S, Stecher G, Li M, Knyaz C, Tamura K (2018) MEGA X: Molecular evolutionary genetics analysis across computing platforms. Mol Biol Evol 35:1547-1549.
- Last PR, Compagno LJV (1999) Dasyatidae - Stingrays. In Carpenter KE, Niem VH (eds) FAO species identification guide for fishery purposes. The living marine resources of the Western Central Pacific. Volume 3. Batoid fishes, chimaeras and bony fishes part 1 (Elopidae to Linophrynidae). FAO, Rome, 1479-1505.
- Last PR, White WT, Naylor G (2016) Three new stingrays (Myliobatiformes: Dasyatidae) from the Indo-West Pacific. Zootaxa 4147:377-402
- Lim KC, Lim PE, Chong VC, Loh KH (2015) Molecular and morphological analyses reveal phylogenetic relationships of stingrays focusing on the family Dasyatidae (Myliobatiformes). PLoS One 10:e0120518

- Manjaji BM (2004) Taxonomy and phylogenetic systematics of the Indo-Pacific whip-tailed stingray genus *Himantura* Müller and Henle 1837 (Chondrichthyes: Myliobatiformes: Dasyatidae). PhD Thesis, University of Tasmania, xxii+607 p
- Manjaji-Matsumoto BM, Last PR (2008) *Himantura leoparda* sp. nov., a new whipray (Myliobatoidei: Dasyatidae) from the Indo-Pacific. CSIRO Mar Atmosph Res Pap 22:292-301
- Manjaji-Matsumoto BM, White WT, Gutteridge AN (2016) *Himantura uarnak*. IUCN Red List Threat Sp 2016:e.T161692A68629130
- McGrouther M.A. (2020)
http://www.boldsystems.org/index.php/Public_RecordView?processid=AMS045-06
- Mohd Arshaad W (2020)
[http://www.boldsystems.org/index.php/Public_RecordView?processid=JTFR099-17, JTFR112-17](http://www.boldsystems.org/index.php/Public_RecordView?processid=JTFR099-17,JTFR112-17)
- Mohd Arshaad W, Jamaludin N-A (2018) DNA barcoding of rays in Malaysia.
<https://www.ncbi.nlm.nih.gov/nuccore/MG774902>, MG774903, MG774913, MG774915, MG774922, MG792078, MG792101, MG792110, MG792112, MG792113, MG792123, MG792124, MG792125
- Muktha M, Akhilesh KV, Sandhya S, Jasmin F, Jishnudev MA, Kizhakudan SJ (2018) Re-description of the longtail butterfly ray, *Gymnura poecilura* (Shaw, 1804) (Gymnuridae: Myliobatiformes) from Bay of Bengal with a neotype designation. Mar Biodiv 48:1085-1096
- Naylor GJP, Caira JN, Jensen K, Rosana KAM, White WT, Last PR (2012) A DNA sequence-based approach to the identification of shark and ray species and its implications for global elasmobranch diversity and parasitology. Bull Am Mus Nat Hist 367:1-262
- Pavan-Kumar A, Kumar R, Pitale P, Shen K-N, Borsa P (2018) *Neotrygon indica* sp. nov., the Indian-Ocean blue spotted maskray (Myliobatoidei, Dasyatidae). C R Biol 341:120-130
- Priyanga K, Thangaraj M, Singh R (2013) Molecular identification of *Urogymnus asperrimus* (family: Dasyatidae) using cytochrome oxidase I (COI) as a molecular marker.
<https://www.ncbi.nlm.nih.gov/nuccore/KC508509>
- Puckridge M, Last PR, White WT, Andreakis N (2013) Phylogeography of the Indo-West Pacific maskrays (Dasyatidae, *Neotrygon*): A complex example of chondrichthyan radiation in the Cenozoic. Ecol Evol 3:217-232
- Randall JE (1983) Red Sea reef fishes. IMMEL Publishing, London, 192 p
- Randall JE (1995) Coastal fishes of Oman. Crawford, Bathurst, xii+439 p
- Ratnasingham S, Hebert PD (2007) BOLD: the barcode of life data system (<http://www.barcodinglife.org>). Mol Ecol Notes 7:55–364
- Ravi RK, Venu S, Bineesh KK, Akhilesh KV, Sajeela KA, Basheer VS, Gopalakrishnan A (2019)
<https://www.ncbi.nlm.nih.gov/nuccore/MK422130>
- Rüppell E (1835) Neue Wirbelthiere zu der Fauna von Abyssinien gehörig. Fische des rothen Meeres. Siegmund Schmerber, Frankfurt am Mein, ii+148 p, 33 pl
- Santos MD, Asis AMM, Lacsamana JKM (2014) Illegal trade of regulated and protected aquatic species in the Philippines detected by DNA barcoding.
<https://www.ncbi.nlm.nih.gov/nuccore/KC970332>
- Segura-Garcia I, Yain Tun T (2018) Unscrambling Myanmar marine fishery biodiversity using DNA barcoding. <https://www.ncbi.nlm.nih.gov/nuccore/MH235646>, MH235647
- Shen K-N, Chang C-W, Tsai S-Y, Wu S-C, Lin Z-H, Chan Y-F, Chen C-H, Hsiao C-D, Borsa P (2016) Next generation sequencing yields the complete mitogenomes of leopard whipray (*Himantura leoparda*) and blue-spotted maskray (*Neotrygon kuhlii*) (Chondrichthyes: Dasyatidae). Mitoch DNA Pt A 27:2613-2614
- Steinke D, Zemlak TS, Connell AD, Heemstra PC, Hebert PDN (2011)
<https://www.ncbi.nlm.nih.gov/nuccore/JF493651>, JF493652

- Ward RD, Holmes BH, White WT, Last PR (2008) DNA barcoding Australasian chondrichthyans: results and potential uses in conservation. *Mar Freshw Res* 59:57–71
- Ward RD, Zemlak TS, Innes BH, Last PR, Hebert PDN (2005) DNA barcoding Australia's fish species. *Phil Trans Roy Soc Lond B* 360:1847–1857
- Yucel N, Sakalli A, Karahan A (2017) First record of the honeycomb stingray *Himantura leoparda* (Manjaji-Matsumoto & Last, 2008) (Myliobatoidei: Dasyatidae) in the Mediterranean Sea, confirmed by DNA barcoding. *J Appl Ichthyol* 33:530–532

Table 1 *Himantura uarnak*. Pigmentation patterns in 12 DNA-barcoded specimens sampled from the Jeddah region. *DW* disk width; *na* no data

Specimen no.	DW (mm)	Dorsal side				Ventral side			
		Spot density between spiracles ^b	Spot density on pectoral fin ^c		Spot diameter (mm) ^d	Outer margin (% DW) ^e		Outer margin (% DW) ^f	
			Left	Right		Left	Right	Left	Right
H001	319	35	14	18	3.1 ± 0.4	15.4	14.9	na	na
H002	346	13	7	13	4.0 ± 0.6	18.7	20.0	na	na
H003	284	2	4	7	4.6 ± 0.5	25.8	25.3	na	na
H004	326	19	7	9	3.3 ± 0.3	11.8	9.8	na	na
H005	317	19	5	5	3.2 ± 0.6	29.1	25.0	na	na
H006 ^a	369	33	13	12	4.9 ± 0.6	2.9	3.1	31.3	25.6
H009	531	28	11	9	5.5 ± 1.1	9.1	6.8	na	na
HC127	278	15	8	11	3.3 ± 0.5	11.3	11.3	na	na
HC131	310	15	10	5	5.2 ± 0.6	9.6	14.1	na	na
HC150	288	21	11	12	4.0 ± 0.6	7.0	8.1	na	na
HC256	748	34	26	25	5.2 ± 0.7	4.1	5.6	12.5	na
HC289	580	16	15	9	6.6 ± 1.2	2.3	2.4	na	na

^a Neotype

^b Number of spots counted in band between spiracles

^c Number of spots crossed along line running from mid-scapular point to extremity of pectoral fin (see figure 1 of Borsa et al. 2013)

^d Average ± SD, from 30-36 randomly-chosen dorsal spots in central area of pectoral fin

^e Thickness of paler outer disc margin at extremity of pectoral fin

^f Thickness of pigmented outer disc margin at extremity of pectoral fin

Fig. 1 Reticulate whipray sampling localities in the central Red Sea. Jeddah is assumed to be the type locality of Forsskal's (1775) *H. uarnak* (Fricke 2008)

Fig. 2 Reticulate whipray *Himantura uarnak* (Gmelin [ex Forsskål], 1789). **a** Dorsal view of neotype, no. CAS-ICH 247241 deposited at the fish collections of the California Academy of Sciences, San Francisco (formerly at the Red Sea Research Center, Thuwal as no. KAUST-RSRC-H006); scale bar: 5 cm. **b** Neotype, ventral view; scale bar: 5 cm. **c** Neotype, spine details in median dorsal region; scale bar: 1cm. **d** Neotype, mouth details; scale bar: 1cm. **e** Sandy mangrove habitat of *H. uarnak* off Thuwal, Red Sea, 22.313°N 39.090°E (credit: M. Bennett-Smith / KAUST)

Fig. 3 Reticulate whipray *Himantura* spp. species complex. Maximum-likelihood tree of *CO1* gene haplotypes based on the HKY+G+I model (MEGA X; Kumar et al. 2018). Tree with highest log likelihood (-16139.6) is shown. Tree is drawn to scale, with branch lengths measured in the number of substitutions per site. Haplotype sequences are listed using their GenBank or BOLD accession number. Scores at a node are percentages of trees in which the associated taxa clustered together (from 500 iterations of bootstrap resampling; Felsenstein 1985). The four main clades are designated by roman numbers *I-IV* following Arlyza et al. (2013); numbers in subscript indicate a sub-clade or an haplogroup within a clade. Arrow indicates placement of *H. uarnak* neotype

Supplementary Table S1 Sampling details for reticulate whipray *Himantura uarnak* specimens from the Red Sea including specimen no. KAUST-RSRC-H006 / CAS-ICH 247241 chosen as neotype (highlighted red)

Supplementary Table S2 Reticulate whipray *Himantura uarnak* species complex. List of nucleotide sequences at the *CO1* locus used for the present work, with GenBank and BOLD identification nos. and other details including sampling locations and references. Sequences were retrieved on 23 March 2020 from the GenBank and BOLD public databases (*N* = 201) or donated by X. Chen (Wenzhou Medical College, China) (*N* = 2) or produced through the present survey (*N* = 13). Sequences are arranged by species and presented in alphabetical order

Figure 1

Supplementary Table S1 Sampling details for reticulate whipray *Himantura uarnak* specimens from the Red Sea including specimen no. KAUST-RSRC-H006 / CAS-ICH 247241 chosen as neotype (highlighted red)

Specimen ID	Location	Date	Approximate depth	Collector	Method of capture	Sex	No. stings	Total length (cm)	Disk width (cm)	Snout-to-cloaca length (cm)
H001	Sirrayn south	14 Feb. 2019	15 cm	AJM, CTW	15-m seine net	F	1	113.2	31.9	23.8
H002	Sirrayn south	14 Feb. 2019	15 cm	AJM, CTW	15-m seine net	F	1	121.8	34.6	26.5
H003	Sirrayn south	14 Feb. 2019	15 cm	AJM, CTW	15-m seine net	M	na	103.2	28.4	22.8
H004	Sirrayn south	14 Feb. 2019	15 cm	AJM, CTW	15-m seine net	F	1	120.2	32.6	25.0
H005	Sirrayn south	15 Feb. 2019	20 cm	AJM, CTW	15-m seine net	M	1	100.2	31.7	24.7
H006	KAUST north beach	25 Apr. 2019	50 cm	AJM, CTW	15-m seine net	M	1	109.9	36.9	28.9
H007	Jeddah fish market	29 Apr. 2019	na	AJM, CTW	Chunk of pectoral fin purchased	na	na	na	na	na
H008	KAUST north beach	22 May 2019	20 cm	AJM, CTW	Handline baited with shrimp	M	1	103.4	27.2	22.1
H009	KAUST harbor	Dec. 2016	na	M. Tietbohl	15-m seine net	F	2	na	53.1	na
HC127	KAUST pier	01 Feb. 2019	39 cm	AJM, CTW	15-m seine net	F	na	102.0	27.8	29.0
HC131	KAUST pier	23 Mar. 2019	30 cm	AJM, CTW	Dip net	F	2	113.4	31.0	24.6
HC150	KAUST north beach	30 Apr. 2019	10 cm	AJM, CTW	15-m seine net	F	1	96.7	28.8	23.2
HC256	KAUST north beach	25 Apr. 2019	80 cm	AJM, CTW	15-m seine net	M	1	197.0	74.8	54.8
HC289	KAUST pier	01 Feb. 2019	74 cm	AJM, CTW	15-m seine net	F	2	184.0	58.0	48.0

Supplementary Table S2 Reticulate whipray *Himantura uarnak* species complex. List of nucleotide sequences at the CO1 locus used for the present work, with GenBank and BOLD identification nos. and other details including sampling locations and references. Sequences were retrieved on 23 March 2020 from the GenBank and BOLD public databases ($N = 201$) or donated by X. Chen (Wenzhou Medical College, China) ($N = 2$) or produced through the present survey ($N = 13$). Sequences are arranged by species and presented in alphabetical order ($N = 2$) or produced through the present survey ($N = 13$). Sequences are arranged by species and presented in alphabetical order. Type material highlighted red

GenBank no.	BOLD no.	Other identifier	Initial identification	Sampling locality	Reference	BIN	Species*
<i>H. australis</i>							
JQ765509	GBGC11930-13	FCP-HUNR08	<i>H. astra</i>	Skeleton Bay, Coral Bay, -23.12 113.77	Cerutti-Pereyra et al. 2012	BOLD:ABZ0063	<i>H. australis</i>
JQ765519	GBGC11920-13	FCP-HUNR22	<i>H. leoparda</i>	Ranger Bay, Ningaloo, 21°54'S 113°59'E	Cerutti-Pereyra et al. 2012	BOLD:ABZ0063	<i>H. australis</i>
JQ765520	GBGC11919-13	FCP-HUNR23	<i>H. leoparda</i>	Ranger Bay, Ningaloo, 21°54'S 113°59'E	Cerutti-Pereyra et al. 2012	BOLD:ABZ0063	<i>H. australis</i>
JQ765521	GBGC11918-13	FCP-HUNR25	<i>H. uarnak</i>	Skeleton Bay, Coral Bay, -23.12 113.77	Cerutti-Pereyra et al. 2012	BOLD:ABZ0063	<i>H. australis</i>
JQ765522	GBGC11917-13	FCP-HUNR26	<i>H. uarnak</i>	Mangrove Bay, Ningaloo Marine Park, -21.97 113.93	Cerutti-Pereyra et al. 2012	BOLD:ABZ0063	<i>H. australis</i>
JQ765523	GBGC11916-13	FCP-HUNR27	<i>H. uarnak</i>	Skeleton Bay, Coral Bay, -23.12 113.77	Cerutti-Pereyra et al. 2012	BOLD:ABZ0063	<i>H. australis</i>
JQ765524	GBGC11915-13	FCP-HUNR28	<i>H. uarnak</i>	Skeleton Bay, Coral Bay, -23.12 113.77	Cerutti-Pereyra et al. 2012	BOLD:ABZ0063	<i>H. australis</i>
JQ765525	GBGC11914-13	FCP-HUNR29	<i>H. uarnak</i>	Point Cloates, Ningaloo, 22°42'S 113°40'E	Cerutti-Pereyra et al. 2012	BOLD:ABZ0063	<i>H. australis</i>
JQ765526	GBGC11913-13	FCP-HUNR30	<i>H. uarnak</i>	Point Cloates, Ningaloo, 22°42'S 113°40'E	Cerutti-Pereyra et al. 2012	BOLD:ABZ0063	<i>H. australis</i>
JQ765527	GBGC11912-13	FCP-HUNR31	<i>H. uarnak</i>	Stanley's Pool, Ningaloo, 22°59'S 113°48'E	Cerutti-Pereyra et al. 2012	BOLD:ABZ0063	<i>H. australis</i>
JQ765528	GBGC11911-13	FCP-HUNR32	<i>H. uarnak</i>	Mangrove Bay, Ningaloo Marine Park, -21.97 113.93	Cerutti-Pereyra et al. 2012	BOLD:ABZ0063	<i>H. australis</i>
JQ765529	GBGC11910-13	FCP-HUNT34	<i>H. uarnak</i>	Shoal Bay, NT, 12°21'S 131°01'E	Cerutti-Pereyra et al. 2012	BOLD:ABZ0063	<i>H. australis</i>
JQ765530	GBGC11909-13	FCP-HUNT35	<i>H. uarnak</i>	Shoal Bay, NT, 12°21'S 131°01'E	Cerutti-Pereyra et al. 2012	BOLD:ABZ0063	<i>H. australis</i>
JQ815394	GBGC11830-13	isolate 24	<i>H. uarnak</i>	Skeleton Bay, Coral Bay, -23.12 113.77	Cerutti-Pereyra et al. 2012	BOLD:ABZ0063	<i>H. australis</i>
JQ815395	GBGC11829-13	isolate 36	<i>H. uarnak</i>	Meckets Creek, NT, 12°20'S 130°57'E	Cerutti-Pereyra et al. 2012	BOLD:ABZ0063	<i>H. australis</i>
JQ929047	GBGC11827-13	isolate 21	<i>H. leoparda</i>	Skeleton Bay, Coral Bay, -23.12 113.77	Cerutti-Pereyra et al. 2012	BOLD:ABZ0063	<i>H. australis</i>
-	AMS045-06	AMS I.39533-003	<i>H. uarnak</i>	Cairns area, Queensland, -16.92 145.77	McGrouther 2020	BOLD:ABZ0063	<i>H. australis</i>
-	FOAI353-09	BW-A6280, GT 4012 #2	<i>H. uarnak</i>	Skeleton Bay, Coral Bay, -23.12 113.77	Appleyard 2020	BOLD:ABZ0063	<i>H. australis</i>
-	FOAI354-09	BW-A6281, GT 4013 #21	<i>H. uarnak</i>	Skeleton Bay, Coral Bay, -23.12 113.77	Appleyard 2020	BOLD:ABZ0063	<i>H. australis</i>
-	FOAI355-09	BW-A6282, GT 4014 #55	<i>H. uarnak</i>	Skeleton Bay, Coral Bay, -23.12 113.77	Appleyard 2020	BOLD:ABZ0063	<i>H. australis</i>
-	FOAI356-09	BW-A6283, GT 4015 #60	<i>H. uarnak</i>	Skeleton Bay, Coral Bay, -23.12 113.77	Appleyard 2020	BOLD:ABZ0063	<i>H. australis</i>
-	FOAI358-09	BW-A6285, GT 4017 #40	<i>H. uarnak</i>	Mangrove Bay, Ningaloo Marine Park, -21.97 113.93	Appleyard 2020	BOLD:ABZ0063	<i>H. australis</i>
-	FOAO1107-18	BW-A14305, GT 10444	<i>H. australis</i>	Western Australia, -19.35 118.12	Appleyard 2020	BOLD:ABZ0063	<i>H. australis</i>
<i>H. leoparda</i>							
JX263361	GBGC12896-13	IR003	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830	<i>H. leoparda</i>
JX263362	GBGC12895-13	IR004	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830	<i>H. leoparda</i>
JX263363	GBGC12894-13	IR008	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830	<i>H. leoparda</i>
JX263364	GBGC12893-13	IR009	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830	<i>H. leoparda</i>
JX263365	GBGC12892-13	IR010	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830	<i>H. leoparda</i>
JX263366	GBGC12891-13	IR011	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830	<i>H. leoparda</i>
JX263367	GBGC12890-13	IR012	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830	<i>H. leoparda</i>
JX263368	GBGC12889-13	IR013	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830	<i>H. leoparda</i>
JX263369	GBGC12888-13	IR014	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830	<i>H. leoparda</i>

JX263370	GBGC12887-13	IR015	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263371	GBGC12886-13	IR016	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263372	GBGC12885-13	IR017	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263373	GBGC12884-13	IR018	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263374	GBGC12883-13	IR019	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263375	GBGC12882-13	IR020	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263376	GBGC12881-13	IR021	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263377	GBGC12880-13	IR022	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263378	GBGC12879-13	IR023	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263379	GBGC12878-13	IR025	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263380	GBGC12877-13	IR027	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263381	GBGC12876-13	IR028	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263382	GBGC12875-13	IR030	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263383	GBGC12874-13	IR031	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263384	GBGC12873-13	IR032	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263385	GBGC12872-13	IR033	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263386	GBGC12871-13	IR034	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263387	GBGC12870-13	IR035	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263388	GBGC12869-13	IR036	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263389	GBGC12868-13	IR039	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263390	GBGC12867-13	IR040	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263391	GBGC12866-13	IR041	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263392	GBGC12865-13	IR042	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263393	GBGC12864-13	IR046	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263394	GBGC12863-13	IR048	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263395	GBGC12862-13	IR050	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263396	GBGC12861-13	IR051	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263397	GBGC12860-13	IR053	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263398	GBGC12859-13	IR054	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263399	GBGC12858-13	IR057	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263400	GBGC12857-13	IR058	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263401	GBGC12856-13	IR059	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263402	GBGC12855-13	IR060	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263403	GBGC12854-13	IR061	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263404	GBGC12853-13	IR062	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263405	GBGC12852-13	IR063	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263406	GBGC12851-13	IR064	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263407	GBGC12850-13	IR065	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263408	GBGC12849-13	IR066	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263409	GBGC12848-13	IR067	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>

JX263410	GBGC12847-13	IR069	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263411	GBGC12846-13	IR070	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263412	GBGC12845-13	IR071	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263413	GBGC12844-13	IR072	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263414	GBGC12843-13	IR074	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263415	GBGC12842-13	IR075	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263416	GBGC12841-13	IR077	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263417	GBGC12840-13	IR078	<i>H. leoparda</i>	Batang	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
JX263418	GBGC12839-13	IR087	<i>H. leoparda</i>	Sunda	Arlyza et al. 2013	BOLD:AAB7830 <i>H. leoparda</i>
MK422130	-	CR03J	<i>H. leoparda</i>	India	Ravi et al. 2019	- <i>H. leoparda</i>
NC_028325	-	wjc329	<i>H. leoparda</i>	Taiwan	Shen et al. 2016	- <i>H. leoparda</i>
-	FOAG126-07	BW-A4003, TWG-58	<i>H. undulata</i>	Tashi fish market, Taiwan	Appleyard 2020	BOLD:AAB7830 <i>H. leoparda</i>
-	FOAO1113-18	BW-A14311	<i>H. leoparda</i>	Western Australia, -19.605 117.292	Appleyard 2017	BOLD:AAB7830 <i>H. leoparda</i>
-	FOAO1135-18	BW-A14335	<i>H. leoparda</i>	Western Australia, -19.724 116.499	Appleyard 2017	BOLD:AAB7830 <i>H. leoparda</i>
-	FOAO1142-18	BW-A14342	<i>H. leoparda</i>	Western Australia, -19.968 116.031	Appleyard 2017	BOLD:AAB7830 <i>H. leoparda</i>
<i>Himantura</i> sp. 1						
JX263337	GBGC12920-13	IR089	<i>H. uarnak</i>	Makassar	Arlyza et al. 2013	BOLD:AAB7832 <i>Himantura</i> sp. 1
JX263338	GBGC12919-13	IR090	<i>H. uarnak</i>	Makassar	Arlyza et al. 2013	BOLD:AAB7832 <i>Himantura</i> sp. 1
JX263339	GBGC12918-13	IR091	<i>H. uarnak</i>	Makassar	Arlyza et al. 2013	BOLD:AAB7832 <i>Himantura</i> sp. 1
JX263340	GBGC12917-13	IR092	<i>H. uarnak</i>	Makassar	Arlyza et al. 2013	BOLD:AAB7832 <i>Himantura</i> sp. 1
JX263341	GBGC12916-13	IR093	<i>H. uarnak</i>	Makassar	Arlyza et al. 2013	BOLD:AAB7832 <i>Himantura</i> sp. 1
JX263342	GBGC12915-13	IR094	<i>H. uarnak</i>	Makassar	Arlyza et al. 2013	BOLD:AAB7832 <i>Himantura</i> sp. 1
JX263343	GBGC12914-13	IR095	<i>H. uarnak</i>	Makassar	Arlyza et al. 2013	BOLD:AAB7832 <i>Himantura</i> sp. 1
JX263344	GBGC12913-13	IR096	<i>H. uarnak</i>	Makassar	Arlyza et al. 2013	BOLD:AAB7832 <i>Himantura</i> sp. 1
JX263345	GBGC12912-13	IR097	<i>H. uarnak</i>	Makassar	Arlyza et al. 2013	BOLD:AAB7832 <i>Himantura</i> sp. 1
JX263346	GBGC12911-13	IR098	<i>H. uarnak</i>	Makassar	Arlyza et al. 2013	BOLD:AAB7832 <i>Himantura</i> sp. 1
JX263347	GBGC12910-13	IR099	<i>H. uarnak</i>	Makassar	Arlyza et al. 2013	BOLD:AAB7832 <i>Himantura</i> sp. 1
JX263348	GBGC12909-13	IR100	<i>H. uarnak</i>	Selayar	Arlyza et al. 2013	BOLD:AAB7832 <i>Himantura</i> sp. 1
JX263349	GBGC12908-13	IR101	<i>H. uarnak</i>	Selayar	Arlyza et al. 2013	BOLD:AAB7832 <i>Himantura</i> sp. 1
JX263350	GBGC12907-13	MZB-20875, LIPI 4419, IR102	<i>H. uarnak</i>	Selayar	Arlyza et al. 2013	BOLD:AAB7832 <i>Himantura</i> sp. 1
JX263351	GBGC12906-13	IR103	<i>H. uarnak</i>	Selayar	Arlyza et al. 2013	BOLD:AAB7832 <i>Himantura</i> sp. 1
JX263352	GBGC12905-13	IR104	<i>H. uarnak</i>	Selayar	Arlyza et al. 2013	BOLD:AAB7832 <i>Himantura</i> sp. 1
JX263353	GBGC12904-13	IR105	<i>H. uarnak</i>	Selayar	Arlyza et al. 2013	BOLD:AAB7832 <i>Himantura</i> sp. 1
JX263354	GBGC12903-13	IR106	<i>H. uarnak</i>	Selayar	Arlyza et al. 2013	BOLD:AAB7832 <i>Himantura</i> sp. 1
JX263355	GBGC12902-13	IR107	<i>H. uarnak</i>	Kendari	Arlyza et al. 2013	BOLD:AAB7832 <i>Himantura</i> sp. 1
JX263356	GBGC12901-13	IR108	<i>H. uarnak</i>	Kendari	Arlyza et al. 2013	BOLD:AAB7832 <i>Himantura</i> sp. 1
JX263357	GBGC12900-13	IR109	<i>H. uarnak</i>	Kendari	Arlyza et al. 2013	BOLD:AAB7832 <i>Himantura</i> sp. 1
JX263358	GBGC12899-13	IR110	<i>H. uarnak</i>	Labuan Bajo	Arlyza et al. 2013	BOLD:AAB7832 <i>Himantura</i> sp. 1
JX263359	GBGC12898-13	IR111	<i>H. uarnak</i>	Labuan Bajo	Arlyza et al. 2013	BOLD:AAB7832 <i>Himantura</i> sp. 1
JX263360	GBGC12897-13	wjc637	<i>H. uarnak</i>	Taiwan	Arlyza et al. 2013	BOLD:AAB7832 <i>Himantura</i> sp. 1

KC970332	BFPHL067-13	DFBOC I	<i>H. uarnak</i>	Philippines	Santos et al. 2014	BOLD:AAB7832 <i>Himantura</i> sp. 1
KM072999	ANGBF12762-15	HUAR3	<i>H. uarnak</i>	Semporna, Malaysia	Lim et al. 2015	BOLD:AAB7832 <i>Himantura</i> sp. 1
MG792110	ANGBF48003-19	-	<i>H. uarnak</i>	Sandakan, Sabah, 5.84N 118.12E	Mohd Arshaad & Jamaludin 2018	BOLD:AAB7832 <i>Himantura</i> sp. 1
MG792123	ANGBF48002-19	-	<i>H. uarnak</i>	Sandakan, Sabah, 5.84N 118.12E	Mohd Arshaad & Jamaludin 2018	BOLD:AAB7832 <i>Himantura</i> sp. 1
-	FOAJ878-09	BW-A7728, IN01396	<i>H. uarnak</i>	Tanjung Luar, Lombok, 8.75S 116.58E	Appleyard 2020	BOLD:AAB7832 <i>Himantura</i> sp. 1
-	IRREK193-08	GN2238, JPAG 036	<i>H. uarnak</i>	Bacolod City, Philippines	Hanner 2020	BOLD:AAB7832 <i>Himantura</i> sp. 1
-	IRREK863-08	GN3119, BO-82	<i>H. uarnak</i> i	Sarawak, Malaysia	Hanner 2020	BOLD:AAB7832 <i>Himantura</i> sp. 1
-	JTFR099-17	PSK11	<i>H. uarnak</i>	Sandakan, Sabah, 5.84N 118.12E	Mohd Arshaad 2020	BOLD:AAB7832 <i>Himantura</i> sp. 1
-	JTFR112-17	PSK26	<i>H. uarnak</i>	Sandakan, Sabah, 5.84N 118.12E	Mohd Arshaad 2020	BOLD:AAB7832 <i>Himantura</i> sp. 1
<i>Himantura</i> sp. 2						
KM073000	ANGBF12763-15	HUAR 1	<i>H. uarnak</i>	Sandakan, Sabah	Lim et al. 2015	BOLD:ACV9539 <i>Himantura</i> sp. 2
-	IRREK752-08	GN3006, HBO-109	<i>H. uarnak</i> ii	Lahad Datu, Sabah	Hanner 2020	BOLD:ACV9539 <i>Himantura</i> sp. 2
<i>H. tutul</i>						
JX263306	GBGC12950-13	IR001	<i>H. tutul</i>	Batang	Arlyza et al. 2013; Borsa et al. 2013 BOLD:AAB7831	<i>H. tutul</i>
JX263307	GBGC12949-13	IR002	<i>H. tutul</i>	Batang	Arlyza et al. 2013; Borsa et al. 2013 BOLD:AAB7831	<i>H. tutul</i>
JX263308	GBGC12948-13	IR005	<i>H. tutul</i>	Batang	Arlyza et al. 2013; Borsa et al. 2013 BOLD:AAB7831	<i>H. tutul</i>
JX263309	GBGC12947-13	IR006	<i>H. tutul</i>	Batang	Arlyza et al. 2013; Borsa et al. 2013 BOLD:AAB7831	<i>H. tutul</i>
JX263310	GBGC12946-13	IR024	<i>H. tutul</i>	Batang	Arlyza et al. 2013; Borsa et al. 2013 BOLD:AAB7831	<i>H. tutul</i>
JX263311	GBGC12945-13	IR026	<i>H. tutul</i>	Batang	Arlyza et al. 2013; Borsa et al. 2013 BOLD:AAB7831	<i>H. tutul</i>
JX263312	GBGC12944-13	IR029	<i>H. tutul</i>	Batang	Arlyza et al. 2013; Borsa et al. 2013 BOLD:AAB7831	<i>H. tutul</i>
JX263313	GBGC12943-13	IR037	<i>H. tutul</i>	Batang	Arlyza et al. 2013; Borsa et al. 2013 BOLD:AAB7831	<i>H. tutul</i>
JX263314	GBGC12942-13	IR043	<i>H. tutul</i>	Batang	Arlyza et al. 2013; Borsa et al. 2013 BOLD:AAB7831	<i>H. tutul</i>
JX263315	GBGC12941-13	IR044	<i>H. tutul</i>	Batang	Arlyza et al. 2013; Borsa et al. 2013 BOLD:AAB7831	<i>H. tutul</i>
JX263316	GBGC12940-13	IR045	<i>H. tutul</i>	Batang	Arlyza et al. 2013; Borsa et al. 2013 BOLD:AAB7831	<i>H. tutul</i>
JX263317	GBGC12939-13	IR047	<i>H. tutul</i>	Batang	Arlyza et al. 2013; Borsa et al. 2013 BOLD:AAB7831	<i>H. tutul</i>
JX263318	GBGC12938-13	IR049	<i>H. tutul</i>	Batang	Arlyza et al. 2013; Borsa et al. 2013 BOLD:AAB7831	<i>H. tutul</i>
JX263319	GBGC12937-13	IR052	<i>H. tutul</i>	Batang	Arlyza et al. 2013; Borsa et al. 2013 BOLD:AAB7831	<i>H. tutul</i>
JX263320	GBGC12936-13	IR055	<i>H. tutul</i>	Batang	Arlyza et al. 2013; Borsa et al. 2013 BOLD:AAB7831	<i>H. tutul</i>
JX263321	GBGC12935-13	IR056	<i>H. tutul</i>	Batang	Arlyza et al. 2013; Borsa et al. 2013 BOLD:AAB7831	<i>H. tutul</i>
JX263322	GBGC12934-13	IR073	<i>H. tutul</i>	Batang	Arlyza et al. 2013; Borsa et al. 2013 BOLD:AAB7831	<i>H. tutul</i>
JX263323	GBGC12933-13	IR076	<i>H. tutul</i>	Batang	Arlyza et al. 2013; Borsa et al. 2013 BOLD:AAB7831	<i>H. tutul</i>
JX263324	GBGC12932-13	IR079	<i>H. tutul</i>	Batang	Arlyza et al. 2013; Borsa et al. 2013 BOLD:AAB7831	<i>H. tutul</i>
JX263325	GBGC12931-13	IR080	<i>H. tutul</i>	Batang	Arlyza et al. 2013; Borsa et al. 2013 BOLD:AAB7831	<i>H. tutul</i>
JX263326	GBGC12930-13	IR081	<i>H. tutul</i>	Batang	Arlyza et al. 2013; Borsa et al. 2013 BOLD:AAB7831	<i>H. tutul</i>
JX263327	GBGC12929-13	IR082	<i>H. tutul</i>	Batang	Arlyza et al. 2013; Borsa et al. 2013 BOLD:AAB7831	<i>H. tutul</i>
JX263328	GBGC12928-13	IR083	<i>H. tutul</i>	Batang	Arlyza et al. 2013; Borsa et al. 2013 BOLD:AAB7831	<i>H. tutul</i>
JX263329	GBGC12927-13	IR084	<i>H. tutul</i>	Batang	Arlyza et al. 2013; Borsa et al. 2013 BOLD:AAB7831	<i>H. tutul</i>
JX263330	GBGC12926-13	IR085	<i>H. tutul</i>	Batang	Arlyza et al. 2013; Borsa et al. 2013 BOLD:AAB7831	<i>H. tutul</i>
JX263331	GBGC12925-13	IR086	<i>H. tutul</i>	Sunda	Arlyza et al. 2013; Borsa et al. 2013 BOLD:AAB7831	<i>H. tutul</i>
JX263332	GBGC12924-13	IR088	<i>H. tutul</i>	Sunda	Arlyza et al. 2013; Borsa et al. 2013 BOLD:AAB7831	<i>H. tutul</i>

JX263333	GBGC12923-13	IR112	<i>H. tutul</i>	Bali	Arlyza et al. 2013; Borsa et al. 2013 BOLD:AAB7831	<i>H. tutul</i>
JX263334	GBGC12922-13	IR113	<i>H. tutul</i>	Southern central Java	Arlyza et al. 2013; Borsa et al. 2013 BOLD:AAB7831	<i>H. tutul</i>
JX263335	GBGC12832-13	MNHN:ICTI5184 (HOLOTYPE)	<i>H. tutul</i>	Mkoani, Tanzania	Arlyza et al. 2013; Borsa et al. 2013 BOLD:AAB7831	<i>H. tutul</i>
KC508509	GBGC11390-13	CASMBAUTRL16	<i>Urogymnus asperrimus</i>	-	Priyanga et al. 2013	BOLD:AAB7831 <i>H. tutul</i>
KF899353	ANGBF10976-15	NBFGR:CHN:HL49	<i>H. leoparda</i>	Mumbai, India	Bineesh et al. 2017	BOLD:AAB7831 <i>H. tutul</i>
KF899507	ANGBF11561-15	NBFGR:CHN:K42	<i>H. uarnak</i>	Kochi, India	Bineesh et al. 2017	BOLD:AAB7831 <i>H. tutul</i>
KF899508	ANGBF11560-15	NBFGR:CHN:K43	<i>H. uarnak</i>	Kochi, India	Bineesh et al. 2017	BOLD:AAB7831 <i>H. tutul</i>
KF899509	ANGBF11559-15	NBFGR:CHN:204	<i>H. uarnak</i>	Kochi, India	Bineesh et al. 2017	BOLD:AAB7831 <i>H. tutul</i>
KF899510	ANGBF11558-15	NBFGR:CHN:K44	<i>H. uarnak</i>	Kochi, India	Bineesh et al. 2017	BOLD:AAB7831 <i>H. tutul</i>
KF899511	ANGBF11557-15	NBFGR:CHN:BM42	<i>H. uarnak</i>	Kochi, India	Bineesh et al. 2017	BOLD:AAB7831 <i>H. tutul</i>
KM072996	ANGBF12752-15	HLEO2	<i>H. leoparda</i>	Sandakan, Malaysia	Lim et al. 2015	BOLD:AAB7831 <i>H. tutul</i>
KM072997	ANGBF12753-15	HLEO1	<i>H. leoparda</i>	Sandakan, Malaysia	Lim et al. 2015	BOLD:AAB7831 <i>H. tutul</i>
KM072998	ANGBF12754-15	HLEO3	<i>H. leoparda</i>	Sandakan, Malaysia	Lim et al. 2015	BOLD:AAB7831 <i>H. tutul</i>
MG774902	JTFR010-17	PDGN04	<i>H. leoparda</i>	Dungun, Malaysia, 4.77 N 103.25 E	Mohd-Arshaad & Jamaludin 2018	BOLD:AAB7831 <i>H. tutul</i>
MG774903	JTFR011-17	PDGN05	<i>H. leoparda</i>	Dungun, Malaysia, 4.77 N 103.25 E	Mohd-Arshaad & Jamaludin 2018	BOLD:AAB7831 <i>H. tutul</i>
MG774913	JTFR021-17	PKTN102	<i>H. leoparda</i>	Kuantan, Malaysia, 3.47 N 103.19 E	Mohd-Arshaad & Jamaludin 2018	BOLD:AAB7831 <i>H. tutul</i>
MG774915	JTFR023-17	PKTN18	<i>H. leoparda</i>	Kuantan, Malaysia, 3.47 N 103.19 E	Mohd-Arshaad & Jamaludin 2018	BOLD:AAB7831 <i>H. tutul</i>
MG774922	JTFR030-17	PKTN35	<i>H. leoparda</i>	Kuantan, Malaysia, 3.47 N 103.19 E	Mohd-Arshaad & Jamaludin 2018	BOLD:AAB7831 <i>H. tutul</i>
MG792078	ANGBF47998-19, JTFR056-17	PMKH19	<i>H. leoparda</i>	Malaysia, 2.54 N 112.05 E	Mohd-Arshaad & Jamaludin 2018	BOLD:AAB7831 <i>H. tutul</i>
MG792101	ANGBF47999-19, JTFR090-17	PSK01	<i>H. leoparda</i>	Sandakan, Malaysia, 5.84 N 118.12 E	Mohd-Arshaad & Jamaludin 2018	BOLD:AAB7831 <i>H. tutul</i>
MG792112	ANGBF47994-19, JTFR101-17	PSK14	<i>H. leoparda</i>	Sandakan, Malaysia, 5.84 N 118.12 E	Mohd-Arshaad & Jamaludin 2018	BOLD:AAB7831 <i>H. tutul</i>
MG792113	ANGBF47995-19, JTFR102-17	PSK15	<i>H. leoparda</i>	Sandakan, Malaysia, 5.84 N 118.12 E	Mohd-Arshaad & Jamaludin 2018	BOLD:AAB7831 <i>H. tutul</i>
MG792124	ANGBF47996-19, JTFR113-17	PSK27	<i>H. leoparda</i>	Sandakan, Malaysia, 5.84 N 118.12 E	Mohd-Arshaad & Jamaludin 2018	BOLD:AAB7831 <i>H. tutul</i>
MG792125	ANGBF47997-19, JTFR114-17	PSK28	<i>H. leoparda</i>	Sandakan, Malaysia, 5.84 N 118.12 E	Mohd-Arshaad & Jamaludin 2018	BOLD:AAB7831 <i>H. tutul</i>
-	IRREK171-08	GN2207	<i>Himantura</i> sp.	Singapore	Hanner 2020	BOLD:AAB7831 <i>H. tutul</i>
-	IRREK662-08	GN2916, BOD-49	<i>H. uarnak</i> iii	Malaysia	Hanner 2020	BOLD:AAB7831 <i>H. tutul</i>
-	IRREK678-08	GN2932, BOD-65	<i>H. uarnak</i> ii	Malaysia	Hanner 2020	BOLD:AAB7831 <i>H. tutul</i>
-	IRREK679-08	GN2933, BOD-66	<i>H. uarnak</i> iii	Malaysia	Hanner 2020	BOLD:AAB7831 <i>H. tutul</i>
-	IRREK687-08	GN2941, BOD-74	<i>H. uarnak</i> iii	Malaysia	Hanner 2020	BOLD:AAB7831 <i>H. tutul</i>
-	IRREK741-08	GN2995, HBO-86	<i>H. uarnak</i> ii	Sandakan, Sabah, Malaysia	Hanner 2020	BOLD:AAB7831 <i>H. tutul</i>
-	IRREK829-08	GN3085, BO-47	<i>H. uarnak</i> i	Sarawak, Malaysia	Hanner 2020	BOLD:AAB7831 <i>H. tutul</i>
-	SAIAB712-08	80368-1, LS07-0449	<i>H. uarnak</i>	Niule Reef, Tanga, 5.08S 39.10 E	Gouws 2020	BOLD:AAB7831 <i>H. tutul</i>
-	SAIAB713-08	80368-2, LS07-0930	<i>H. uarnak</i>	Niule Reef, Tanga, 5.08S 39.10 E	Gouws 2020	BOLD:AAB7831 <i>H. tutul</i>
-	SAIAB714-08	80368-3, LS07-0934	<i>H. uarnak</i>	Niule Reef, Tanga, 5.08S 39.10 E	Gouws 2020	BOLD:AAB7831 <i>H. tutul</i>
-	SAIAB715-08	80368-4, LS07-0750	<i>H. uarnak</i>	Niule Reef, Tanga, 5.08S 39.10 E	Gouws 2020	BOLD:AAB7831 <i>H. tutul</i>
-	-	ZB01210334(W3)5440F	<i>H. tutul</i>	Thailand	X. Chen (pers. comm., 2015)	- <i>H. tutul</i>
<i>H. uarnak</i>						
JF493651	TZMSC474-05	Smith 30.10-2	<i>H. uarnak</i>	Cape Vidal 28.13 S 32.56 E	Steinke et al. 2011	BOLD:AAB7830 <i>H. uarnak</i>
JF493652	TZMSC232-05	ADC30.10-1	<i>H. uarnak</i>	Cape Vidal 28.13 S 32.56 E	Steinke et al. 2011	BOLD:AAB7830 <i>H. uarnak</i>
KF899500	ANGBF10975-15	NBFGR:CHN:PK60	<i>H. leoparda</i>	Mumbai, India	Bineesh et al. 2017	BOLD:AAB7830 <i>H. uarnak</i>

KF899501	ANGBF10974-15	NBFGR:CHN:PK61	<i>H. leoparda</i>	Mumbai, India	Bineesh et al. 2017	BOLD:AAB7830 <i>H. uarnak</i>
KF899502	ANGBF10973-15	NBFGR:CHN:PK62	<i>H. leoparda</i>	Mumbai, India	Bineesh et al. 2017	BOLD:AAB7830 <i>H. uarnak</i>
-	FEE001-16	IMSMETU-HL-01	<i>H. leoparda</i>	off Hatay, Turkey 36°11'N 35°48'E	Yucel et al. 2017	BOLD:AAB7830 <i>H. uarnak</i>
-	-	H002	-	Sirrayn South	Present survey	- <i>H. uarnak</i>
-	-	H003	-	Sirrayn South	Present survey	- <i>H. uarnak</i>
-	-	H004	-	Sirrayn South	Present survey	- <i>H. uarnak</i>
-	-	H005	-	Sirrayn North	Present survey	- <i>H. uarnak</i>
-	-	CAS-ICH 247241 (NEOTYPE)	<i>H. uarnak</i>	KAUST North Beach, Thuwal, Saudi Arabia	Present survey	- <i>H. uarnak</i>
-	-	H007	-	Jeddah Fish Market	Present survey	- <i>H. uarnak</i>
-	-	H008	-	KAUST North Beach	Present survey	- <i>H. uarnak</i>
-	-	H009	-	KAUST Harbor	Present survey	- <i>H. uarnak</i>
-	-	HC127	-	KAUST Pier	Present survey	- <i>H. uarnak</i>
-	-	HC131	-	KAUST Pier	Present survey	- <i>H. uarnak</i>
-	-	HC150	-	KAUST North Beach	Present survey	- <i>H. uarnak</i>
-	-	HC256	-	KAUST North Beach	Present survey	- <i>H. uarnak</i>
-	-	HC289	-	KAUST Pier	Present survey	- <i>H. uarnak</i>
<i>H. undulata</i>						
DQ108167	FOA221-04	BW-A221	<i>H. fava</i>	Sandakan, Malaysia	Ward et al. 2005	BOLD:AAF0692 <i>H. undulata</i>
JX263336	GBGC12921-13	IR007	<i>H. undulata</i>	Batang fish landing place, Java Sea	Arlyza et al. 2013	BOLD:AAF0692 <i>H. undulata</i>
KF899503	ANGBF11565-15	NBFGR:CHN:BM3	<i>H. undulata</i>	Mumbai, India	Bineesh et al. 2017	BOLD:AAF0692 <i>H. undulata</i>
KF899504	ANGBF11564-15	NBFGR:CHN:BM4	<i>H. undulata</i>	Mumbai, India	Bineesh et al. 2017	BOLD:AAF0692 <i>H. undulata</i>
KF899505	ANGBF11563-15	NBFGR:CHN:BM48	<i>H. undulata</i>	Mumbai, India	Bineesh et al. 2017	BOLD:AAF0692 <i>H. undulata</i>
KF899506	ANGBF11562-15	NBFGR:CHN:BM49	<i>H. undulata</i>	Mumbai, India	Bineesh et al. 2017	BOLD:AAF0692 <i>H. undulata</i>
KM073001	ANGBF12764-15	HUND1	<i>H. undulata</i>	Tawau, Malaysia	Lim et al. 2015	BOLD:AAF0692 <i>H. undulata</i>
MG792127	ANGBF48004-19	JTFR116-17, PSK31	<i>H. undulata</i>	Sandakan, Malaysia, 5.84 N 118.12 E	Mohd-Arshaad & Jamaludin 2018	BOLD:AAF0692 <i>H. undulata</i>
MH235646	-	USNM:444112	<i>H. undulata</i>	Tanintharyi, Myanmar, 12.43N 98.60E	Segura-Garcia & Yain Tun 2018	- <i>H. undulata</i>
MH235647	-	ISGDG1	<i>H. undulata</i>	Tanintharyi, Myanmar, 12.43N 98.60E	Segura-Garcia & Yain Tun 2018	- <i>H. undulata</i>
MN013427	GBGC15127-19	DUZM_MF_036B	<i>H. undulata</i>	Cox's Bazar, Bangladesh, 21.28N 91.25E	Ahmed et al. 2019	BOLD:AAF0692 <i>H. undulata</i>
-	-	ZB01210333(W2)5440F	<i>H. undulata</i>	Thailand	X. Chen (pers. comm., 2015)	- <i>H. undulata</i>

* identification of individual to species based on haplotype placement in CO1 gene phylogeny