



ELSEVIER

Salt resistant crop plants[☆]

Stuart J Roy¹, Sónia Negrão^{2,3} and Mark Tester³

Soil salinity is a major constraint to agriculture. To improve salinity tolerance of crops, various traits can be incorporated, including ion exclusion, osmotic tolerance and tissue tolerance. We review the roles of a range of genes involved in salt tolerance traits. Different tissues and cells are adapted for specific and often diverse function, so it is important to express the genes in specific cell-types and to pyramid a range of traits. Modern biotechnology (marker-assisted selection or genetic engineering) needs to be increasingly used to introduce the correct combination of genes into elite crop cultivars. Importantly, the effects of introduced genes need to be evaluated in the field to determine their effect on salinity tolerance and yield improvement.

Addresses

¹ Australian Centre for Plant Functional Genomics, School of Agriculture,

Food and Wine, University of Adelaide, Urrbrae, SA 5064, Australia

² Instituto de Tecnologia Química e Biológica, Universidade Nova de Lisboa, Genomics of Plant Stress Laboratory (GPlantS Lab), Av. da República, 2780-157 Oeiras, Portugal

³ Center for Desert Agriculture, Division of Biological and Environmental Sciences and Engineering, King Abdullah University of Science and Technology, Thuwal 23955-6900, Saudi Arabia

Corresponding author: Tester, Mark (mark.test@kaust.edu.sa)

Current Opinion in Biotechnology 2014, **26**:115–124

This review comes from a themed issue on **Plant biotechnology**

Edited by **Birger Lindberg Møller** and **R George Ratcliffe**

0958-1669/\$ – see front matter, © 2013 The Authors. Published by Elsevier Ltd. All rights reserved.

<http://dx.doi.org/10.1016/j.copbio.2013.12.004>

Introduction

Soil salinity is a major environmental constraint to crop production, affecting an estimated 45 million hectares of irrigated land, and is expected to increase due to global climate changes and as a consequence of many irrigation practices [1,2]. The deleterious effects of salt stress on agricultural yield are significant, mainly because crops exhibit slower growth rates, reduced tillering and, over months, reproductive development is affected [2]. The ultimate aim of salinity tolerance research is to increase the ability of plants to maintain growth and productivity in saline soils relative to their

growth in non-saline soils — that is, to reduce effects of salinity on growth and yield. A range of biotechnologies can facilitate this by speeding gene (and allele) discovery and speeding the delivery of crops with improved salt tolerance (using both marker-assisted selection and genetic modification).

Growth is reduced by salinity via several quite distinct processes, which are related either to the accumulation of salt in the shoot, or which are independent of shoot salt accumulation. These can be experimentally distinguished by measuring effects immediately (within minutes to a few days) upon addition of salt (before there has been time for salt to accumulate in the shoot) or measured after much longer times (several days to weeks), after there has been time for salt to accumulate in the shoot and affect shoot growth. Within minutes of application of salt in an experimental system, there are several rapid responses occurring. Because of their rapid onset, these effects are clearly independent of the accumulation of salts in the shoot. The two best-documented effects are stomatal closure, with concomitant increases in leaf temperature [3]; and inhibition of shoot elongation [4,5]. Hence, the primary consequence is the overall reduction in production of new leaves and a significant reduction in shoot growth. This was termed the ‘osmotic phase’ by Munns and Tester [2], but there is evidence consistent with it not being due just to the effect of salt on water potential [4]. It is perhaps better described as a ‘shoot salt accumulation independent effect’.


In the second phase of plant responses to salinity, there is a slower onset inhibition of growth (occurring over several days to weeks), which is due to accumulation over time of salt, especially in the older leaves, causing the premature senescence of those older leaves. This is termed the ‘ionic phase’ of salt toxicity. This is due to both the accumulation of salts, and the inability of the shoot to tolerate the salt that has accumulated to toxic concentrations [2].

Mechanisms of salinity tolerance — overview

Just as salinity has many different effects on a plant, so there are also many mechanisms for plants to tolerate this stress. These mechanisms can be classified into three main categories: firstly, osmotic tolerance, which is regulated by long distance signals that reduce shoot growth and is triggered before shoot Na^+ accumulation; secondly, ion exclusion, where Na^+ and Cl^- transport processes in roots reduce the accumulation of toxic concentrations of Na^+ and Cl^- within leaves; and finally, tissue tolerance, where high salt concentrations are found

[☆] This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial-No Derivative Works License, which permits non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

Figure 1



The three main mechanisms of salinity tolerance in a crop plant. Tissue tolerance, where high salt concentrations are found in leaves but are compartmentalized at the cellular and intracellular level (especially in the vacuole), a process involving ion transporters, proton pumps and synthesis of compatible solutes. Osmotic tolerance, which is related to minimizing the effects on the reduction of shoot growth, and may be related to as yet unknown sensing and signaling mechanisms. Ion exclusion, where Na^+ and Cl^- transport processes, predominantly in roots, prevent the accumulation of toxic concentrations of Na^+ and Cl^- within leaves. Mechanisms may include retrieval of Na^+ from the xylem, compartmentation of ions in vacuoles of cortical cells and/or efflux of ions back to the soil.

in leaves but are compartmentalized at the cellular and intracellular level (especially in the vacuole) (Figure 1).

Very little, if anything, is known about tolerance to the 'osmotic phase'. This process must involve rapid, long-distance signaling, perhaps via processes such as ROS waves [6], Ca^{2+} waves (Simon Gilroy, personal communication), or even long distance electrical signaling [7]. Differences in osmotic tolerance may be due to differences in this long-distance signaling, or they may involve differences in the initial perception of the salt or differences in the response to the signals. This is still an area of salinity research with many unknowns, and further research is required to obtain a better understanding of osmotic tolerance (Figure 1).

Much more is known about the 'ionic phase', which is due to the accumulation of Na^+ and Cl^- in the leaf blade (a trait that is relatively easy to phenotype). Plants can reduce toxicity during the ionic phase by reducing accumulation of toxic ions in the leaf blades (Na^+ and

Cl^- exclusion), and/or by increasing their ability to tolerate the salts that they have failed to exclude from the shoot, such as by compartmentation into vacuoles (tissue tolerance) (Figure 1). Both of these processes involve a range of transporters and their controllers at both plasma membrane and tonoplast [8,9]. Tissue tolerance, involving the removal of Na^+ from the cytosol and compartmentalizing it in the vacuole before the ion has a detrimental effect on cellular processes, is also likely to require the synthesis of compatible solutes and higher level controls to coordinate transport and biochemical processes, thus having a role in both osmoprotection and osmotic adjustment [2,10] (Figure 1).

Suffice to say, it is clear from this brief overview that there are many mechanisms of salinity tolerance, and many of these can be present in a particular plant. To date, there is neither evidence that these mechanisms are mutually exclusive (i.e. that, e.g., ion exclusion prevents tolerance to the 'osmotic phase' of salt toxicity), nor that a particular plant is committed to only one strategy (e.g. a plant may

have ion exclusion as its primary tolerance mechanism at moderate salinity, but then has tissue tolerance as its main tolerance mechanism when the exclusion processes are 'swamped' at high salinity). It is possible that some tolerance mechanisms are more effective in particular circumstances. For example, Na⁺ exclusion may be more effective in conditions of higher salinity (e.g. see [11^{••}]), whereas 'osmotic tolerance' may be more important in moderately saline conditions. Interactions with other abiotic stresses, such as low water availability, are also likely to be important.

As such, it is clear that salinity tolerance can be complex and involve many genes, as has been pointed out for several decades — for example, programs designed to specifically introgress salinity tolerance using traditional breeding methods appear to have frequently failed (as measured by the apparent absence of commercial products), which has usually been attributed to the multi-genic nature of salinity tolerance [12,13]. It is therefore necessary not to study the molecular genetic basis of salinity tolerance as a particular trait in itself, but to study the mechanisms of traits that are hypothesized to contribute to salinity tolerance. The most intensively studied of these traits is exclusion of Na⁺ from leaf blades, mainly because it is relatively straightforward to phenotype. Focusing on this has led to significant increases in salinity tolerance, as measured by yield in the field, at least in durum wheat (which has poor Na⁺ exclusion) when grown in highly saline sites where yield has already been greatly reduced [11^{••},14].

Differences in traits must then be correlated with differences in tolerance, as measured by performance in the field — yield in a saline site relative to yield in a less saline site. Traits can be measured in any system that enables the trait to be quantified, using necessary experimental manipulations, such as described earlier to access the 'osmotic tolerance' trait — key is to test the relevance of the trait being measured with salinity tolerance in field conditions.

Candidate genes likely to contribute to tolerance traits in crops

There are numerous candidate genes that might be usefully used to transform crops to improve salinity tolerance. We focus in this review on the three main traits that are proposed by Munns and Tester [2] to be the primary mechanisms for salinity tolerance — shoot ion exclusion, shoot tissue tolerance and 'osmotic' tolerance — and propose genes that might contribute to each of these traits. A particular gene (or gene family) may well contribute to more than one trait, just as one trait can be conferred by more than one gene. We discuss a particular gene family in the section where it is likely to be making a significant contribution — this does not preclude members of this gene family also contributing to other traits,

and we try to acknowledge this in the text below. We have taken this approach to try to encourage a focus more on the traits that contribute to salinity tolerance, rather than on the traditional functional categorizations of gene families. This is an attempt to come more 'from the plant's perspective' of functional effects of the genes at the whole plant level, rather than categorizing genes based on the immediate activity of their encoded proteins.

- Ion exclusion

The *high affinity potassium transporter (HKT)* gene family [2,15–22] and the salt overly sensitive (SOS) pathway [23–29] have both been implicated in having an important role in regulating Na⁺ transport within a plant. Manipulation of the expression of these genes has been frequently reported to alter accumulation of Na⁺ in the shoot (Table 1 and Supplementary Table 1). However, to date the application of this knowledge into generating successful crop plants in the field has been limited. Of the two families, the *HKT1* group of *HKTs* have perhaps the greatest potential for improving the salinity tolerance of crops, frequently appearing as the most likely candidate for quantitative trait loci when phenotyping for salt tolerance and/or Na⁺ exclusion in mutant and mapping populations [30–33]. A marker assisted selection (MAS) approach was used successfully to incorporate novel *HKT* alleles from *Triticum monococcum* to improve the salinity tolerance of durum wheat [11^{••},14]. In contrast, transgenic approaches to improve salinity tolerance using *HKTs* have been only moderately successful (Table 1 and Supplementary Table 1). An *HKT2* has been reported to increase salinity tolerance, although not through Na⁺ exclusion [34[•]]. These genes appear to require cell type specific expression to be effective (Table 1 and Supplementary Table 1) [35]. If stress inducible and cell type specific expression of these genes can be realized in an effective way in crops, the potential for improving crop salinity tolerance through ion exclusion is possible.

- Shoot tissue tolerance

To date, three main mechanisms contributing to shoot tissue tolerance have been targeted: accumulation of Na⁺ in the vacuole, synthesis of compatible solutes and production of enzymes catalyzing detoxification of reactive oxygen species (Table 1 and Supplementary Table 2). Increasing the abundance of vacuolar Na⁺/H⁺ antiporters (NHX), vacuolar H⁺ pyrophosphatases (e.g. AVP1), proteins involved in the synthesis of compatible solutes (such as proline and glycinebetaine) and enzymes responsible for the detoxification of reactive oxygen species have had differing degrees of success in improving crop salinity tolerance (Table 1 and Supplementary Table 2). Although there is still uncertainty about the primary ions being transported by NHX proteins *in planta* [36,37,38[•]] (and the potential role of these proteins in K⁺ transport needs

Table 1

Genes that have been overexpressed to improve specific salinity tolerance traits in crops

Transgene	Gene isolated from	Promoters used	Transgenic crop	Reported transgenic plant performance during salt stress
<i>Ion exclusion (transporters)</i>				
Na ⁺ /H ⁺ antiporter (<i>SOS1</i>)	<i>Arabidopsis</i>	Constitutive	Tobacco	Altered shoot and root accumulation of Na ⁺ and K ⁺
Na ⁺ /H ⁺ antiporter (<i>SOD2</i>)	<i>Salicornia brachiata</i>	Stress inducible	Rice	Improved biomass production
Na ⁺ transporter (<i>HKT</i> subfamily 1)	Yeast	ABA responsive	Barley	Improved germination
Na ⁺ /K ⁺ transporter (<i>HKT</i> subfamily 2)	Barley			
Na ⁺ ATPase (<i>ENA</i>)	<i>Physcomitrella patens</i>			
<i>Tissue tolerance (transporters/proton pumps)</i>				
Na ⁺ /H ⁺ antiporter (<i>NHX</i>)	<i>Arabidopsis</i>	Constitutive	Buckwheat	Improved shoot and root biomass production
Na ⁺ /H ⁺ antiporter (<i>nhaA</i>)	<i>Atriplex gmelini</i>		Cotton	Altered Na ⁺ and K ⁺ accumulation
Vacuolar H ⁺ pyrophosphatase (<i>vacuolar H⁺-PPase</i>)	Rice Cotton <i>Hordeum brevisubulatum</i> <i>Pennisetum glaucum</i> <i>Aeluropus littoralis</i> <i>Salicornia brachiata</i> <i>Salsola soda</i> <i>Malus domestica</i> <i>E. coli</i>		Tomato Poplar Kiwifruit Fescue Rice Wheat Brassica Bentgrass Sugar beet Alfalfa Tobacco Apple	Increased proline content
<i>Tissue tolerance (Compatible solutes)</i>				
Trehalose-6-phosphate synthase (<i>TPS</i>)	Yeast	Constitutive	Alfalfa	Increased compatible solute accumulation
Trehalose-6-phosphate phosphatase (<i>TPP</i>)	Rice	Stress inducible	Tomato	Improved plant survival
Mannitol-1-phosphate dehydrogenase (<i>mt1D</i>)	<i>E. coli</i>	Shoot expression	Rice	Increased growth
L-myo-Inositol-1-phosphate synthase (<i>MIP</i>)	<i>Porteresia coarctata</i>	Protein often targeted to chloroplast	Tobacco	Reduced wilting
Myo-inositol O-methyltransferase	<i>Mesembryanthemum crystallinum</i>		Wheat	Maintenance of photosynthetic efficiency
Betaine aldehyde dehydrogenase (<i>BADH</i>)	Spinach		Sweet potato	
Choline oxidase/dehydrogenase (<i>codA/betA</i>)	Moth bean		Wheat	
Δ1-pyrroline-5-carboxylate synthetase (<i>P5CS</i>)			Potato	
<i>Tissue tolerance (Degradation of reactive oxygen species)</i>				
Ascorbate peroxidase (<i>APX</i>)	<i>Arabidopsis</i>	Constitutive	Tobacco	Maintenance of photosynthetic efficiency
Glutathione S-transferase (<i>GST</i>)	Tomato	Protein often targeted to chloroplast or cytosol	Rice	Maintenance of growth
Superoxide dismutase monodehydroascorbate reductase (<i>MDR</i>)	Tobacco			Improved maintenance of photosynthesis
Catalase	Mangrove Pea <i>E. coli</i>			Improved germination Improved growth of seedlings Increased antioxidant enzyme activity
<i>Signaling/regulatory pathways</i>				
Calcineurin-B like interacting protein kinases (<i>CIPK</i>)	<i>Arabidopsis</i>	Constitutive	Barley	Altered Na ⁺ , K ⁺ and Cl ⁻ accumulation
Mitogen-activated protein kinase (<i>MAPK</i>)	Chickpea		Tobacco	Improved biomass production

Table 1 (Continued)

Transgene	Gene isolated from	Promoters used	Transgenic crop	Reported transgenic plant performance during salt stress
Sucrose nonfermenting1-1 type protein kinase	Rice		Rice	Reduced leaf senescence
	Apple		Tomato	
	Cotton			
	Tomato			
<i>Transcription factors</i> <i>DREB</i>	<i>Pennisetum glaucum</i>	Constitutive and inducible	Tobacco	Improved germination
<i>AP2/ERF</i>	Cotton		Wheat	Improved biomass
<i>MYB</i>	Soybean		Tomato	Improved chlorophyll retention
<i>NAC</i>	Tomato		Rice	Altered Na ⁺ accumulation
	Rice			
	Chrysanthemum			
	Wheat			

to be kept in mind [38[•]]), and a new role has recently been proposed for AVP1 [39[•]], salinity tolerant plants appear to have been developed by the overexpression of *NHX* and vacuolar pyrophosphatase genes (Table 1 and Supplementary Table 2). While approaches to improve the tissue tolerance of crops through increasing compatible solutes (Supplementary Table 2) and enzymes involved in ROS metabolism (Table 1 and Supplementary Table 2) also appear to have been successful, there are often reports of low performance by the transgenic lines in low stress environments [40–45]. Such negative effects might be avoided by use of tightly regulated stress-inducible promoters [41,46–49].

With a large number of papers reporting success in increasing plant salinity tolerance by improving shoot tissue tolerance (Table 1 and Supplementary Table 2) in the greenhouse or growth room, particularly by improving accumulation of Na⁺ in vacuoles, it might be tempting to suggest this is the best mechanism for improving crop performance. However, there is, as yet, insufficient quantitative data and field trial data available to be able to draw such a conclusion. Exclusion of ions from the shoot has frequently been shown to be an important salt tolerance mechanism. MAS for improved salinity tolerance through improved exclusion has been successful in the field [11^{••},14]. The limited success to date of transgenic manipulations to increase ion exclusion, for example using the *HKT* gene family, is more likely to be due to the inability to express important exclusion genes in a cell type specific manner rather than a reflection of either the candidate gene or the effectiveness of the tolerance mechanism. *HKT1* proteins are reported to affect salinity tolerance by retrieving Na⁺ from the xylem in roots [16,17,19–22]. However, it is often the case that genes encoding transporters involved in the transport of Na⁺ across the plasma membrane cannot be constitutively overexpressed, unlike those genes encoding transporters involved in ion tissue tolerance that transport Na⁺ across

the vacuole. The cell type in which the transport of Na⁺ across the plasma membrane occurs has a fundamental effect on the overall accumulation in the whole plant [9]. To retrieve Na⁺ from the root xylem, enhanced Na⁺ influx into stelar cells are required — a phenotype that is opposite to that desired in root epidermal cells, where Na⁺ influx should be minimized. Therefore successful manipulation of transporter genes involved in ion exclusion processes requires a cell type-specific expression, which is not yet possible in most crops.

- Osmotic tolerance

We are currently not aware of any specific candidate genes for osmotic tolerance, although some genes highlighted above may be involved. Differences in osmotic tolerance are likely to involve long-distance signaling, control of cell cycle and processes involving perception of signals from the roots in the shoots. Given crop plants will usually be exposed to low levels of salinity throughout the growing season, or levels of salinity that start low but which build up toward the end of the growing season, genes encoding for osmotic tolerance traits have the potential to have a significantly larger impact on the salinity tolerance of crops than those involved in ion exclusion. The introgression of *TmHKT1;5-A* from *T. monococcum* into the durum wheat, Tamaroi, resulted in a significant improvement in grain yield in salt stressed, field grown durum by increasing its ion exclusion, but only in plots with highly saline soils [11^{••},14]. However, the yield of Tamaroi with *TmHKT1;5-A* was similar to that observed in the Tamaroi cultivar without the introgressed gene, under low and moderate saline conditions [11^{••},14] suggesting that osmotic stress was having a greater effect on the end yield of these plants growing in low to moderate salinity, than ionic stress. The identification of genes for osmotic tolerance should therefore be a priority for improving salinity tolerance of crops growing in low to moderate saline soils.

- Signaling/regulatory pathways

An attractive alternative to manipulating specific salinity stress tolerance mechanisms is to adjust the detection, signaling and regulatory pathways involved in global salinity tolerance. Potentially altering one component of these pathways can have significant secondary effects for downstream processes, such as ion exclusion, tissue tolerance and osmotic tolerance. ROS signaling has recently been shown to be important for regulating plant responses to many abiotic stresses [6^{*},50^{*}], and is involved controlling shoot Na⁺ accumulation by regulating vasculature Na⁺ concentrations [51^{*}]. Many aspects of plant growth and development are mediated by Ca²⁺. Environmental cues are perceived by receptors on the cell membrane, activating a Ca²⁺ signaling cascade, resulting in the regulation of gene expression and protein activities [23,24,52–54]. Overexpression of genes encoding proteins in Ca²⁺ signaling pathways have been shown to improve the growth of crop plants, such as rice, apple, barley, tobacco and tomato during salt stress (Table 1 and Supplementary Table 3).

Similarly, overexpression of transcription factors as a viable way of improving crop salinity tolerance has also been described numerous times in model species, such as *Arabidopsis* [55–58], and to a lesser extent in crops, such as rice, wheat, tomato and alfalfa (Table 1 and Supplementary Table 3). As key transcription factors may regulate the induction and/or repression of a range of salinity tolerance genes and mechanisms, it is tempting to suggest that manipulation of transcription factors may result in the greatest effect on crop salinity tolerance with the least amount of genetic modification. However, as for genes encoding compatible solute synthesis, yield penalties can often be apparent in low stress environments when transcription factors are constitutively expressed [59–61], presumably due to the effects of the transcription factors on a wide range of processes. Therefore, while these genes maybe have the potential to have the greatest effect on improving salinity tolerance, care should be taken in manipulating the expression levels of transcription factors. Use of stress inducible promoters may address many of these issues.

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.copbio.2013.12.004>.

It is important to recognize that different tissues and cells in a plant are adapted for specific and often very different purposes. It is therefore not surprising that the expression levels of genes will be different from tissue to tissue and cell to cell, depending on the cell's function — as an obvious example, genes involved in photosynthesis will not normally be expressed in cells not exposed to light, for example roots. Care must therefore be taken in the choice

of promoter used to drive the expression of salinity tolerance transgenes, so as not to disrupt or negatively affect the plant phenotype. There exist many examples where the constitutive expression of genes encoding transcription factors, ion transporters or proteins involved in compatible solute synthesis can lead to undesirable phenotypes, especially under non-stressed conditions [35,40–45,47,55,59–62]. However, when these genes are controlled by a promoter that is either cell type specific and/or stress inducible or have a tag on the protein that directs it to the correct cellular organelle, then the desired salt tolerant phenotype can be observed [35,40,42,48,49,55,63–68]. It is therefore imperative that promoters of genes which allow cell type and/or stress inducible expression are identified for controlling the expression of genes which can be used to obtain the desired phenotype. The gene is only half the story.

It is also important to recognize that altering expression of a gene is only one way to alter the protein function. In addition to allelic variation of the primary protein sequence, post-translational modifications that activate or suppress the protein's activity along with the location of the protein within a cell are also important factors that can be modified to alter a plant's salinity tolerance [24,69–71]. The calcium signaling pathways, which involve protein kinases to phosphorylate key proteins involved in salinity tolerance (such as SOS1) [23,24,26,27,52,72,73^{*}], are a good example of where post-translation modifications can enable a plant to respond quickly (and reversibly) to environmental changes.

It must be made clear that the choice of salinity tolerance mechanism to manipulate in crops plants will be largely dependent on the underlying salt tolerance mechanisms within individual crop species. For example, barley has greater salinity tolerance than wheat, and a greater ability to accumulate high concentrations of Na⁺ in its shoot (higher tissue tolerance). Barley may therefore benefit from the addition of, for example, osmotic tolerance mechanisms, rather than improved ion exclusion. However, transferring the tissue tolerance mechanisms of barley into crops such as wheat and rice may significantly increase salinity tolerance of these crops. Again, incorporation of osmotic tolerance and improved signaling traits may also benefit these species. Unfortunately, there is not enough data available to make critical comments on which salinity tolerance strategy would work best for a certain crop. Proper field evaluations of crops developed by MAS or transgenic approaches are now required to determine the effect of the modification on crop yield and the best strategy to improve the salinity tolerance of specific crop species. To date very few studies have been able to examine the effect of modifying salinity tolerance traits on the yield of field grown crop plants, and assess their yield penalty in non-saline soils.

Delivery to crops

Discoveries of mechanisms of salinity tolerance now need to be applied to crops to improve crop performance in the field. One way that this can be done is by using conventional breeding, accelerated by the use of molecular markers linked to the tolerance trait being introgressed, MAS. The effectiveness of the research outputs in salinity tolerance can be observed in the newly developed durum wheat lines with *TmHKT1;5-A* (a gene conferring Na⁺ exclusion) introgressed by MAS, which showed a yield increase of 25% when compared to control wheat, at least when grown in saline fields [11^{**},14]. Several studies evaluated natural allelic variants of HKT genes in response to salinity in *Arabidopsis* [33,74^{*}], rice [32,75^{*},76,77] and barley [78]. This allelic variation in the gene of interest can provide both a novel source of genetic material for MAS, and can also be used to design an easy, cheap molecular marker, thus providing a perfect marker for salinity tolerance — or, at least, for a trait contributing to salinity tolerance. If variation in a particular gene is insufficient within the relevant germplasm or if the gene is only found in a model plant species [79^{*}], then alteration of the trait using GM technologies can be deployed. It is essential that those experiments involving GM plants have not only greenhouse evaluation of the salinity tolerance of the plant but field evaluation as well. A plant that shows salinity tolerant in the greenhouse may not necessarily have improved grain yield in the field. Only a few studies (e.g. [80^{**},81]) have evaluated the performance of GM plants for any abiotic stress tolerance mechanism in the field. It also must be taken into consideration that a GM approach for edible crops may not be desirable due to public perception; therefore the use of biotechnology techniques that will not generate a GM crop may often be more appropriate.

These emerging biotechnologies include ones where plants go through a GM phase, then have the GM construct segregated back out. Designer nuclease-based approaches to alter genetic composition, such as Transcription Activator-Like Effector Nucleases (TALEN) [82–85], are reported to increase the mutagenesis efficiency of endogenous targets by specifying their binding sites with single nucleotide precision [85]. The potential promise of this strategy was tested in rice to enhance disease resistance [86]. Another emerging genome engineering tool is the CRISPR/Cas system (short for Clustered Regularly Interspersed Short Palindromic Repeats (CRISPR)/CRISPR-associated (Cas)) [87,88]. Several groups are optimizing and exploring the potential of Cas9 RNA-guided endonuclease system in human cells [89,90] and zebra fish [91]. These approaches are examples of new technologies that give promise to the future of crop biotechnology, whilst avoiding the presence of problematic transgenes or antibiotic markers.

To conclude, salinity tolerance is too complex to be easily amenable to improvement through selection as a trait in itself, but traits that are hypothesized to contribute to salinity tolerance are more genetically tractable and genes underlying these can be discovered using molecular genetics tools and genomics. Alterations in crops can then be made using both marker-assisted selection and genetic modification, and the relevance of such traits on whole plant salinity tolerance can then be tested, as measured by yield maintenance in saline conditions.

Acknowledgements

The authors would like to thank the Australian Research Council (SR and MT), Grains Research and Development Corporation (SR and MT), FCT-Portugal (grant SFRH/BPD/34593/2007) (SN) and KAUST (MT and SN) for their generous support.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Rengasamy P: **Soil processes affecting crop production in salt-affected soils.** *Funct Plant Biol* 2010, **37**:613-620.
2. Munns R, Tester M: **Mechanisms of salinity tolerance.** *Annu Rev Plant Biol* 2008, **59**:651-681.
3. Sirault XRR, James RA, Furbank RT: **A new screening method for osmotic component of salinity tolerance in cereals using infrared thermography.** *Funct Plant Biol* 2009, **36**:970-977.
4. Munns R, Passioura JB: **Hydraulic resistance of plants. 3. Effects of NaCl in barley and lupin.** *Aust J Plant Physiol* 1984, **11**:351-359.
5. Rajendran K, Tester M, Roy SJ: **Quantifying the three main components of salinity tolerance in cereals.** *Plant Cell Environ* 2009, **32**:237-249.
6. Mittler R, Vanderauwera S, Suzuki N, Miller G, Tognetti VB, Vandepoel K, Gollery M, Shulaev V, Van Breusegem F: **ROS signaling: the new wave?** *Trends Plant Sci* 2011, **16**:300-309.
7. Maischak H, Zimmermann MR, Felle HH, Boland W, Mithofer A: **Alamethicin-induced electrical long distance signaling in plants.** *Plant Signal Behav* 2010, **5**:988-990.
8. Plett DC, Moller IS: **Na⁺ transport in glycophytic plants: what we know and would like to know.** *Plant Cell Environ* 2010, **33**:612-626.
9. Tester M, Davenport R: **Na⁺ tolerance and Na⁺ transport in higher plants.** *Ann Bot* 2003, **91**:503-527.
10. Flowers TJ, Colmer TD: **Salinity tolerance in halophytes.** *New Phytol* 2008, **179**:945-963.
11. Munns R, James RA, Xu B, Athman A, Conn SJ, Jordans C, Byrt CS, Hare RA, Tyerman SD, Tester M *et al.*: **Wheat grain yield on saline soils is improved by an ancestral Na⁺ transporter gene.** *Nat Biotechnol* 2012, **30**:360-364.
12. Dewey DR: **Breeding crested wheatgrass *Agropyron desertorum* for salt tolerance.** *Crop Sci* 1962, **2**:403-407.

An example where the use of non-GM gene technology has been used to develop salt-tolerant wheat. A novel allele of the root stelar-specific Na⁺-selective transporter *HKT1;5* (*TmHKT1;5-A*) was introduced into commercial durum wheat from a near wild relative (*T. monococcum*) using marker assisted selection. Durum wheat with the *TmHKT1;5-A* gene had lower leaf [Na⁺] and a 25% increase in yield when grown on saline soils (when compared to near-isogenic lines without *TmHKT1;5-A*).

13. Flowers TJ, Yeo AR: **Breeding for salinity resistance in crop plants: where next?** *Aust J Plant Physiol* 1995, **22**:875-884.
14. James RA, Blake C, Zwart AB, Hare RA, Rathjen AJ, Munns R: **Impact of ancestral wheat sodium exclusion genes *Nax1* and *Nax2* on grain yield of durum wheat on saline soils.** *Funct Plant Biol* 2012, **39**:609-618.
15. Platten JD, Cotsaftis O, Berthomieu P, Bohnert H, Davenport RJ, Fairbairn DJ, Horie T, Leigh RA, Lin H-X, Luan S *et al.*: **Nomenclature for HKT transporters, key determinants of plant salinity tolerance.** *Trends Plant Sci* 2006, **11**:372-374.
16. Horie T, Hauser F, Schroeder JI: **HKT transporter-mediated salinity resistance mechanisms in *Arabidopsis* and monocot crop plants.** *Trends Plant Sci* 2009, **14**:660-668.
17. Hauser F, Horie T: **A conserved primary salt tolerance mechanism mediated by HKT transporters: a mechanism for sodium exclusion and maintenance of high K^+/Na^+ ratio in leaves during salinity stress.** *Plant Cell Environ* 2010, **33**:552-565.
18. Ali Z, Park HC, Ali A, Oh D-H, Aman R, Kropornicka A, Hong H, Choi W, Chung WS, Kim W-Y *et al.*: ***TsHKT1;2*, a *HKT1* homolog from the extremophile *Arabidopsis* relative *Thellungiella salsuginea*, shows K^+ specificity in the presence of NaCl.** *Plant Physiol* 2012, **158**:1463-1474.
19. Byrt CS, Platten JD, Spielmeier W, James RA, Lagudah ES, Dennis ES, Tester M, Munns R: ***HKT1;5*-like cation transporters linked to Na^+ exclusion loci in wheat, *Nax2* and *Kna1*.** *Plant Physiol* 2007, **143**:1918-1928.
20. Davenport RJ, Munoz-Mayor A, Jha D, Essah PA, Rus ANA, Tester M: **The Na^+ transporter *AtHKT1;1* controls retrieval of Na^+ from the xylem in *Arabidopsis*.** *Plant Cell Environ* 2007, **30**:497-507.
21. Horie T, Horie R, Chan W-Y, Leung H-Y, Schroeder JI: **Calcium regulation of sodium hypersensitivities of *sos3* and *athkt1* mutants.** *Plant Cell Physiol* 2006, **47**:622-633.
22. Xue S, Yao X, Luo W, Jha D, Tester M, Horie T, Schroeder JI: ***AtHKT1;1* mediates nernstian sodium channel transport properties in *Arabidopsis* root stelar cells.** *PLoS ONE* 2011, **6**:e24725.
23. Kudla J, Batistic O, Hashimoto K: **Calcium signals: the lead currency of plant information processing.** *Plant Cell* 2010, **22**:541-563.
24. Weinel S, Kudla J: **The CBL-CIPK Ca^{2+} -decoding signaling network: function and perspectives.** *New Phytol* 2009, **184**:517-528.
25. Mahajan S, Pandey GK, Tuteja N: **Calcium- and salt-stress signaling in plants: shedding light on SOS pathway.** *Arch Biochem Biophys* 2008, **471**:146-158.
26. Qiu Q-S, Guo Y, Dietrich MA, Schumaker KS, Zhu J-K: **Regulation of *SOS1*, a plasma membrane Na^+/H^+ exchanger in *Arabidopsis thaliana*, by *SOS2* and *SOS3*.** *Proc Natl Acad Sci U S A* 2002, **99**:8436-8441.
27. Ji H, Pardo JM, Batelli G, Van Oosten MJ, Bressan RA, Li X: **The salt overly sensitive (SOS) pathway: established and emerging roles.** *Mol Plant* 2013, **6**:275-286.
28. Shi H, Lee B-h, Wu S-J, Zhu J-K: **Overexpression of a plasma membrane Na^+/H^+ antiporter gene improves salt tolerance in *Arabidopsis thaliana*.** *Nat Biotechnol* 2003, **21**:81-85.
29. Yang Q, Chen Z-Z, Zhou X-F, Yin H-B, Li X, Xin X-F, Hong X-H, Zhu J-K, Gong Z: **Overexpression of *SOS* (salt overly sensitive) genes increases salt tolerance in transgenic *Arabidopsis*.** *Mol Plant* 2009, **2**:22-31.
30. Ahmadi N, Negrão S, Katsantonis D, Frouin J, Ploux J, Letourmy P, Droc G, Babo P, Trindade H, Bruschi G *et al.*: **Targeted association analysis identified *japonica* rice varieties achieving Na^+/K^+ homeostasis without the allelic make-up of the salt tolerant *indica* variety Nona Bokra.** *Theor Appl Genet* 2011, **123**:881-895.
31. James RA, Davenport RJ, Munns R: **Physiological characterization of two genes for Na^+ exclusion in durum wheat, *Nax1* and *Nax2*.** *Plant Physiol* 2006, **142**:1537-1547.
32. Ren ZH, Gao JP, Li LG, Cai XL, Huang W, Chao DY, Zhu MZ, Wang ZY, Luan S, Lin HX: **A rice quantitative trait locus for salt tolerance encodes a sodium transporter.** *Nat Genet* 2005, **37**:1141-1146.
33. Rus A, Baxter I, Muthukumar B, Gustin J, Lahner B, Yakubova E, Salt DE: **Natural variants of *AtHKT1* enhance Na^+ accumulation in two wild populations of *Arabidopsis*.** *PLoS Genet* 2006, **2**:1964-1973.
34. Mian A, Oomen RJFJ, Isayenkov S, Sentenac H, Maathuis FJM, Véry A-A: **Over-expression of an Na^+ - and K^+ -permeable HKT transporter in barley improves salt tolerance.** *Plant J* 2011, **68**:468-479.
- A study in which the electrophysiological properties of *HvHKT2;1* are characterized, along with the expression profile of the native *HvHKT2;1* gene in barley, before generating transgenic barley overexpressing *HvHKT2;1* to determine its role in salinity tolerance. Interestingly, constitutive overexpression of this subfamily 2 *HKT* gene in barley results in increased shoot Na^+ concentrations and improved biomass production under salt stress (at least in the conditions of the assay used in this work), in contrast to what is observed when subfamily 1 *HKT* genes are constitutively expressed in *Arabidopsis* [89].
35. Møller I, Gilliam M, Jha D, Mayo G, Roy SJ, Coates J, Haseloff J, Tester M: **Salinity tolerance engineered by cell type-specific over-expression of a Na^+ transporter in the *Arabidopsis* root.** *Plant Cell* 2009, **21**:2163-2178.
36. Bassil E, Ohto M-a, Esumi T, Tajima H, Zhu Z, Cagnac O, Belmonte M, Peleg Z, Yamaguchi T, Blumwald E: **The *Arabidopsis* intracellular Na^+/H^+ antiporters *NHX5* and *NHX6* are endosome associated and necessary for plant growth and development.** *Plant Cell* 2011, **23**:224-239.
37. Bassil E, Tajima H, Liang Y-C, Ohto M-a, Ushijima K, Nakano R, Esumi T, Coku A, Belmonte M, Blumwald E: **The *Arabidopsis* Na^+/H^+ antiporters *NHX1* and *NHX2* control vacuolar pH and K^+ homeostasis to regulate growth, flower development, and reproduction.** *Plant Cell* 2011, **23**:3482-3497.
38. Barragán V, Leidi EO, Andrés Z, Rubio L, De Luca A, Fernández JA, Cubero B, Pardo JM: **Ion exchangers *NHX1* and *NHX2* mediate active potassium uptake into vacuoles to regulate cell turgor and stomatal function in *Arabidopsis*.** *Plant Cell* 2012, **24**:1127-1142.
- Members of the Na^+/H^+ antiporter gene family (*NHX*), particularly *AtNHX1*, have frequently been overexpressed in plants to improve vacuolar sequestration of Na^+ . In this study, the *Arabidopsis* double knockout mutant *nhx1 nhx2* showed high sensitivity to external K^+ concentrations but not high external Na^+ , suggesting a role for these proteins in the vacuolar sequestration of K^+ .
39. Ferjani A, Segami S, Horiguchi G, Muto Y, Maeshima M, Tsukaya H: **Keep an eye on PPI: the vacuolar-type H^+ -pyrophosphatase regulates postgerminative development in *Arabidopsis*.** *Plant Cell* 2011, **23**:2895-2908.
- This paper challenges the traditional view that the main role of vacuolar inorganic pyrophosphatases (PPases), such as AVP1, is vacuolar acidification. Work on the *fugu5* vacuolar H^+ -PPase mutant suggests that a primary role of PPases is to remove inorganic pyrophosphate that inhibits gluconeogenesis during germination. Transformation of *fugu5* with a yeast soluble pyrophosphatase recovered the early growth inhibition effects resulting from a build-up of PPI in the cytosol.
40. Romero C, Bellés J, Vayá J, Serrano R, Culiáñez-Macià F: **Expression of the yeast *trehalose-6-phosphate synthase* gene in transgenic tobacco plants: pleiotropic phenotypes include drought tolerance.** *Planta* 1997, **201**:293-297.
41. Sheveleva E, Chmara W, Bohnert HJ, Jensen RG: **Increased salt and drought tolerance by D-ononitol production in transgenic *Nicotiana tabacum* L.** *Plant Physiol* 1997, **115**:1211-1219.
42. Suárez R, Calderón C, Iturriaga G: **Enhanced tolerance to multiple abiotic stresses in transgenic Alfalfa accumulating trehalose.** *Crop Sci* 2009, **49**:1791-1799.
43. Cortina C, Culiáñez-Macià FA: **Tomato abiotic stress enhanced tolerance by trehalose biosynthesis.** *Plant Sci* 2005, **169**:75-82.
44. Karakas B, Ozias-Akins P, Stushnoff C, Sufferheld M, Rieger M: **Salinity and drought tolerance of mannitol-accumulating transgenic tobacco.** *Plant Cell Environ* 1997, **20**:609-616.

45. Hmida-Sayari A, Gargouri-Bouزيد R, Bidani A, Jaoua L, Savouré A, Jaoua S: **Overexpression of $\Delta 1$ -pyrroline-5-carboxylate synthetase increases proline production and confers salt tolerance in transgenic potato plants.** *Plant Sci* 2005, **169**:746-752.
46. Garg AK, Kim J-K, Owens TG, Ranwala AP, Choi YD, Kochian LV, Wu RJ: **Trehalose accumulation in rice plants confers high tolerance levels to different abiotic stresses.** *Proc Natl Acad Sci U S A* 2002, **99**:15898-15903.
47. Vendruscolo ECG, Schuster I, Pileggi M, Scapim CA, Molinari HBC, Marur CJ, Vieira LGE: **Stress-induced synthesis of proline confers tolerance to water deficit in transgenic wheat.** *J Plant Physiol* 2007, **164**:1367-1376.
48. Zhu B, Su J, Chang M, Verma DPS, Fan Y-L, Wu R: **Overexpression of a $\Delta 1$ -pyrroline-5-carboxylate synthetase gene and analysis of tolerance to water- and salt-stress in transgenic rice.** *Plant Sci* 1998, **139**:41-48.
49. Su J, Wu R: **Stress-inducible synthesis of proline in transgenic rice confers faster growth under stress conditions than that with constitutive synthesis.** *Plant Sci* 2004, **166**:941-948.
50. Suzuki N, Koussevitzky S, Mittler R, Miller G: **ROS and redox signaling in the response of plants to abiotic stress.** *Plant Cell Environ* 2012, **35**:259-270.
- A comprehensive review on the current knowledge of ROS and redox signaling during abiotic stress. Discusses in depth the various systems involved in communication, development, metabolism and coordinating gene expression between the nucleus, chloroplasts and mitochondria. Proposes a model for the involvement of ROS in local and systemic signaling in abiotic stressed plants.
51. Jiang C, Belfield EJ, Mithani A, Visscher A, Ragoussis J, Mott R, Smith JAC, Harberd NP: **ROS-mediated vascular homeostatic control of root-to-shoot soil Na delivery in *Arabidopsis*.** *EMBO J* 2012, **31**:4359-4370.
- AtrbohF* mutants are shown to be salt sensitive, due to higher xylem sap Na^+ concentrations and thus increased shoot Na^+ accumulation. This provides interesting possibilities for linking salt-induced elevations in ROS with specific mechanisms of salinity tolerance.
52. Luan S: **The CBL-CIPK network in plant calcium signaling.** *Trends Plant Sci* 2009, **14**:37-42.
53. Dodd AN, Kudla J, Sanders D: **The language of calcium signaling.** *Annu Rev Plant Biol* 2010, **61**:593-620.
54. Batistić O, Kudla J: **Analysis of calcium signaling pathways in plants.** *Biochim Biophys Acta* 2012, **1820**:1283-1293.
55. Kasuga M, Liu Q, Miura S, Yamaguchi-Shinozaki K, Shinozaki K: **Improving plant drought, salt, and freezing tolerance by gene transfer of a single stress-inducible transcription factor.** *Nat Biotechnol* 1999, **17**:287-291.
56. Mao X, Jia D, Li A, Zhang H, Tian S, Zhang X, Jia J, Jing R: **Transgenic expression of *TaMYB2A* confers enhanced tolerance to multiple abiotic stresses in *Arabidopsis*.** *Funct Integr Genomics* 2011, **11**:445-465.
57. Yang W, Liu X-D, Chi X-J, Wu C-A, Li Y-Z, Song L-L, Liu X-M, Wang Y-F, Wang F-W, Zhang C *et al.*: **Dwarf apple *MbDREB1* enhances plant tolerance to low temperature, drought, and salt stress via both ABA-dependent and ABA-independent pathways.** *Planta* 2011, **233**:219-229.
58. Zhou J, Li F, Wang J-I, Ma Y, Chong K, Xu Y-y: **Basic helix-loop-helix transcription factor from wild rice (*OrbHLH2*) improves tolerance to salt- and osmotic stress in *Arabidopsis*.** *J Plant Physiol* 2009, **166**:1296-1306.
59. Liu Q, Kasuga M, Sakuma Y, Abe H, Miura S, Yamaguchi-Shinozaki K, Shinozaki K: **Two transcription factors, *DREB1* and *DREB2*, with an EREBP/AP2 DNA binding domain separate two cellular signal transduction pathways in drought- and low-temperature-responsive gene expression, respectively, in *Arabidopsis*.** *Plant Cell* 1998, **10**:1391-1406.
60. Morran S, Eini O, Pyvovarenko T, Parent B, Singh R, Ismagul A, Eliby S, Shirley N, Langridge P, Lopato S: **Improvement of stress tolerance of wheat and barley by modulation of expression of *DREB/CBF* factors.** *Plant Biotechnol J* 2011, **9**:230-249.
61. AbuQamar S, Luo H, Laluk K, Mickelbart MV, Mengiste T: **Crosstalk between biotic and abiotic stress responses in tomato is mediated by the *AIM1* transcription factor.** *Plant J* 2009, **58**:347-360.
62. Rus A, Lee B-h, Munoz-Mayor A, Sharkhuu A, Miura K, Zhu J-K, Bressan RA, Hasegawa PM: ***AtHKT1* facilitates Na^+ homeostasis and K^+ nutrition in *Planta*.** *Plant Physiol* 2004, **136**:2500-2511.
63. Mallikarjuna G, Mallikarjuna K, Reddy M, Kaul T: **Expression of *OsDREB2A* transcription factor confers enhanced dehydration and salt stress tolerance in rice (*Oryza sativa* L.).** *Biotechnol Lett* 2011, **33**:1689-1697.
64. Holmström KO, Somersalo S, Mandal A, Palva TE, Welin B: **Improved tolerance to salinity and low temperature in transgenic tobacco producing glycine betaine.** *J Exp Bot* 2000, **51**:177-185.
65. Lee YP, Baek K-H, Lee H-S, Kwak S-S, Bang J-W, Kwon S-Y: **Tobacco seeds simultaneously over-expressing Cu/Zn-superoxide dismutase and ascorbate peroxidase display enhanced seed longevity and germination rates under stress conditions.** *J Exp Bot* 2010, **61**:2499-2506.
66. Badawi GH, Kawano N, Yamauchi Y, Shimada E, Sasaki R, Kubo A, Tanaka K: **Over-expression of ascorbate peroxidase in tobacco chloroplasts enhances the tolerance to salt stress and water deficit.** *Physiol Plant* 2004, **121**:231-238.
67. Sakamoto A, Murata A: **Metabolic engineering of rice leading to biosynthesis of glycinebetaine and tolerance to salt and cold.** *Plant Mol Biol* 1998, **38**:1011-1019.
68. Verma D, Singla-Pareek S, Rajagopal D, Reddy MK, Sopory SK: **Functional validation of a novel isoform of Na^+/H^+ antiporter from *Pennisetum glaucum* for enhancing salinity tolerance in rice.** *J Biosci* 2007, **32**:621-628.
69. Castro P, Tavares R, Bejarano E, Azevedo H: **SUMO, a heavyweight player in plant abiotic stress responses.** *Cell Mol Life Sci* 2012, **69**:3269-3283.
70. Bond AE, Row PE, Dudley E: **Post-translation modification of proteins; methodologies and applications in plant sciences.** *Phytochemistry* 2011, **72**:975-996.
71. Mazzucotelli E, Mastrangelo AM, Crosatti C, Guerra D, Stanca AM, Cattivelli L: **Abiotic stress response in plants: when post-transcriptional and post-translational regulations control transcription.** *Plant Sci* 2008, **174**:420-431.
72. Halfter U, Ishitani M, Zhu J-K: **The *Arabidopsis* *SOS2* protein kinase physically interacts with and is activated by the calcium-binding protein *SOS3*.** *Proc Natl Acad Sci U S A* 2000, **97**:3735-3740.
73. Huertas R, Olias R, Eljakaoui Z, Galvez FJ, Li JUN, De Morales PA, Belver A, Rodríguez-Rosales MP: **Overexpression of *SISOS2* (*SICIPK24*) confers salt tolerance to transgenic tomato.** *Plant Cell Environ* 2012, **35**:1467-1482.
- Investigated the *SISOS2* gene and its role in the SOS pathway of tomato, *Solanum lycopersicum*. Confirmed that *SISOS2* protein interacts with members of the CBL family, specifically *SOS3* (*CBL4*) and could be used to recover the salt sensitive phenotype of *Atsos2-2 Arabidopsis* plants. Characterisation of transgenic tomato overexpressing *SISOS2* at the seedling, vegetative and flowering/fruitlet stage showed that these plants were able to maintain growth and produce more fruit under salt stress, while accumulating more shoot Na^+ , than non-transformed controls. The study also looked at the effect of overexpressing a single transgene on the expression of genes encoding downstream targets of *SISOS2* in an attempt to explain the observed phenotype.
74. Baxter I, Brazelton JN, Yu D, Huang YS, Lahner B, Yakubova E, Li Y, Bergelson J, Borevitz JO, Nordborg M *et al.*: **A coastal cline in sodium accumulation in *Arabidopsis thaliana* is driven by natural variation of the sodium transporter *AtHKT1;1*.** *PLoS Genet* 2010:6.
- Demonstrated that a weak allele of *AtHKT1;1* in *Arabidopsis*, associated with increased leaf $[\text{Na}^+]$, was found to be enriched in populations of *Arabidopsis* that were found in coastal and saline soils of Europe. They suggest that the fixation of this weaker allele of *AtHKT1;1* in these populations supports local adaptations to the environment, perhaps through enhanced tissue tolerance.

75. Negrão S, Almadanim MC, Pires IS, Abreu IA, Maroco J, Courtois B, Gregorio GB, McNally KL, Oliveira MM: **New allelic variants found in key rice salt-tolerance genes: an association study.** *Plant Biotechnol J* 2013, **11**:87-100.

This work provides new insights into allelic variants in five salt-related genes in 392 rice accessions. Forty new allelic variants were found at the coding sequence level, and seven significant SNPs (in *OsHKT1;5*, *OsNHX1* and *SaT*) leading to amino acid changes were associated with salinity tolerance. By addressing the putative consequences of these SNPs at the protein level, the authors hypothesize their functional relevance. This allelic variation can provide novel sources of genetic material to be used for MAS by rice breeders.

76. Oomen RJFJ, Benito B, Sentenac H, Rodriguez-Navarro A, Talon M, Very A-A, Domingo C: **HKT2;2/1, a K⁺-permeable transporter identified in a salt-tolerant rice cultivar through surveys of natural genetic polymorphism.** *Plant J* 2012, **71**:750-762.
77. Platten JD, Egdane JA, Ismail AM: **Salinity tolerance, Na⁺ exclusion and allele mining of HKT1;5 in *Oryza sativa* and *O. glaberrima*: many sources, many genes, one mechanism?** *BMC Plant Biol* 2013, **13**:32.
78. Qiu L, Wu D, Ali S, Cai S, Dai F, Jin X, Wu F, Zhang G: **Evaluation of salinity tolerance and analysis of allelic function of HvHKT1 and HvHKT2 in Tibetan wild barley.** *Theor Appl Genet* 2011, **122**:695-703.
79. Roy SJ, Huang W, Wang XJ, Evrard A, SchmÖckel SM, Zafar ZU, Tester M: **A novel protein kinase involved in Na⁺ exclusion revealed from positional cloning.** *Plant Cell Environ* 2013, **36**:553-568.

While many genes for improving plant salinity tolerance are known, new genes or novel alleles of known genes are still being discovered which can be used to improve crops. Using a forward genetics approach, the authors found that a difference in the expression of a CIPK was responsible for differences in shoot Na⁺ accumulation in *Arabidopsis*. Over-expressing of the *Arabidopsis* gene in barley resulted in reduced shoot Na⁺ accumulation and improved biomass production during salinity stress.

80. Pasapula V, Shen G, Kuppu S, Paez-Valencia J, Mendoza M, Hou P, Chen J, Qiu X, Zhu L, Zhang X *et al.*: **Expression of an *Arabidopsis* vacuolar H⁺-pyrophosphatase gene (*AVP1*) in cotton improves drought- and salt tolerance and increases fibre yield in the field conditions.** *Plant Biotechnol J* 2011, **9**:88-99.

A good example of how to perform research using transgenic crops, importantly using null segregants as biological controls in addition to wild

type plants. The authors not only show that the transgene was present and expressed in their transformed cotton but additionally demonstrated the location of the AVP1 protein. Transgenic cotton expressing AVP1 produces more biomass and have higher yields under both salinity and drought stress in the greenhouse, and also yielded more in dry-land field conditions over two years of field trials.

81. Schilling RK, Marschner P, Shavrukov Y, Berger B, Tester M, Roy SJ, Plett DC: **Expression of the *Arabidopsis* vacuolar H⁺-pyrophosphatase gene (*AVP1*) improves the shoot biomass of transgenic barley and increases grain yield in a saline field.** *Plant Biotechnol J* 2013 <http://dx.doi.org/10.1111/pbi.12145>.
82. Bogdanove AJ, Voytas DF: **TAL effectors: customizable proteins for DNA targeting.** *Science* 2011, **333**:1843-1846.
83. Christian M, Cermak T, Doyle EL, Schmidt C, Zhang F, Hummel A, Bogdanove AJ, Voytas DF: **Targeting DNA double-strand breaks with TAL effector nucleases.** *Genetics* 2010, **186**:757-761.
84. Curtin SJ, Voytas DF, Stupar RM: **Genome engineering of crops with designer nucleases.** *Plant Genome* 2012, **5**:42-50.
85. Voytas DF: **Plant genome engineering with sequence-specific nucleases.** *Annu Rev Plant Biol* 2013, **64**:327-350.
86. Li T, Liu B, Spalding MH, Weeks DP, Yang B: **High-efficiency TALEN-based gene editing produces disease-resistant rice.** *Nat Biotechnol* 2012, **30**:390-392.
87. Haurwitz RE, Jinek M, Wiedenheft B, Zhou K, Doudna JA: **Sequence- and structure-specific RNA processing by a CRISPR endonuclease.** *Science* 2010, **329**:1355-1358.
88. Wiedenheft B, Sternberg SH, Doudna JA: **RNA-guided genetic silencing systems in bacteria and archaea.** *Nature* 2012, **482**:331-338.
89. Cho SW, Kim S, Kim JM, Kim J-S: **Targeted genome engineering in human cells with the Cas9 RNA-guided endonuclease.** *Nat Biotechnol* 2013, **31**:230-232.
90. Mali P, Yang LH, Esvelt KM, Aach J, Guell M, DiCarlo JE, Norville JE, Church GM: **RNA-guided human genome engineering via Cas9.** *Science* 2013, **339**:823-826.
91. Hwang WY, Fu Y, Reyon D, Maeder ML, Tsai SQ, Sander JD, Peterson RT, Yeh JRJ, Joung JK: **Efficient genome editing in zebrafish using a CRISPR-Cas system.** *Nat Biotechnol* 2013, **31**:227-229.