



# Complete Genome Sequence of the Halophilic Methylophilic Methanogen Archaeon *Methanohalophilus portucalensis* Strain FDF-1<sup>T</sup>

Stéphane L'Haridon,<sup>a,b,c</sup> Erwan Corre,<sup>d</sup> Yue Guan,<sup>e</sup> Manikandan Vinu,<sup>e</sup> Violetta La Cono,<sup>f</sup> Michail Yakimov,<sup>f</sup> Ulrich Stingl,<sup>e,g</sup> Laurent Toffin,<sup>a,b,c</sup> Mohamed Jebbar<sup>a,b,c</sup>

<sup>a</sup>Université de Brest (UBO), Institut Universitaire Européen de la Mer (IUEM), UMR6197, Laboratoire de Microbiologie des Environnements Extrêmes (LM2E), Plouzané, France

<sup>b</sup>CNRS, IUEM-UMR 6197, Laboratoire de Microbiologie des Environnements Extrêmes (LM2E), Plouzané, France

<sup>c</sup>Ifremer, UMR6197, Laboratoire de Microbiologie des Environnements Extrêmes (LM2E), Plouzané, France

<sup>d</sup>CNRS, Université Pierre et Marie Curie, Station Biologique de Roscoff, Plateforme ABI/MS, Roscoff, France

<sup>e</sup>Red Sea Research Center, King Abdullah University of Science and Technology, Thuwal, Saudi Arabia

<sup>f</sup>Institute for Coastal Marine Environment CNR, Messina, Italy

<sup>g</sup>Department for Microbiology and Cell Sciences, University of Florida, UF/IFAS Fort Lauderdale Research and Education Center, Davie, Florida, USA

**ABSTRACT** We report here the complete genome sequence (2.08 Mb) of *Methanohalophilus portucalensis* strain FDF-1<sup>T</sup>, a halophilic methylophilic methanogen isolated from the sediment of a saltern in Figeria da Foz, Portugal. The average nucleotide identity and DNA-DNA hybridization analyses show that *Methanohalophilus mahii*, *M. halophilus*, and *M. portucalensis* are three different species within the *Methanosarcinaceae* family.

*Methanohalophilus portucalensis* strain FDF-1<sup>T</sup> (DSM 7471<sup>T</sup>, OCM 59) was described as a new species by Boone et al. (1). The strictly anaerobic species is able to produce methane by reducing methyl compounds and grows optimally at 40°C within a pH range of 6.5 to 7.5 and a salinity range of 0.5 to 2 M NaCl.

The complete genome of *M. portucalensis* was sequenced using a combination of the following three sequencing approaches: a 300-bp paired-end library sequenced on an Illumina MiSeq platform (Bioscience Core Lab, KAUST, Thuwal, Saudi Arabia), a 100-bp paired-end library sequenced on an Illumina HiSeq platform (Beckman Coulter Genomics, Inc., Danvers, MA, USA), and a PacBio RS library (Genotoul, Toulouse, France). The 13,835,892 paired reads of 300 bp were quality trimmed (Q30) and *de novo* assembled into contigs using SPAdes version 3.6.1 (2). The 17 resulting contigs were then scaffolded with SSPACE version 3.0 (3) using the 60,686,211 paired-end reads of 100 bp, leading to an intermediate version of the assemblage containing 9 scaffolds. A final scaffolding step was performed with SSPACE LongRead version 1.1 (4) using the 317,258 PacBio RS-filtered subreads. We finally obtained 3 scaffolds with a total size of 2,084,275 bp (without an unspecified base) and an average coverage of approximately 5,300×. Finally, a fully circularized version of the genome with expected gaps between oriented scaffolds was produced after comparison with the *M. mahii* genome using CONTIGuator (5).

The *M. portucalensis* genome consists of a circular chromosome of 2,084,875 bp with a GC content of 41.95%. A total of 2,198 coding DNA sequences were identified with the MaGe platform (6, 7), as well as 2 6S-23S operons, 3 5S rRNAs, 46 tRNAs, and 4 miscellaneous RNAs.

The *M. portucalensis* genome size is close to that of *M. mahii* strain SLP<sup>T</sup> (2,012,424 bp) (8) and *M. halophilus* strain Z-7982 (2,022,959 bp) (9). *M. mahii*, *M. halo-*

Received 27 November 2017 Accepted 1 December 2017 Published 18 January 2018

**Citation** L'Haridon S, Corre E, Guan Y, Vinu M, La Cono V, Yakimov M, Stingl U, Toffin L, Jebbar M. 2018. Complete genome sequence of the halophilic methylophilic methanogen archaeon *Methanohalophilus portucalensis* strain FDF-1<sup>T</sup>. *Genome Announc* 6:e01482-17. <https://doi.org/10.1128/genomeA.01482-17>.

**Copyright** © 2018 L'Haridon et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Mohamed Jebbar, [mohamed.jebbar@univ-brest.fr](mailto:mohamed.jebbar@univ-brest.fr).

*philus*, and *M. portucalensis* are physiologically very similar, and their separation into three different species was previously based on DNA reassociation and electrophoretic analysis of whole-cell proteins and on the need to maintain the genus *Methanohalophilus* for taxonomic stability (1). The average nucleotide identity (ANI) scores are 92.59% (SD, 2.91) between *M. halophilus* and *M. portucalensis*, 92.59% (SD, 2.91) between *M. halophilus* and *M. mahii*, and 92.59% (SD, 2.91) between *M. mahii* and *M. portucalensis*. An ANI score below 95% has been defined for the delineation of a new species (10). Thus, the ANI score comparisons indicate that the three strains are on the boundary of the species delineation. Afterward, we calculated the *in silico* DNA-DNA hybridization (DDH) values using the genome-to-genome distance calculator GGDC2.1 (11), which indicated values of 44.8% between *M. halophilus* and *M. mahii*, 44.4% between *M. portucalensis* and *M. mahii*, and 50.50% between *M. halophilus* and *M. mahii*. The ANI and DDH values confirm that *M. mahii*, *M. halophilus*, and *M. portucalensis* represent three phylogenetically closely related species.

**Accession number(s).** This whole-genome sequence has been deposited at GenBank under the accession number [CP017881](https://ncbi.nlm.nih.gov/nucl/CP017881).

## ACKNOWLEDGMENTS

The research leading to these results received funding from the European Union Seventh Framework Programme (FP7/2007–2013), project MaCuMBA, under grant agreement 311975. The LABGeM (CEA/IG/Genoscope and CNRS UMR 8030) and the France Génomique National Infrastructure (funded as part of the Investissement d'Avenir program managed by the Agence Nationale pour la Recherche, contract ANR-10-INBS-09) are acknowledged for their support within the MicroScope annotation platform.

## REFERENCES

- Boone DR, Mathrani IM, Liu YT, Menaia J, Mah RA, Boone JE. 1993. Isolation and characterization of *Methanohalophilus portucalensis* sp. nov. and DNA reassociation study of the genus *Methanohalophilus*. *Int J Syst Bacteriol* 43:431–437.
- Nurk S, Bankevich A, Antipov D, Gurevich A, Korobeynikov A, Lapidus A, Pribelsky A, Pyshkin A, Sirotkin A, Sirotkin Y, Stepanauskas R, McLean J, Lasken R, Clingenpeel SR, Woyke T, Tesler G, Alekseyev MA, Pevzner PA. 2013. Assembling genomes and mini-metagenomes from highly chimeric reads, p 158–170. *In* Deng M, Jiang R, Sun F, Zhang X (ed), *Research in computational molecular biology. Proceedings of the 17th Annual International Conference, RECOMB 2013, Beijing, China, April 7–10, 2013*. Springer, Berlin, Germany.
- Boetzer M, Henkel CV, Jansen HJ, Butler D, Pirovano W. 2011. Scaffolding pre-assembled contigs using SSPACE. *Bioinformatics* 27:578–579. <https://doi.org/10.1093/bioinformatics/btq683>.
- Boetzer M, Pirovano W. 2014. SSPACE-LongRead: scaffolding bacterial draft genomes using long read sequence information. *BMC Bioinformatics* 15:211. <https://doi.org/10.1186/1471-2105-15-211>.
- Galardini M, Biondi EG, Bazzicalupo M, Mengoni A. 2011. CONTIGuator: a bacterial genomes finishing tool for structural insights on draft genomes. *Source Code Biol Med* 6:11. <https://doi.org/10.1186/1751-0473-6-11>.
- Vallenet D, Labarre L, Rouy Z, Barbe V, Bocs S, Cruveiller S, Lajus A, Pascal G, Scarpelli C, Médigue C. 2006. MaGe: a microbial genome annotation system supported by synteny results. *Nucleic Acids Res* 34:53–65. <https://doi.org/10.1093/nar/gkj406>.
- Vallenet D, Belda E, Calteau A, Cruveiller S, Engelen S, Lajus A, Le Fèvre F, Longin C, Mornico D, Roche D, Rouy Z, Salvignol G, Scarpelli C, Thil Smith AAT, Weiman M, Médigue C. 2013. MicroScope—an integrated microbial resource for the curation and comparative analysis of genomic and metabolic data. *Nucleic Acids Res* 41:D636–D647. <https://doi.org/10.1093/nar/gks1194>.
- Spring S, Scheuner C, Lapidus A, Lucas S, Del Rio TG, Tice H, Copeland A, Cheng JF, Chen F, Nolan M, Saunders E, Pitluck S, Liolios K, Ivanova N, Mavromatis K, Lykidis A, Pati A, Chen A, Palaniappan K, Land M, Hauser L, Chang YJ, Jeffries CD, Goodwin L, Detter JC, Brettin T, Rohde M, Goker M, Woyke T, Bristow J, Eisen JA, Markowitz V, Hugenholtz P, Kyrpides NC, Klenk HP. 2010. The genome sequence of *Methanohalophilus mahii* SLP<sup>T</sup> reveals differences in the energy metabolism among members of the *Methanosarcinaceae* inhabiting freshwater and saline environments. *Archaea* 2010:690737.
- L'Haridon S, Corre E, Guan Y, Vinu M, La Cono V, Yakimov M, Stingl U, Toffin L, Jebbar M. 2017. Complete genome sequence of *Methanohalophilus halophilus* DSM 3094<sup>T</sup>, isolated from a cyanobacterial mat and bottom deposits at Hamelin Pool, Shark Bay, northwestern Australia. *Genome Announc* 5(7):e01604-16. <https://doi.org/10.1128/genomeA.01604-16>.
- Goris J, Konstantinidis KT, Klappenbach JA, Coenye T, Vandamme P, Tiedje JM. 2007. DNA-DNA hybridization values and their relationship to whole-genome sequence similarities. *Int J Syst Evol Microbiol* 57:81–91. <https://doi.org/10.1099/ijs.0.64483-0>.
- Meier-Kolthoff JP, Auch AF, Klenk HP, Göker M. 2013. GBDP on the grid: a genome-based approach for species delimitation adjusted for an automated and highly parallel processing of large data sets, Hochleistungsrechnen in Baden-Württemberg—Ausgewählte Aktivitäten im bwGRID 2012. KIT Scientific Publishing, Karlsruhe, Germany.