



Draft Genome Sequences of Four *Salmonella enterica* subsp. *enterica* Serovar Enteritidis Strains Implicated in Infections of Avian and Human Hosts

Ran An,^a Pengpeng Lin,^b Salim Bougouffa,^c Magbubah Essack,^c David Boxrud,^d Vladimir B. Bajic,^c Sinisa Vidovic^a

^aDepartment of Veterinary and Biomedical Sciences, University of Minnesota, Saint Paul, Minnesota, USA

^bDepartment of Mathematics, Statistics, and Computer Sciences, University of Wisconsin—Stout, Menomonie, Wisconsin, USA

^cKing Abdullah University of Science and Technology (KAUST), Computational Biosciences Research Center, Thuwal, Kingdom of Saudi Arabia

^dPublic Health Laboratory, Minnesota Department of Health, Saint Paul, Minnesota, USA

ABSTRACT *Salmonella enterica* subsp. *enterica* serovar Enteritidis is a wide-host-range pathogen. Occasionally, it is involved in invasive infections, leading to a high mortality rate. Here, we present the draft genome sequences of four *S. Enteritidis* strains obtained from human and avian hosts that had been involved in bacteremia, gastroenteritis, and primary infections.

Salmonella enterica subsp. *enterica* serovar Enteritidis (here referred to as *Salmonella* Enteritidis) is a Gram-negative rod-shaped non-spore-forming bacterium capable of infecting a wide range of mammalian (1), bird (2), and reptile (3) species.

The *S. Enteritidis* strains 15-00488 and 16-03217, implicated in the infection of avian hosts, were obtained from Alabama and Wisconsin in 2015 and 2016, respectively. Two human *S. Enteritidis* strains, 2016010175 and 2011007870, were obtained from Minnesota in 2015 and 2011, respectively. The strain 2016010175 was isolated from a stool sample of a patient who suffered from gastrointestinal infection, and the 2011007870 strain was obtained from a patient who experienced bacteremia. The primary goal of this research project was to identify differences on the genome level between avian-associated (15-00488 and 16-03217) and human-associated (2016010175 and 2011007870) *S. Enteritidis* strains.

Genomic DNA was extracted using the Genra Puregene kit (Qiagen, Inc., Valencia, CA). The whole-genome sequencing libraries were prepared using the Nextera XT DNA library preparation kit (Illumina, San Diego, CA, USA). The genomes were sequenced using an Illumina HiSeq 2500 platform with V4 125-bp paired-end sequencing chemistry. Sequencing reads were *de novo* assembled by the A5 pipeline version 5.0 (4). The draft genomes were annotated using the NCBI Prokaryotic Genome Automated Annotation Pipeline (https://www.ncbi.nlm.nih.gov/genome/annotation_prok/). Plasmid-Finder 1.3 (5) and Mauve (6) were used for plasmid and single nucleotide polymorphism (SNP) identification, respectively.

The draft genomes of the avian strains, 15-00488 and 16-03217, were 4,699,188 bp and 4,771,079 bp, with average genome coverages of 60× and 183×, respectively. The genomes of the 15-00488 and 16-03217 strains resulted in 4,736 and 4,851 protein-coding sequences, respectively. The two draft genomes of the human strains, 2016010175 and 2011007870, resulted in 4,771,079 bp and 4,698,250 bp, with average genome coverages of 75× and 146×, respectively. The 2016010175 and 2011007870 strains contained 4,737 and 4,740 protein-coding regions, respectively.

Received 12 December 2017 Accepted 19 December 2017 Published 25 January 2018

Citation An R, Lin P, Bougouffa S, Essack M, Boxrud D, Bajic VB, Vidovic S. 2018. Draft genome sequences of four *Salmonella enterica* subsp. *enterica* serovar Enteritidis strains implicated in infections of avian and human hosts. *Genome Announc* 6:e01550-17. <https://doi.org/10.1128/genomeA.01550-17>.

Copyright © 2018 An et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Sinisa Vidovic, svidovic@umn.edu.

Comparative genome analysis revealed that the *S. Enteritidis* strains of avian origin possessed all four genes, *galK*, *galM*, *galT*, and *galE*, that encode the enzymes galactokinase, galactose-1-epimerase, galactose-1-phosphate uridylyltransferase, and UDP-galactose 4-epimerase, respectively, which are necessary for conversion of galactose into glucose. Besides the enzymes of the Leloir pathway, the avian strains possessed the protein-coding sequences involved in acquisitions of molybdate (*modFBAE*) and iron (*fepCD*), as well as multidrug efflux (*ybhT*), and a protein of unknown function (DUF2167 domain-containing protein). Recently, it has been shown that molybdate uptake is important for anaerobic growth and multihost virulence in *Pseudomonas aeruginosa* (7, 8), which may indicate that the human-associated *S. Enteritidis* strains undergo a host restriction process compared to that of the avian strains.

In addition to loss of the genes involved in conversion of galactose into glucose and acquisition of molybdate, the human-associated strains showed a unique host-associated amino acid substitution at position 111, threonine (T) to isoleucine (I), in the gene *lpfA*, which encodes an adhesin “long polar fimbria protein A” that mediates bacterial attachment to the Peyer’s patches (9).

Accession number(s). The complete genome sequences of avian-associated (15-00488 and 16-03217) and human-associated (2016010175 and 2011007870) strains have been deposited in the NCBI GenBank database under the accession numbers [NQNX00000000](#), [NQUZ00000000](#), [NQVA00000000](#), and [NQVB00000000](#), respectively.

ACKNOWLEDGMENTS

This work was supported by the start-up fund provided to S.V. by the Department of Veterinary and Biomedical Sciences, University of Minnesota. S.B., M.E., and V.B.B. were supported by the KAUST research grants URF/1/2302-01-01, FCC/1/1976-02-01, and BAS/1/1606-01-01, respectively.

REFERENCES

1. Kingsley RA, Msefula CL, Thomson NR, Kariuki S, Holt KE, Gordon MA, Harris D, Clarke L, Whitehead S, Sangal V, Marsh K, Achtman M, Molyneux ME, Cormican M, Parkhill J, MacLennan CA, Heyderman RS, Dougan G. 2009. Epidemic multiple drug resistant *Salmonella* Typhimurium causing invasive disease in sub-Saharan Africa have a distinct genotype. *Genome Res* 19:2279–2287. <https://doi.org/10.1101/gr.091017.109>.
2. Han J, Gokulan K, Barnette D, Khare S, Rooney AW, Deck J, Nayak R, Stefanova R, Hart ME, Foley SL. 2013. Evaluation of virulence and antimicrobial resistance in *Salmonella enterica* serovar Enteritidis isolates from humans and chicken- and egg-associated sources. *Foodborne Pathog Dis* 10:1008–1015. <https://doi.org/10.1089/fpd.2013.1518>.
3. Bosch S, Tauxe RV, Behravesh CB. 2016. Turtle-associated salmonellosis, United States, 2006–2014. *Emerg Infect Dis* 22:1149–1155. <https://doi.org/10.3201/eid2207.150685>.
4. Coil D, Jospin G, Darling AE. 2015. A5-miseq: an updated pipeline to assemble microbial genomes from Illumina MiSeq data. *Bioinformatics* 31:587–589. <https://doi.org/10.1093/bioinformatics/btu661>.
5. Carattoli A, Zankari E, García-Fernández A, Voldby Larsen M, Lund O, Villa L, Møller Aarestrup F, Hasman H. 2014. *In silico* detection and typing of plasmids using PlasmidFinder and plasmid multilocus sequence typing. *Antimicrob Agents Chemother* 58:3895–3903. <https://doi.org/10.1128/AAC.02412-14>.
6. Darling AE, Mau B, Perna NT. 2010. progressiveMauve: multiple genome alignment with gene gain, loss and rearrangement. *PLoS One* 5:e11147. <https://doi.org/10.1371/journal.pone.0011147>.
7. Pederick VG, Eijkelkamp BA, Ween MP, Begg SL, Paton JC, McDevitt CA. 2014. Acquisition and role of molybdate in *Pseudomonas aeruginosa*. *Appl Environ Microbiol* 80:6843–6852. <https://doi.org/10.1128/AEM.02465-14>.
8. Périnet S, Jeukens J, Kukavica-Ibrulj I, Ouellet MM, Charette SJ, Levesque RC. 2016. Molybdate transporter ModABC is important for *Pseudomonas aeruginosa* chronic lung infection. *BMC Res Notes* 9:23. <https://doi.org/10.1186/s13104-016-1840-x>.
9. Bäumler AJ, Tsolis RM, Heffron F. 1996. The *l.p.f.* fimbrial operon mediates adhesion of *Salmonella* Typhimurium to murine Peyer’s patches. *Proc Natl Acad Sci U S A* 93:279–283. <https://doi.org/10.1073/pnas.93.1.279>.