

Nonseparable, Space-Time Covariance Functions with Dynamical Compact Supports

Emilio Porcu, Moreno Bevilacqua, and Marc G. Genton

Abstract: The paper provides new classes of nonseparable space-time covariance functions with spatial (or temporal) margin belonging to the Generalized Wendland class of compactly supported covariance functions. An interesting feature of our covariances, from the computational viewpoint, is that the compact support is a decreasing function of the temporal (spatial) lag. We provide conditions for the validity of the proposed class, and analyze the problem of differentiability at the origin for the temporal (spatial) margin. A simulation study explores the finite sample properties and the computational burden associated with the maximum likelihood estimation of the covariance parameters. Finally, we use the proposed covariance models

School of Mathematics, Statistics, and Physics, Newcastle University,
Newcastle upon Tyne, NE1 7RU, United Kingdom.
E-mail: georgepolya01@gmail.com & emilio.porcu@newcastle.ac.uk

Universidad de Valparaiso, Department of Statistics, 2360102 Valparaiso, Chile.
E-mail: moreno.bevilacqua@uv.cl

Statistics Program, King Abdullah University of Science and Technology, Thuwal 23955-6900, Saudi Arabia. E-mail: marc.genton@kaust.edu.sa

on Irish wind speed data and compare them with Gneiting-Matérn models in terms of fitting, prediction efficiency and computational burden. Necessary and sufficient conditions together with other results on dynamically varying compact supports are provided in the Online Supplement to this paper.

Key words and phrases: Generalized Wendland covariance function; Geostatistics; Kriging; Random field; Sparse matrices.

1. Introduction

There has been an increasing interest for space-time modeling through covariance functions in the last decades; we refer the reader to Gneiting (2002a), Stein (2005), Zastavnyi and Porcu (2011), Gneiting *et al.* (2007) and Schlather (2010) as examples of this interest. Typically, data observed over space and time are frequently modeled as a realization of a stationary Gaussian random field having a covariance function that is spatially isotropic and temporally symmetric (Gneiting, 2002a). Specifically, for a stationary random field $Z(\mathbf{x}, t)$, with \mathbf{x} a point of \mathbb{R}^d and t denoting time, spatial isotropy is coupled with temporal symmetry through a continuous function, $\phi : [0, \infty) \times [0, \infty) \rightarrow \mathbb{R}$ such that

$$\text{cov} \{Z(\mathbf{x}, t), Z(\mathbf{x} + \mathbf{h}, t + u)\} = \phi(\|\mathbf{h}\|, |u|), \quad (1.1)$$

with $(\mathbf{h}, u) \in \mathbb{R}^d \times \mathbb{R}$ being the space-time lag vector and with $\phi(0, 0) = \sigma^2$ denoting the variance of Z . For the remainder of the paper, we use r for $\|\mathbf{h}\|$ and we use the abuse of notation u for $|u|$. Also, the margins $\phi(r, 0)$ and $\phi(0, u)$ are called, respectively, spatial and temporal margins. A covariance function ϕ is called separable if ϕ factors into the product of a purely spatial with a purely temporal covariance function.

A popular example of nonseparable space-time covariance functions of the type (1.1) is the Gneiting class (Gneiting, 2002a; Zastavnyi and Porcu, 2011). We define it here as

$$\phi(r, u) = \frac{\sigma^2}{\psi(u^2)^{d/2}} g\left(\frac{r^2}{\psi(u^2)}\right), \quad (r, u) \in [0, \infty) \times [0, \infty), \quad (1.2)$$

where g is completely monotonic on the positive real line, so that it is infinitely differentiable on $(0, \infty)$ and $(-1)^k g^{(k)}(t) \geq 0$ for $t \geq 0$. The function ψ is strictly positive and has a completely monotonic derivative. Additionally, with no loss of generality, we assume that $g(0) = \psi(0) = 1$ so that $\phi(0, 0) = \sigma^2$. Sufficient conditions for the validity of this class have been found by Gneiting (2002a). Then Zastavnyi and Porcu (2011) found the necessary conditions and additionally relaxed the hypothesis on the function ψ . A subclass of the Gneiting class in (1.2) that became especially popular has expression

$$\phi(r, u) = \frac{\sigma^2}{\psi(u^2)^{d/2}} \mathcal{M}_\mu\left(\frac{r}{\psi(u^2)}\right), \quad (r, u) \in [0, \infty) \times [0, \infty), \quad (1.3)$$

where

$$\mathcal{M}_\mu(r) = \frac{2^{1-\mu}}{\Gamma(\mu)} r^\mu \mathcal{K}_\mu(r), \quad r \geq 0,$$

with $\mu > 0$ and \mathcal{K}_μ a modified Bessel function of the second kind of order μ , is the so called Matérn class (Stein, 1999). Hence, the class in Equation (1.3) has been termed Gneiting-Matérn class. The parameter μ characterizes the differentiability at the origin and, as a consequence, the differentiability of the sample paths of a Gaussian field in \mathbb{R}^d with Matérn covariance function. In particular for a positive integer k , the sample paths are k times differentiable, in any direction, if and only if $\mu > k$.

A spatial covariance function is called compactly supported if it vanishes after a given spatial distance. There is a large literature about compactly supported covariance functions in many branches of probability theory, geostatistics and approximation theory, and the reader is referred to Golubov (1981), Wendland (1995), Schaback and Wu (1995), Wu (1995), Buhmann (2000), Gneiting (2002b), Zastavnyi and Trigub (2002), Zastavnyi (2006), Schaback (2011), Zhu (2012), Hubbert (2012), Porcu and Zastavnyi (2014), Chernih *et al.* (2014) as well as to the more recent results in Bevilacqua *et al.* (2018) and the review by Porcu *et al.* (2018).

Compactly supported covariance functions are relevant to various ends: computationally efficient spatial prediction (Furrer *et al.*, 2006, with the references

therein) and estimation (Kaufman *et al.*, 2008) in the covariance tapering technique, fast and exact simulation, and appeal among practitioners (Gneiting, 2002b). The recent work by Bevilacqua *et al.* (2018) shed some light on their importance for kriging predictions, since it was shown that the Generalized Wendland class (Zastavnyi and Trigub, 2002; Gneiting, 2002a) is compatible with the Matérn class. This implies that, under fixed domain asymptotics, and under some specific conditions on the parameters indexing the covariance functions, the misspecified linear unbiased predictor with the Generalized Wendland class is asymptotically as efficient as the true simple kriging predictor using a Matérn class. Thus, kriging prediction can be performed with a compactly supported function without any loss of asymptotic prediction efficiency.

The problem of construction of nonseparable compactly supported space-time covariance functions is almost unexplored. A mathematical formulation of the problem was made in Zastavnyi and Porcu (2011), who suggested to replace the function g in Equation (1.2) with another function having compact support. They could not find any solution to such problem, a characterization of which remains elusive. The present paper challenges this problem. Specifically, we show how to generate covariance functions of the type (1.2) by replacing the function g with another function with compact support. Further, we replace the Matérn function used in the Gneiting-Matérn class in (1.3) with generalized Wendland

functions that are compactly supported, and that have the same properties of the Matérn class in terms of differentiability at the origin (Bevilacqua *et al.*, 2018).

A simulation study explores the finite sample properties of the maximum likelihood (ML) estimation of the covariance parameters. Finally, we apply our models on Irish wind speed data and compare them with Gneiting models in terms of fitting, prediction efficiency through some predictive scores and computational burden.

The plan of the paper is the following. Section 2.1 contains the necessary background and introduces the Generalized Wendland class. Section 2.2 provides the results on the new classes of space-time covariance functions proposed in this paper. Section 2.3 discusses examples and parameterization. Section 2.4 provides conditions to improve the differentiability of the temporal margins of the proposed classes. Section 3 explores our findings through both simulation and real data. Section 4 concludes the paper.

In the Online Supplement (OS throughout) we provide a collection of more technical results: on the one hand, we generalize the results in Section 2 to wider classes of functions with compact support. On the other hand, Fourier analysis and completely monotone functions are used to explore necessary and sufficient conditions. In the OS we also provide some figures that are discussed

in the paper.

2. Compactly Supported Space-Time Covariance Functions

2.1 Background Material

To favor a neater exposition, some preliminaries are needed. Space-time covariance functions as in Equation (1.1) are positive semidefinite. That is, for any finite collection $\{(\mathbf{x}_k, t_k)\}_{k=1}^N \subset \mathbb{R}^d \times \mathbb{R}$ and for any system of constants $\{c_k\}_{k=1}^N \subset \mathbb{R}$, we have

$$\sum_{k=1}^N \sum_{h=1}^N c_k c_h \phi(\|\mathbf{x}_k - \mathbf{x}_h\|, |t_k - t_h|) \geq 0.$$

In what follows we propose a class of candidate functions with compact support that can be used to replace the function g in Equation (1.2) while preserving positive definiteness. We introduce the Generalized Wendland class (Gneiting, 2002b; Zastavnyi and Trigub, 2002) $\varphi_{\nu, \kappa} : [0, \infty) \rightarrow \mathbb{R}$, defined through

$$\varphi_{\nu, \kappa}(r) = \frac{1}{B(2\kappa + 1, \nu)} \int_r^\infty (t^2 - r^2)^\kappa \varphi_{\nu-1, 0}(t) dt \quad r \geq 0, \quad (2.1)$$

where $\kappa > 0$ and with B denoting the beta function, that is

$$B(2\kappa + 1, \nu) = \frac{\Gamma(2\kappa + 1)\Gamma(\nu)}{\Gamma(2\kappa + \nu + 1)}.$$

Here, $\varphi_{\nu, 0}$ denotes the Askey family of functions (Askey, 1973), defined by

$$\varphi_{\nu, 0}(r) := (1 - r)_+^\nu, \quad \nu > 0, \quad (2.2)$$

where $(\cdot)_+$ denotes positive part. Let d be a positive integer. The function $\varphi_{\nu,0}(r)$ is positive definite in \mathbb{R}^d if and only if $\nu \geq (d+1)/2$ (Golubov, 1981). Arguments in Zastavnyi and Trigub (2002) show that $\varphi_{\nu,\kappa}$ is positive definite in \mathbb{R}^d if and only if $\nu \geq (d+1)/2 + \kappa$. Additionally, $\varphi_{\nu,\kappa}(\cdot/b)$ is compactly supported over the ball of \mathbb{R}^d with radius $b > 0$. Closed form solutions of the integral in Equation (2.1) can be obtained when $\kappa = k$, a non negative integer. In this case,

$$\varphi_{\nu,k}(r) = \varphi_{\nu+k,0}(r)P_k(r), \quad r \geq 0,$$

with P_k a polynomial of order k . See the first column of Table 1 for some examples with $k = 0, 1, 2, 3$. These functions, termed (original) Wendland functions, were originally proposed by Wendland (1995). Other closed form solutions of the integral (2.1) can be obtained when $\kappa = k + 1/2$, using some results in Schaback (2011). Hubbert (2012) showed some other closed forms based on hypergeometric functions. Finally, Chernih *et al.* (2014) showed that, for κ tending to infinity, a rescaled version of the model (2.1) converges to a Gaussian model. As noted by Gneiting (2002b), the Generalized Wendland and Matérn models exhibit the same behavior at the origin when the smoothness parameters of the two covariance models are related by the equation $\nu = \kappa + 1/2$. This fact is depicted by Table 1, where some specific cases of Wendland functions are compared with the Matérn covariance in terms of sample paths differentiability of

2.1 Background Material

Table 1: Generalized Wendland correlation $\varphi_{\nu,\kappa}(r)$ and Matérn correlation $\mathcal{M}_\mu(r)$ with increasing smoothness parameters κ and μ . $SP(k)$ means that the sample paths of the associated Gaussian field are k times differentiable. Taken from Bevilacqua *et al.* (2018).

κ	$\varphi_{\nu,\kappa}(r)$	μ	$\mathcal{M}_\mu(r)$	$SP(k)$
0	$(1-r)_+^\nu$	0.5	e^{-r}	0
1	$(1-r)_+^{\nu+1}(1+r(\nu+1))$	1.5	$e^{-r}(1+r)$	1
2	$(1-r)_+^{\nu+2}(1+r(\nu+2)+r^2(\nu^2+4\nu+3)\frac{1}{3})$	2.5	$e^{-r}(1+r+\frac{r^2}{3})$	2
3	$(1-r)_+^{\mu+3}(1+r(\nu+3)+r^2(2\nu^2+12\nu+15)\frac{1}{5}+r^3(\nu^3+9\nu^2+23\nu+15)\frac{1}{15})$	3.5	$e^{-r}(1+\frac{r}{2}+r^2\frac{6}{15}+\frac{r^3}{15})$	3

the associated Gaussian random field. Generalized Wendland functions include many other popular classes of covariance functions with compact support, and for a recent review the reader is referred to Porcu *et al.* (2018).

We finish this section with a new definition that opens to the results provided in the subsequent section. Let ϕ be a space-time covariance functions as in Equation (1.1). We call a temporally dynamical radius, ψ , the continuous mapping from $[0, \infty)$ into $(0, \infty)$ such that, for each $u_o \in [0, \infty)$, the margin $\phi(\cdot, u_o)$ is compactly supported on the interval $[0, \psi(u_o))$. Clearly, both Askey and Generalized Wendland classes are special cases of dynamical compact support, when $\psi \equiv b > 0$ is the constant function.

2.2 Space-time Gneiting-Wendland functions with dynamical compact support

The results coming subsequently are based on a constructive criterion provided by Porcu and Zastavnyi (2012).

Lemma 1. Let d be a positive integer. Let (Ω, \mathcal{F}, P) be a measure space with P a positive measure. Let $H(\cdot; \cdot) : \Omega \times [0, \infty) \rightarrow \mathbb{R}$ and $F(\cdot; \cdot) : [0, \infty) \times \Omega \rightarrow \mathbb{R}$ such that

1. $H(\xi; \cdot)$ is a temporal covariance function for all $\xi \in \Omega$;
2. $F(\cdot; \xi)$ is an isotropic spatial covariance function in \mathbb{R}^d for all $\xi \in \Omega$;
3. $H(\cdot; u)F(r; \cdot) \in L_1(\Omega, \mathcal{F}, P)$ for any $r, u \geq 0$.

Then, the mapping

$$\phi(r, u) = \sigma^2 \frac{\int_{\Omega} F(r; \xi) H(\xi; u) P(d\xi)}{\int_{\Omega} F(0; \xi) H(\xi; 0) P(d\xi)}, \quad (r, u) \in [0, \infty) \times [0, \infty), \quad (2.3)$$

with $\sigma^2 > 0$, defines a space-time covariance function in $\mathbb{R}^d \times \mathbb{R}$ that is isotropic in the spatial argument and symmetric in time.

An intuitive way to understand the formal statement in Lemma 1 is to see the integral in Equation (2.3) as a scale mixture of a spatial and a temporal covariance. Conditions 1 and 2 are needed to have well defined spatial and temporal covariances. Condition 3 ensures the integral (2.3) to be well defined.

2.2 Space-time Gneiting-Wendland functions with dynamical compact support

Another relevant comment is that the denominator in Equation (2.3) is a normalization constant, so that $\phi(0, 0) = \sigma^2$.

Theorem 2.1. *Let d be a positive integer. Let $\varphi_{\nu,0}$ be the Askey function in (2.2). Let ψ be a continuous and positive function on the positive real line, with $\psi(0) = 1$ and such that $1/\psi$ is increasing and concave on the positive real line, with $\lim_{t \rightarrow \infty} \psi(t) = 0$. Then, the mapping*

$$\phi(r, u) = \psi(u)^\alpha \varphi_{\nu,0} \left(\frac{r}{\psi(u)} \right), \quad (r, u) \in [0, \infty) \times [0, \infty), \quad (2.4)$$

defines a space-time covariance function in $\mathbb{R}^d \times \mathbb{R}$ provided $\nu \geq (d + 5)/2$ and $\alpha \geq (d + 3)/2$.

Proof. The proof is an application of Lemma 1. Specifically, we use the scale mixture argument of Equation (2.3) under some appropriate choices of the functions H and F .

We now proceed formally and consider the mapping $F(r; \xi) = \varphi_{n,0}(r/\xi)$. Arguments in Golubov (1981) show that $F(\cdot; \xi)$ is a isotropic spatial covariance function in \mathbb{R}^d for any $\xi > 0$ provided $n \geq (d + 1)/2$. Thus, Condition 2 in Lemma 1 is satisfied. As for the choice of the function H , we consider the mapping

$$H(\xi; u) = H_{n,\gamma}(\xi; u) = \xi^n \varphi_{\gamma,0} \left(1 - \frac{\xi}{\psi(u)} \right)_+, \quad \xi > 0, u \geq 0, \quad \gamma \geq 1, n > 0.$$

2.2 Space-time Gneiting-Wendland functions with dynamical compact support

In virtue of the properties of the function ψ , we have that $H_{n,\gamma}$ is positive, decreasing and convex, with $\lim_{t \rightarrow \infty} H_{n,\gamma}(\xi; t) = 0$ for any $\xi > 0$. Thus, we can invoke the Pólya criterion (Pólya, 1949) to show that $H_{n,\gamma}(\xi; u)$ is a covariance function in \mathbb{R} . This fact shows that Condition 1 in Lemma 1 is satisfied. Finally, observe that Condition 3 of Lemma 1 holds trivially. Thus, we can now apply the scale mixture in Equation (2.3) with $\Omega = [0, \infty)$ and P being the Lebesgue measure:

$$\begin{aligned}
 \phi(r, u) &= \int_{(0, \infty)} F(r; \xi) H(\xi; u) d\xi \\
 &= \int_{(0, \infty)} \left(1 - \frac{r}{\xi}\right)_+^n \xi^n \left(1 - \frac{\xi}{\psi(u)}\right)_+^\gamma d\xi \\
 &= \frac{1}{\psi(u)^\gamma} \int_r^{\psi(u)} (\xi - r)^n (\psi(u) - \xi)^\gamma d\xi \\
 &= \frac{1}{\psi(u)^\gamma} \int_0^{\psi(u)-r} t^n (\psi(u) - r - t)^\gamma dt \\
 &= \frac{1}{\psi(u)^\gamma} \int_0^1 (\psi(u) - r)^{n+\gamma+1} v^n (1 - v)^\gamma dv \\
 &= B(n+1, \gamma+1) \psi(u)^{n+1} \left(1 - \frac{r}{\psi(u)}\right)^{n+\gamma+1} \\
 &= B(n+1, \gamma+1) \psi(u)^{n+1} \varphi_{n+\gamma+1,0} \left(\frac{r}{\psi(u)}\right), \tag{2.5}
 \end{aligned}$$

with B denoting the beta function, and where the third line in the chain of equalities is justified by the fact that, by definition, ϕ is identically equal to zero whenever $r > \psi(u)$. We now let $\alpha = n + 1$ and $\nu = n + \gamma + 1$. Thus, Equations (2.4) and (2.5) agree modulo a positive factor, that is the normalization constant.

2.2 Space-time Gneiting-Wendland functions with dynamical compact support

This fact completes the proof. The conditions on α and ν are easily verified from the previous identities. □

Theorem 2.2. *Let d be a positive integer and $\kappa > 0$. Let $\varphi_{\nu,\kappa}$ be the Generalized Wendland class of functions in (2.1). Let ψ be a continuous and positive function on the positive real line, with $\psi(0) = 1$ and such that $1/\psi(\cdot)$ is increasing and concave on the positive real line, with $\lim_{t \rightarrow \infty} \psi(t) = 0$. Then, the mapping ϕ defined through*

$$\phi(r, u) = \psi(u)^\alpha \varphi_{\nu,\kappa} \left(\frac{r}{\psi(u)} \right), \quad (r, u) \in [0, \infty) \times [0, \infty), \quad (2.6)$$

defines a space-time covariance function in $\mathbb{R}^d \times \mathbb{R}$ provided that $\nu \geq (d+5)/2 + \kappa$ and $\alpha \geq (d+3)/2 + 2\kappa$.

Proof. We give a constructive proof by applying again Lemma 1 for some specific choices of the functions H and F in the scale mixture (2.3). As for the choice of the function F , let $\kappa > 0$, $n \geq (d+1)/2 + \kappa$ and $F(r; \xi) = \varphi_{n,\kappa}(r/\xi)$, with $\varphi_{n,\kappa}$ as defined in Equation (2.1). Clearly, Condition 1 in Lemma 1 is satisfied. Further, we observe that arguments in Zastavnyi and Trigub (2002) show that $\varphi_{n,\kappa}$ can be rewritten as:

$$\varphi_{n,\kappa}(w) = \frac{1}{B(n, 2\kappa + 1)} \int_w^1 (1-t)^{n-1} (t^2 - w^2)^\kappa dt, \quad w \geq 0.$$

2.2 Space-time Gneiting-Wendland functions with dynamical compact support

In particular, following Daley *et al.* (2015), we have that, for $0 < y < \xi \leq 1$,

$$\begin{aligned}\varphi_{n,\kappa}\left(\frac{y}{\xi}\right) &= \frac{1}{B(n, 2\kappa + 1)} \int_{y/\xi}^1 (1-t)^{n-1} \left(t^2 - \frac{y^2}{\xi^2}\right)^\kappa dt \\ &= \frac{1}{B(n, 2\kappa + 1)} \int_y^\xi \left(1 - \frac{v}{\xi}\right)^{n-1} (v^2 - y^2)^\kappa \frac{dv}{\xi^{2\kappa+1}}.\end{aligned}$$

We now choose the function

$$H(\xi; u) = H_{n,\kappa,\gamma}(\xi; u) = \xi^{n+2\kappa} \left(1 - \frac{\xi}{\psi(u)}\right)_+^\gamma, \quad \xi > 0, u \geq 0, \quad \gamma \geq 1, n > 0,$$

with κ positive and ψ as stated. Again, it is easy to show that $H_{n,\kappa,\gamma}(\xi; \cdot)$ is positive, decreasing and convex with $\lim_{t \rightarrow \infty} H_{n,\kappa,\gamma}(\xi; t) = 0$ for all $\xi > 0$. Thus, Condition 2 of Lemma 1 is satisfied. Condition 3 holds trivially. We can thus apply the scale mixture argument in Equation (2.3) with $\Omega = [0, \infty)$ and P being the Lebesgue measure. We write ψ for $\psi(u)$ and have

$$\begin{aligned}& \int_0^\infty \varphi_{n,\kappa}\left(\frac{r}{\xi}\right) H_{n,\kappa,\gamma}(\xi; u) d\xi = \int_r^\psi \varphi_{n,\kappa}\left(\frac{r}{\xi}\right) \xi^{n+2\kappa} \left(1 - \frac{\xi}{\psi}\right)^\gamma d\xi \\ &= \frac{1}{B(n, 2\kappa + 1)} \int_r^\psi \xi^{n+2\kappa} \left(1 - \frac{\xi}{\psi}\right)^\gamma d\xi \int_r^\xi \left(1 - \frac{v}{\xi}\right)^{n-1} (v^2 - r^2)^\kappa \frac{dv}{\xi^{2\kappa+1}} \\ &= \frac{\psi^{-\gamma}}{B(n, 2\kappa + 1)} \int_r^\psi (v^2 - r^2)^\kappa dv \int_v^\psi (\psi - \xi)^\gamma (\xi - v)^{n-1} d\xi \\ &= \frac{\psi^{-\gamma}}{B(n, 2\kappa + 1)} B(n, \gamma + 1) \int_r^\psi (\psi - v)^{n+\gamma} (v^2 - r^2)^\kappa dv \\ &= \psi^{n+2\kappa+1} \frac{B(n, \gamma + 1)}{B(n, 2\kappa + 1)} \int_{r/\psi}^1 (1-t)^{n+\gamma} \left(t^2 - \frac{r^2}{\psi^2}\right)^\kappa dt \\ &= \frac{B(n, \gamma + 1)}{B(n, 2\kappa + 1)} B(n + \gamma + 1, 2\kappa + 1) \psi^{n+2\kappa+1} \varphi_{n+\gamma+1,\kappa}\left(\frac{r}{\psi}\right) \\ &= B(n + 2\kappa + 1, \gamma + 1) \psi^{n+2\kappa+1} \varphi_{n+\gamma+1,\kappa}\left(\frac{r}{\psi}\right),\end{aligned}$$

2.3 Examples and Parameterizations

with B being the beta function as before. We now let $\nu = n + \gamma + 1$ and $\alpha = n + 2\kappa + 1$. Rescaling at the origin and using the same arguments as in Theorem 2.1, we easily arrive at the assertion. \square

Using the arguments in Gneiting (2002b), it can be shown that for any increasing sequence $\{c_n\}_{n \geq 0}$ we have $\varphi_{c_n, \kappa} \left\{ \frac{r}{\psi(u)c_n} \right\} \rightarrow \mathcal{M}_{1/2+\kappa} \{r/\psi(u)\}$, with the convergence being uniform on any bounded set. Thus, the class in Equation (2.6) converges to the Gneiting-Matérn class and when $u = 0$ the smoothness parameters of the two covariance models are related by the equation $\mu = \kappa + 1/2$ (see again Table 1 for an illustration).

2.3 Examples and Parameterizations

Several examples from the mappings ψ that satisfy the requirements in Theorems 2.1 and 2.2 can be found in Table 1 in Porcu and Schilling (2011). A notable example comes from the choice

$$\psi(t; \delta, \beta) = (1 + t^\delta)^{-\beta/\delta}, \quad t \geq 0, \quad (2.7)$$

for $0 < \delta \leq 1$ and $0 \leq \beta \leq \delta$. In particular, in the following sections, we work with the special case $\psi(\cdot; \beta) := \psi(\cdot; 1, \beta)$, valid for $\beta \in [0, 1]$.

For $\kappa = k$, a nonnegative integer, we find that the classes in (2.6) can be

written as

$$\phi(r, u) = \psi(u)^\alpha \varphi_{\nu+k,0} \left(\frac{r}{\psi(u)} \right) P_k \left(\frac{r}{\psi(u)} \right), \quad r, u \geq 0,$$

where the constraints on α and ν are specified in Theorems 2.1 and 2.2, with P_k being a polynomial of degree k . In particular, we make use of the cases $k = 0, 1, 2$ for ease of illustration. Using the first three entries in Table 1 coupled with Equations (2.4) and (2.6), we obtain

$$\begin{aligned} \phi(r, u) &= \psi(u)^\alpha \left(1 - \frac{r}{\psi(u)} \right)_+^\nu, \\ \phi(r, u) &= \psi(u)^\alpha \left(1 - \frac{r}{\psi(u)} \right)_+^{\nu+1} \left(1 + (\nu+1) \frac{r}{\psi(u)} \right), \\ \phi(r, u) &= \psi(u)^\alpha \left(1 - \frac{r}{\psi(u)} \right)_+^{\nu+2} \left(1 + (\nu+2) \frac{r}{\psi(u)} + \frac{1}{3} \left((\nu+2)^2 - 1 \right) \left(\frac{r}{\psi(u)} \right)^2 \right), \end{aligned} \tag{2.8}$$

where α and ν must be determined according to Theorems 2.1 (for the first example) and 2.2 (for the other two examples). For geostatistical applications, it is useful to consider rescaled versions $\phi(r/b, u/a)$, which in turn allow to consider the marginal spatial compact support $b > 0$, the dynamical compact support $b\psi(u/a)$, and the temporal scale parameter $a > 0$. In many instances, a reparameterization of the proposed covariance models is useful. For instance, using the function (2.7) in the construction (2.8), and replacing $\beta\alpha$ with $\tau > 0$, we obtain the space-time correlation functions:

$$\phi(r, u) = \frac{1}{(1 + u/a)^\tau} \varphi_{\nu,k} \left(\frac{r}{b(1 + u/a)^\beta} \right), \quad r, u \geq 0, \tag{2.9}$$

with $\tau \geq 2.5 + 2k$ and $k = 0, 1, 2$. If we fix τ , we obtain a parametric family with an easily interpretable space-time nonseparable parameter $0 \leq \beta \leq 1$ which includes as special case a separable covariance, obtained when $\beta = 0$.

Figure S1 (OS) shows the contour-plot of the nonseparable space-time correlation functions with dynamical compact support in Equation (2.9). Specifically, we fix $b = 0.15$, $a = 0.2$, $\nu = 3.5 + \kappa$ and $\tau = 2.5 + 2\kappa$ for $\kappa = 0, 1$ and $\beta = 0, 0.5, 1$. Apparently, when increasing β , the rate of decay of the dynamical compact support is more severe. Thus, this parameter affects the dynamical compact support, that is, the sparseness of the associated correlation matrix.

Figure S2 (OS) shows a simulation on a regular grid of 12,544 location sites over the unit square and over temporal instants $u = 1, 1.5$, obtained through Cholesky decomposition, of a space-time Gaussian field with correlation (2.9) (top) fixing $\kappa = 1$, $\tau = 6.5$, $\nu = 4.5$, $b = 0.15$, $a = 0.2$ and $\beta = 0.5$. The same figure depicts a realization of a space-time Gaussian random field with covariance function from the Gneiting-Matérn class:

$$\phi(r, u) = \frac{1}{(1 + u/0.2)^{6.5}} \mathcal{M}_{1.5} \left(\frac{r}{0.0226(1 + u/0.2)^{1/4}} \right), \quad r, u \geq 0. \quad (2.10)$$

The two simulations share the same Gaussian realization in the Cholesky decomposition method. The two covariance models have the same marginal temporal correlation and the spatial scale parameter in the Gneiting-Matérn model is chosen such that the marginal spatial correlation is lower than 0.01 when $r > 0.15$,

i.e., greater than the marginal compact support of the Generalized Wendland model. It is apparent from Figure S2 that the two simulations look very similar.

Remark 2.1. The members of the classes in Theorems 2.1 and 2.2 are dynamically compactly supported in space. Thus, they are computationally suitable covariance models for space-time data with a relatively large number of location sites with respect to the temporal instants.

It is important to remark that the constructions in Theorems 2.1 and 2.2 can be interchanged, so to have space-time covariances being compactly supported over time, and with a compact support which evolves dynamically according to spatial distance. We omit such a specification of the mathematical conditions because the analogue specification is literal. Then, for instance, an analogue version of the model in Equation (2.9) is:

$$\phi(r, u) = \frac{1}{(1 + r/b)^\tau} \varphi_{\nu, \kappa} \left(\frac{u}{a(1 + r/b)^\beta} \right), \quad r, u \geq 0. \quad (2.11)$$

In this model the parameter a is the marginal temporal compact support and the decreasing dynamical compact support is given by $a\psi(r/b)$.

This kind of models is computationally more suitable for space-time data with a relatively large number of temporal instants with respect to the location sites as in the Irish wind speed data in Section 3.2.

2.4 Improving Temporal Differentiability at the Origin

The ψ functions that can be used for the purposes of Theorems 2.1 and 2.2 are, by construction, non-differentiable at the origin. This implies that we can govern the degree of differentiability in the spatial component, but not in the temporal one. This issue is studied in detail in the OS, where we show necessary conditions based on Fourier analysis that preserve positive definiteness of the constructions proposed in Theorems 2.1 and 2.2.

Having a model that allows for different degrees of temporal differentiability at the origin is important for attaining more flexibility in the analysis of space-time datasets. Another important fact is that differentiability at the origin has a crucial impact on spatial and temporal prediction (Stein, 1999). Since we are approximating the Gneiting-Matérn class with a compactly supported structure, it is important to be able to attain the same level of differentiability of both spatial and temporal margins.

Sufficient conditions that allow to improve the differentiability of the temporal margin can be improved on the basis of the following facts. The function

$$\varpi_{\tau,\lambda}(r) := \varphi_{\tau,0}(r^\lambda) = (1 - r^\lambda)_+^\tau, \quad r \geq 0, \quad \lambda \in (0, 2), \quad \tau > 0, \quad (2.12)$$

has attracted the interest of several mathematicians in the past, and we refer the reader to Gneiting (2001), with the references therein. In particular, the univari-

ate case, $d = 1$, has an interesting history and we again refer to Gneiting (2000) in his tour de force. We have that $\varpi_{\nu,2}$ is not positive definite on \mathbb{R} , regardless of the value of ν . Kuttner (1944) showed that there exists a function $\kappa_1(\lambda)$, $\lambda \in (0, 2)$, such that $\varpi_{\tau,\lambda}(r)$ is positive definite on \mathbb{R} if and only if $\tau > \kappa_1(\lambda)$. The function $\kappa_1(\lambda)$ is continuous and strictly increasing, with $\lim_{\lambda \rightarrow 0} \kappa_1(\lambda) > 0$, $\kappa_1(1) = 1$, $\lim_{\lambda \rightarrow 2} \kappa_1(\lambda) = \infty$, and $\kappa_1(\lambda) > \lambda$ if $\lambda \neq 1$.

We now apply our results to Theorem 2.2 and consider the function

$$\phi(r, u) = (1 + u^\lambda)^{-\alpha} \varphi_{\nu(\tau), \kappa} \left(\frac{r}{(1 + u^\lambda)} \right), \quad (r, u) \in [0, \infty) \times [0, \infty),$$

where ν is a function of τ as described through Equation (2.12). The same scale mixture arguments as in the proof of Theorem 2.2 apply (see Lemma 1), hence we omit them. We have that, for a given $d \in \mathbb{N}$, ϕ is positive definite on $\mathbb{R}^d \times \mathbb{R}$ provided $\alpha \geq (d + 3)/2$ and

$$\nu \geq (d + 3)/2 + \kappa + \tau, \quad \tau \geq \kappa_1(\lambda), \quad \lambda \in (0, 2).$$

Table 2.4, taken from Gneiting (2000), allows to obtain the corresponding values for a given $\lambda \in (0, 2)$.

3. Numerical Results

We start by describing the performance of the ML estimation of the parameters of the Gneiting-Wendland model. Then we compare the Gneiting-Matérn model

Table 2: Lower bounds for $\kappa_1(\lambda)$ for given values of λ . Taken from Gneiting (2000).

λ	1.05	1.15	1.25	1.45	1.55	1.75	1.95
$\kappa_1(\lambda)$	1.0507	1.1572	1.2706	1.5247	1.7234	2.3462	3.9084

with the proposed Gneiting-Wendland model from a modeling, prediction performance and computational point of view when used as space-time covariance models for the Irish wind speed data.

3.1 Simulation Studies

Following Remark 2.1, we consider two possible scenarios:

1. A dataset with many spatial location sites and few temporal observations. Specifically, we have \mathbf{x}_i , $i = 1, 2, \dots, 60$ location sites uniformly distributed on the unit square, and $u = 0, 0.25, \dots, 2.25$ temporal instants;
2. A dataset with few spatial location sites and many temporal observations, that is \mathbf{x}_i , $i = 1, 2, \dots, 10$ location sites uniformly distributed on the unit square and $u = 0, 0.25, \dots, 14.75$ temporal instant;

For both scenarios, the total number of observations is kept relatively small (600 observations) in order to make ML estimation feasible. Under Scenario 1,

we simulate 1000 zero mean space-time Gaussian random fields with covariance in Equation (2.9), setting $k = 0, 1, 2$ in order to consider different levels of differentiability in the spatial covariance margin. Then, following Theorems 2.1 and 2.2, we fix $\tau = 2.5 + 2\kappa$ and $\nu = 3.5 + \kappa$.

We set $\sigma^2 = 1$, $b = 0.15$, $a = 0.2$ and we fix $\beta = 0, 0.5, 1$. For each simulation we estimate with ML the parameters σ^2 , a and b . Table 3 reports bias and variance associated to ML estimation of σ^2 , a and b , for $k = 0.1, 2$ and $\beta = 0, 0.5, 1$.

Similarly, under the Scenario 2, we simulate 1000 zero mean space-time Gaussian random fields with covariance as in Equation (2.11), with $k = 0, 1, 2$, fixing $\tau = 2.5 + 2k$ and $\nu = 3.5 + k$. We set $\sigma^2 = 1$, $a = 0.75$, $b = 0.2$ and we consider $\beta = 0, 0.5, 1$. For each simulation we estimate with ML the parameters σ^2 , a and b . The spatial and temporal scales parameters in both scenarios have been chosen in order to attain a small dependence in space and time. Table 3 reports bias and standard deviation (SD) associated to ML estimation of σ^2 , a and b , for $k = 0.1, 2$ and $\beta = 0, 0.5, 1$. The bias is overall negligible, and increasing β does not affect the bias and the SDs of the ML estimation. Under Scenario 1 the SD of the spatial marginal compact support b increases considerably when increasing k . Similarly, under Scenario 2, the SD of the temporal marginal compact support a increases when increasing k .

The bottleneck when evaluating a Gaussian likelihood is the computation of the inverse and the determinant of the covariance matrix, and both can be easily obtained from its Cholesky decomposition. Some computational gains can be achieved in the Gaussian likelihood computation using specific algorithms for sparse matrices using our models. The sparsity of the covariance matrix changes at each iteration of the maximization algorithm. In our implementation, Gaussian likelihood optimization is performed exploiting algorithms for sparse matrices as implemented in the R package `spam` (Furrer and Sain, 2010) using the maximization algorithm proposed in Nelder and Mead (1965) and implemented in the `optim` function of the R software (R Development Core Team, 2016).

Substantial further computational gains are achieved when performing kriging prediction, since in this case the sparsity of the covariance matrix is fixed. More details are given in the next section.

3.2 Irish Wind Speed Data

The main goal of this section is to compare the Gneiting-Matérn model with the proposed Gneiting-Wendland model from a modeling, prediction performance and computational point of view when used as space-time covariance models for the Irish wind speed data (Haslett and Raftery, 1989).

We consider daily wind speeds collected over 18 years, from 1961 to 1978, at 12 sites in Ireland. Following Gneiting *et al.* (2007) we omit the Rosslare station, we consider the square root transformation of the data and we remove the seasonal component. The latter is estimated by calculating the average of the square roots of the daily means over all years and stations and regressing this on a set of annual harmonics. The resulting transformed data, $\{z(\mathbf{x}_i, t_j), i = 1, \dots, 11, j = 1, \dots, 6574\}$, are assumed to be a realization from a zero-mean space-time Gaussian random field. Since we perform ML estimation, we focus on a subset of the data for computational reasons. Specifically, we focus on $\mathbf{z} = \{z(\mathbf{x}_i, t_j), i = 1, \dots, 11, j = 366, \dots, 910\}$. Thus, we have $11 \times 545 = 5,995$ observations and ML estimation is still feasible.

From Figure S3 (OS) it becomes apparent that the empirical temporal marginal semivariogram attains the sill at a temporal distance of 3 days approximately. Thus, following Remark 2.1, a nonseparable, temporally compactly supported covariance model, as defined through Equation (2.11), seems to be a natural choice for this kind of data. We compare the following space-time covariance models. A Gneiting-Matérn model:

$$C_M(r, u; \boldsymbol{\theta}_M) = \frac{\sigma_M^2}{\psi(r/a_M)^{\tau_M}} \mathcal{M}_\mu \left(\frac{u}{b_M \psi(r/a_M)^{\beta_M/2}} \right), \quad \mu = 0.5, 1.5, 2.5, \quad (3.1)$$

and our Gneiting-Wendland model:

$$C_W(r, u; \boldsymbol{\theta}_W) = \frac{\sigma_W^2}{\psi(r/a_W)^{\tau_W}} \varphi_{\nu, k} \left(\frac{u}{b_W \psi(r/a_W)^{\beta_W}} \right), \quad k = 0, 1, 2, \quad (3.2)$$

where $\psi(r) = 1+r$, $r \geq 0$, and $\boldsymbol{\theta}_M = (\sigma_M^2, a_M, b_M, \beta_M)^\top$ and $\boldsymbol{\theta}_W = (\sigma_W^2, a_W, b_W)^\top$.

For the model in Equation (3.2), according to the choices $k = 0, 1, 2$, we fix $\tau_W = 2.5 + 2k$ and $\nu = 3.5 + k$ according to Theorem 2.2, so that positive definiteness is attained. Then, for each k , we deliberately choose β_W equal to 0, 0.5, 1 in order to increase the sparsity of the associated covariance matrix, and we estimate $\boldsymbol{\theta}_W$ using ML. Similarly, for model (3.1) we consider the cases $\mu = 0.5, 1.5, 2.5$ fixing $\tau_M = 2.5 + 2(\mu - 0.5)$ and we estimate $\boldsymbol{\theta}_M$ using ML.

This setting makes the models (3.1) and (3.2) comparable, because they share the same spatial margin. Besides, the temporal margins are of the Matérn and Generalized Wendland type respectively, with the same level of differentiability at the origin for $\mu = 0.5, 1.5, 2.5$ and $k = 0, 1, 2$ respectively.

Table 4 (top) reports the ML estimation of $\boldsymbol{\theta}_W$ for each $k = 0, 1, 2$ and for each $\beta_W = 0, 0.5, 1$ with associated loglikelihood, and Table 4 (bottom) reports the ML estimation of $\boldsymbol{\theta}_M$ for each $\mu = 0.5, 1.5, 2.5$ with associated loglikelihood.

A comparison of the two models in terms of loglikelihood shows that the best models are obtained when $k = 0$ and $\mu = 0.5$, that is when the temporal margin is not differentiable at the origin for both cases. For the Gneiting-Wendland model, the best fitting is obtained for the case $\beta_W = 0$. Increasing this parameter

leads to a small loss in terms of fitting and, at the same time, a decreasing number of non-zero values in the associated covariance matrix. The estimation of the spatial scale and the variance parameters are overall very similar, as expected, for $k = 0, 1, 2$ and for $\mu = 0.5, 1.5, 2.5$. A graphical comparison between empirical and estimated temporal semivariograms using model (3.1) when $\mu = 0.5$ and model (3.2) when $k = 0$ and $\beta_W = 0$ is provided in Figure S3 (OS).

In order to compare covariance models (3.1) and (3.2) from prediction performance and computational viewpoint, we use three predictive scores as described in Gneiting and Raftery (2007) and Zhang and Wang (2010). Let $\hat{Z}(\mathbf{x}_i, t_j)$ be the best linear prediction of Z at the space-time location (\mathbf{x}_i, t_j) based on all data except $z(\mathbf{x}_i, t_j)$. The first prediction score is the root mean square error (RMSE) defined as

$$\text{RMSE} = \left[\frac{1}{545 \times 11} \sum_{i=1}^{11} \sum_{j=366}^{910} \left(z(\mathbf{x}_i, t_j) - \hat{Z}(\mathbf{x}_i, t_j) \right)^2 \right]^{1/2}. \quad (3.3)$$

The logarithmic score is defined as

$$\log S = \frac{1}{545 \times 11} \sum_{i=1}^{11} \sum_{j=366}^{910} \left[\frac{1}{2} \log(2\pi\sigma(\mathbf{x}_i, t_j)) + \frac{1}{2} (Y(\mathbf{x}_i, t_j))^2 \right], \quad (3.4)$$

where $Y(\mathbf{x}_i, t_j) = \frac{z(\mathbf{x}_i, t_j) - \hat{Z}(\mathbf{x}_i, t_j)}{\sigma(\mathbf{x}_i, t_j)}$ and $\{\sigma(\mathbf{x}_i, t_j)\}^2$ is the prediction variance associated with $\hat{Z}(\mathbf{x}_i, t_j)$. Finally, we consider the continuous ranked probability

score (CRPS) which can be written in the Gaussian case as

$$\text{CRPS} = \frac{1}{545 \times 11} \sum_{i=1}^{11} \sum_{j=366}^{910} \sigma(\mathbf{x}_i, t_j) \left(Y(\mathbf{x}_i, t_j) [2F\{Y(\mathbf{x}_i, t_j)\} - 1] + 2F\{Y(\mathbf{x}_i, t_j)\} - \frac{1}{\sqrt{\pi}} \right), \quad (3.5)$$

where F is the Gaussian cumulative distribution. In Table 4, RMSE, logS and CRPS are shown for each considered covariance model. Comparing the covariances (3.1) and (3.2) for $\mu = 0.5, 1.5, 2.5$ and $k = 0, 1, 2$ respectively, a very small loss of prediction efficiency can be appreciated for the compactly supported models. For instance, when $\mu = 0.5$ and $k = 0$ and $\beta_W = 0$ the associated RMSE is 0.2174 and 0.2198, respectively.

The three prediction scores can be computed efficiently avoiding to calculate iteratively all these drop-one predictions involved in (3.3), (3.4) and (3.5) (Zhang and Wang, 2010). This efficient computation depends on the inverse of the covariance matrix. Let $\Sigma(\hat{\boldsymbol{\theta}})$ be the estimated covariance matrix associated to one of the covariance models considered in Equations (3.1) or (3.2). Then, for instance, RMSE can be written as $\text{RMSE} = (\mathbf{f}^\top \mathbf{f} / 5995)^{\frac{1}{2}}$ where $\mathbf{f} = D \Sigma(\hat{\boldsymbol{\theta}})^{-1} \mathbf{z}$ and $D = (\text{diag}(\Sigma(\hat{\boldsymbol{\theta}})^{-1}))^{-1}$.

As outlined in Furrer *et al.* (2006), efficient computation of the inverse of a (possibly large) symmetric positive definite matrix, with a given Cholesky matrix factorization, requires the solution of two triangular linear systems trough

back substitution. In our implementation, for covariance models obtained from Equation (3.2), the solution can be obtained through Cholesky factorization using the block sparse Cholesky algorithm of Ng and Peyton (1993) implemented in the *spam* package (Furrer and Sain, 2010). In Table 4, for a given percentage of non-zero values in the covariance matrix, we report the total time (in seconds) needed for computing the Cholesky factor and the inverse through back substitution for Gneiting-Wendland models. In Table 4 (bottom), we show the total time needed for computing the Cholesky factor through classical Cholesky decomposition, and the inverse through back substitution for Gneiting-Matérn models. Time in seconds is expressed in terms of elapsed time using the function *system.time* of the R software on a laptop with 2.4 GHz processor and 16 GB of memory. As expected, the computational gains obtained using Gneiting-Wendland models are huge. For instance, computation of the inverse is approximately 30 times faster with respect to the Gneiting-Matérn when comparing the cases $k = 0$, $\beta_W = 1$ and $\mu = 0.5$, and approximately 157 times faster when comparing the cases $k = 2$, $\beta_W = 1$ and $\mu = 2.5$. Similar computational gains can be achieved when computing classical space-time kriging and when performing simulation using Cholesky decomposition.

In conclusion, we have shown that our models allow for a substantial computational gain at the expense of a very small loss in terms of fitting and prediction

performance.

4. Conclusion

As outlined in Bevilacqua *et al.* (2018), in recent years the dataset sizes associated to spatially or spatio-temporally correlated random processes have steadily increased, so that straightforward statistical tools are computationally too expensive. The use of covariance functions with an (inherent or induced) compact support, leading to sparse matrices, is a very accessible and scalable approach. The nonseparable compactly supported space-time covariance models introduced in this paper have spatial (temporal) marginal covariance of the Generalized Wendland type and a dynamical decreasing compact support, an appealing feature from a computational viewpoint, in particular when dealing with datasets with a large number of location sites (temporal instants) and a relatively small number of temporal instants (location sites).

The recent work of Bevilacqua *et al.* (2018) gives a central importance to our covariance models with dynamical compact supports for prediction. In fact, Bevilacqua *et al.* (2018) showed that under some specific conditions, Matérn and Generalized Wendland covariance models are compatible, i.e., the induced Gaussian measures are equivalent. This implies that, under fixed domain asymptotics, the misspecified linear unbiased predictor with a Generalized Wendland

model is asymptotically as efficient as the true simple kriging predictor using a Matérn model. This fact applies to some extent to our space-time dynamical supports, albeit some caution is needed because of the apparent lack of a solid asymptotic framework that allows to merge fixed domain asymptotic in space with increasing domain in time.

Finally, the construction of nonseparable covariance models with both marginal covariances of the Generalized Wendland type and with dynamical decreasing compact support is very challenging from a theoretical point of view. This is an interesting possible topic for future research.

Online Supplement

The OS contains some material that integrates the main results provided in the manuscript. Specifically, Section 2 provides a generalization of Theorems 2.1 and 2.2 in the paper to a broad class of functions, called multiply monotonic. Section 3 explores necessary and sufficient conditions in a general framework through Fourier analysis. The OS reports the figures discussed through the paper.

Acknowledgement

The research work conducted by Moreno Bevilacqua was supported in part by FONDECYT grant 1160280 Chile. Emilio Porcu was partly supported by FONDECYT 1130647 Chile. The work of Marc G. Genton was supported by King Abdullah University of Science and Technology (KAUST).

References

- Askey, R. (1973) Radial characteristic functions. *Tech. Report No. 1262. - Math. Research Center*, University of Wisconsin-Madison.
- Buhmann, M. (2000) A new class of radial basis functions with compact support. *Mathematics of Computation*, 70, 233, 307-318.
- Bevilacqua, M., Faouzi, T., Furrer, R. and Porcu, E. (2018). Estimation and prediction using generalized Wendland functions under fixed domain asymptotics. *Annals of Statistics, Forthcoming*.
- Chernih, A., Sloan, I. H., and Womersley, R. S. (2014) Wendland functions with increasing smoothness converge to a Gaussian *Advances in Computational Mathematics.*, **142**, 17–33.
- Daley, D. J., Porcu, E. and Bevilacqua, M. (2015) Classes of compactly sup-

REFERENCES

- ported correlation functions for multivariate random fields. *Stochastic Environmental Research Risk Assessment*, **29**, 1249–1263.
- Furrer, R., Genton, M. G. and Nychka, D. (2006) Covariance tapering for interpolation of large spatial datasets. *Journal of Computational and Graphical Statistics*, **15**, 502–523.
- Furrer, R. and Sain, S. R. (2010) spam: A sparse matrix R package with emphasis on MCMC methods for Gaussian Markov random fields. *Journal of Statistical Software*, **36**, 1–25.
- Gneiting, T. (1999a) On the derivatives of radial positive definite functions. *Journal of Mathematical Analysis and Applications*, **236**, 86–93.
- Gneiting, T. (2000) Kuttner’s problem and a Pólya type criterion for characteristic functions. *Proceedings of the American Mathematical Society*, **128**, 1721–1728.
- Gneiting, T. (2001) Criteria of the Pólya type for radial positive definite functions. *Proceedings of the American Mathematical Society*, **129**, 8, 2309–2318.
- Gneiting, T. (2002a) Nonseparable, stationary covariance functions for space-time data. *Journal of the American Statistical Association*, **97**, 590–600.

REFERENCES

- Gneiting, T. (2002b) Compactly supported correlation functions. *J. Multiv. Anal.*, **83**, 493–508.
- Gneiting, T., Genton, M. G. and Guttorp, P. (2007) Geostatistical space-time models, stationarity, separability and full symmetry. In Finkenstadt, B., Held, L. and Isham, V. (eds.), *Statistical Methods for Space-Time Systems*, Chapman & Hall/CRC, Boca Raton, pp. 151–175.
- Gneiting, T. and Raftery, A. E. (2007) Strictly proper scoring rules, prediction, and estimation. *Journal of the American Statistical Association*, **102**, 359–378.
- Golubov, B. I. (1981) On Abel-Poisson type and Riesz means. *Analysis Mathematica*, **7**, 161–184.
- Haslett, J. and Raftery, A. (1989) space-time modelling with long-memory dependence: Assessing Ireland’s wind power resource. *Applied Statistics*, **38**, 1–50.
- Hubbert, S. (2012) Closed form representations for a class of compactly supported radial basis functions. *Advances in Computational Mathematics*, **36** (1), 115–136.
- Kaufman, C., Schervish, M. and Nychka, D. (2008) Covariance tapering for

REFERENCES

- likelihood-based estimation in large spatial datasets. *Journal of the American Statistical Association*, **103**, 1545–1569.
- Kuttner, B. (1944) On the Riesz means of a Fourier series (II). *Journal of the London Mathematical Society*, **19**, 77–84.
- Ng, E. G., and Peyton, B. W. (1993a) Block sparse Cholesky algorithms on advanced uniprocessor computers *SIAM Journal on Scientific Computing*, **14(5)**, 1064-8275.
- Nelder, J.A. and Mead, R (1965).A simplex algorithm for function minimization *Computer Journal* , **7**, 308-313.
- Pólya, G. (1949) Remarks on Characteristic Functions. *Proceedings of the [First] Berkeley Symposium on Mathematical Statistics and Probability*, 115–123, University of California Press, Berkeley, Calif.
- Porcu, E. and Schilling, R. (2011) From Schoenberg to Pick–Nevanlinna: Toward a complete picture of the variogram class. *Bernoulli*, **17**, 441–455.
- Porcu, E. and Zastavnyi, V. P. (2012). Characterization theorems for some classes of covariance functions associated to vector valued random fields. *Journal of Multivariate Analysis*, **102**, 1293–1301.

REFERENCES

- Porcu, E. and Zastavnyi, V. P. (2014) Generalized Askey functions and their walks through dimensions. *Exposithiones Mathematicae*, **32**, 190–198.
- Porcu, E., Zastavnyi, V. P. and Bevilacqua, M. (2018). Buhmann covariance functions, their compact supports, and their smoothness. *Dolomite Research Notes on Approximation*, to appear.
- R Development Core Team (2016) R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.
- Schaback, R. (2011) The missing Wendland functions. *Advances in Computational Mathematics*, **34**, 1, 67–81.
- Schaback, R. and Wu, Z. (1995) Operators on radial functions, *Journal of Computational and Applied Mathematics*, **73**(1-2), 257–270.
- Schlather, M. (2010) Covariance functions generated by Gaussian scale mixtures. *Bernoulli*, **16**, 3, 780–797.
- Stein, M. L. (1999) Statistical Interpolation of Spatial Data: Some Theory for Kriging. Springer, New York.
- Stein, M. L. (2005) Space-time covariance functions. *Journal of the American Statistical Association*, **100**, 310–321.

REFERENCES

- Wendland, H. (1995) Piecewise polynomial, positive definite and compactly supported radial functions of minimal degree. *Advances in Computational Mathematics*, **4**, 389–396.
- Wu, Z. (1995) Compactly supported positive definite radial functions. *Advances in Computational Mathematics*, **4**(3), 283–292.
- Zastavnyi, V. P. (2006) On some properties of Buhmann functions. *Ukrainian Mathematical Journal*, **58**, 8, 1184–12-9. Translated from *Ukrainskyi Matematychnyi Zhurnal*, 2006, **58**, 8, 1045–1067.
- Zastavnyi, V. P. and Porcu, E. (2011) Characterization theorems for the Gneiting class of covariance functions. *Bernoulli*, **17**, 456–465.
- Zastavnyi, V. P. and Trigub, R. M. (2002) Positive-definite splines of special form. *Matematicheskii Sbornik*, **193**, 1771–1800.
- Zhang, H. and Wang, Y. (2010) Kriging and cross-validation for massive spatial data. *Environmetrics*, **21**, 290–304.
- Zhu, S. (2012). Compactly supported radial basis functions: How and Why, *Oxford Preprint*.

REFERENCES

Table 3: Top: Bias and Standard Deviation (SD) for ML estimation of spatial and temporal scales and variance for the Gneiting-Wendland model in Equation (2.9) for $\kappa = 0, 1, 2$ and $\beta = 0, 0.5, 1$ under Scenario 1. Bottom: Scenario 2.

		a		b		σ^2	
κ	β	bias	SD	bias	SD	bias	SD
0	0	-0.00016	0.02168	0.00076	0.04889	0.00024	0.06181
	0.5	-0.00018	0.02145	0.00092	0.04796	0.00021	0.06164
	1	-0.00017	0.02121	0.00105	0.04690	0.00019	0.06156
1	0	0.00036	0.01342	-0.01943	0.12685	0.00027	0.06156
	0.5	0.00038	0.01342	-0.01841	0.12194	0.00028	0.06156
	1	0.00042	0.01342	-0.01596	0.11485	0.00034	0.06156
2	0	0.00054	0.01225	0.01040	0.18746	0.00034	0.06132
	0.5	0.00053	0.01225	0.00910	0.18185	0.00035	0.06132
	1	0.00054	0.01225	0.00636	0.17438	0.00037	0.06140
0	0	-0.00228	0.05000	0.00209	0.06812	0.00038	0.06419
	0.5	-0.00158	0.04743	0.00251	0.06745	0.00043	0.06411
	1	-0.00140	0.04506	0.00257	0.06626	0.00037	0.06409
1	0	-0.01809	0.11091	0.00064	0.03768	0.00084	0.06395
	0.5	-0.01763	0.10266	0.00086	0.03768	0.00040	0.06496
	1	-0.01741	0.09644	0.00103	0.03782	0.00095	0.06496
2	0	-0.01131	0.16199	0.00021	0.03674	0.00040	0.06380
	0.5	-0.01388	0.15556	0.00033	0.03688	0.00095	0.06372
	1	-0.01692	0.15063	0.00040	0.03688	0.00098	0.06372

REFERENCES

Table 4: Top: ML estimation of σ_W^2 , a_W , b_W for covariance model in equation (3.2) for $\kappa = 0, 1, 2$ and $\beta_W = 0, 0.5, 1$. RMSE, logS and CRPS computed using the ML estimated covariance matrix, the percentage of non zero values associated and time (in seconds) needed to compute inverse. Bottom part: ML estimation of σ_M^2 , a_M , b_M , β_M for covariance model in equation (3.1) for $\mu = 0, 1, 2$.

	β_W	a_W	b_W	σ_W^2	Loglik	RMSE	logS	CRPS	%	Time
$\kappa = 0$	0	1313.13	4.64	0.325	-691.23	0.2198	-0.1212	0.4399	1.64	4.5
	0.5	1274.87	3.95	0.323	-724.74	0.2210	-0.1140	0.4396	1.28	3.8
	1	1342.21	3.12	0.335	-788.79	0.2234	-0.1020	0.4419	0.95	3.6
$\kappa = 1$	0	2451.25	3.21	0.319	-765.29	0.2234	-0.1020	0.4419	1.28	3.7
	0.5	2500.77	2.88	0.324	-795.82	0.2240	-0.1036	0.4487	0.09	2.8
	1	2648.16	2.56	0.338	-829.81	0.2246	-0.1018	0.4504	0.09	2.8
$\kappa = 2$	0	3586.95	3.33	0.319	-773.65	0.2235	-0.1065	0.4492	1.28	3.7
	0.5	3637.85	3.09	0.323	-795.05	0.2239	-0.1043	0.4494	1.20	3.5
	1	3768.07	2.86	0.332	-818.62	0.2244	-0.1028	0.4503	0.09	2.8
	β_M	a_M	b_M	σ_M^2	Loglik	RMSE	logS	CRPS	%	Time
$\mu = 0.5$	0.54	1374.01	1.322	0.333	-634.44	0.2174	-0.1343	0.4375	100.00	109
$\kappa = 1.5$	1.0	2498.58	0.528	0.326	-702.77	0.2205	-0.1196	0.4437	100.00	393
$\kappa = 2.5$	1.0	3604.60	0.368	0.323	-724.16	0.2214	-0.1153	0.4454	100.00	441