

LEAST-SQUARES FORMULATIONS FOR EIGENVALUE PROBLEMS ASSOCIATED WITH LINEAR ELASTICITY

FLEURIANNE BERTRAND AND DANIELE BOFFI

ABSTRACT. We study the approximation of the spectrum of least-squares operators arising from linear elasticity. We consider a two-field (stress/displacement) and a three-field (stress/displacement/vorticity) formulation; other formulations might be analyzed with similar techniques.

We prove a priori estimates and we confirm the theoretical results with simple two-dimensional numerical experiments.

1. INTRODUCTION

In this paper we continue the investigations started in [3] about the spectral properties of operators associated with finite element least-squares formulations. While [3] deals with the Poisson eigenvalue problem, here we consider linear elasticity in the stress/displacement formulation. We discuss a two-field least-squares formulation introduced in [8] and a three-field least-squares formulation studied in [5].

We show that under natural approximation properties of the involved finite element spaces the discretization of the eigenvalue problems converges optimally by using the standard Babuška–Osborn theory [2, 6].

An important difference with respect to the case of the Poisson equation, is that for elasticity problem the resulting formulation is not symmetric. We will see that in most cases the eigenvalues are nevertheless real, but there are situations in which complex solutions are found.

As for the analysis presented in [3], it should be clear that the aim of this study is not to present a competitive method for the computation of elasticity eigenmodes. However, knowing the behavior of the spectrum of least-squares operators can be useful for several reasons; for instance, people interested in implementing least-squares approximation for transient problems, could benefit from such information.

Some numerical tests confirm our theoretical results.

2. PROBLEM SETTING

Given a polytopal domain $\Omega \in \mathbb{R}^d$ ($d = 2, 3$) with boundary $\partial\Omega$ divided into two parts Γ_D and Γ_N , we consider the stress/displacement formulation of linear elasticity as a system of first order as follows: find a symmetric d -by- d stress tensor $\underline{\sigma}$ and a displacement vectorfield \mathbf{u} such that

$$(1) \quad \begin{aligned} \mathcal{A}\underline{\sigma} - \underline{\varepsilon}(\mathbf{u}) &= \mathbf{0} && \text{in } \Omega \\ \mathbf{div} \underline{\sigma} &= -\mathbf{f} && \text{in } \Omega \\ \mathbf{u} &= \mathbf{0} && \text{on } \Gamma_D \\ \underline{\sigma}\mathbf{n} &= \mathbf{0} && \text{on } \Gamma_N, \end{aligned}$$

where the compliance tensor is defined in terms of the Lamé constants μ and λ

$$\mathcal{A}\underline{\boldsymbol{\tau}} = \frac{1}{2\mu} \left(\underline{\boldsymbol{\tau}} - \frac{\lambda}{2\mu + d\lambda} \operatorname{tr}(\underline{\boldsymbol{\tau}}) \mathbf{I} \right),$$

the symmetric gradient is defined as

$$\underline{\boldsymbol{\varepsilon}}(\mathbf{v}) = \frac{1}{2} (\underline{\nabla} \mathbf{v} + (\underline{\nabla} \mathbf{v})^\top),$$

and \mathbf{n} denotes the outward unit normal vector to Γ_N .

The formulation we are discussing has the important property to be robust in the incompressible limit. Other formulations could be studied as well. We refer the interested reader, for instance, to the introduction of [5] for an historical perspective.

A least-squares formulation of (1) was considered in [8] by minimizing the functional

$$\mathcal{F}(\underline{\boldsymbol{\tau}}, \mathbf{v}; \mathbf{f}) = \|\mathcal{A}\underline{\boldsymbol{\sigma}} - \underline{\boldsymbol{\varepsilon}}(\mathbf{u})\|_0^2 + \|\mathbf{div} \underline{\boldsymbol{\sigma}} + \mathbf{f}\|_0^2$$

in $\underline{\mathbf{X}}_N \times H_{0,D}^1(\Omega)^d$, with

$$\underline{\mathbf{X}} = \begin{cases} \mathbf{H}(\mathbf{div}; \Omega)^d & \text{if } \Gamma_N \neq \emptyset \\ \{\underline{\boldsymbol{\tau}} \in \mathbf{H}(\mathbf{div}; \Omega)^d : \int_\Omega \operatorname{tr}(\underline{\boldsymbol{\tau}}) d\mathbf{x} = 0\} & \text{if } \Gamma_N = \emptyset \end{cases}$$

and $\underline{\mathbf{X}}_N$ the subset of $\underline{\mathbf{X}}$ corresponding to the boundary condition $\underline{\boldsymbol{\tau}}\mathbf{n} = \mathbf{0}$ on Γ_N .

Remark 1. This formulation does not require a symmetry constraint in the definition of $\underline{\mathbf{X}}_N$ since the constitutive equation implies automatically that the solution satisfies $\underline{\boldsymbol{\sigma}} = \underline{\boldsymbol{\sigma}}^\top$. We shall refer to this formulation as the *two-field* formulation.

Other two formulations have been introduced in [5] which make use of three and four fields, respectively, by the introduction of the vorticity and the pressure as new unknowns. We describe the three-field formulation, similar considerations could be derived for the four-field formulation.

The three-field formulation seeks a minimizer of the functional

$$\mathcal{G}(\underline{\boldsymbol{\tau}}, \mathbf{v}, \boldsymbol{\varphi}; \mathbf{f}) = \|\mathcal{A}\underline{\boldsymbol{\tau}} - \underline{\nabla} \mathbf{v} + (-1)^d \underline{\boldsymbol{\chi}} \boldsymbol{\varphi}\|_0^2 + \|\mathbf{div} \underline{\boldsymbol{\sigma}} + \mathbf{f}\|_0^2 + \|\operatorname{as} \underline{\boldsymbol{\tau}}\|_0^2$$

in $\underline{\mathbf{X}}_N \times H_{0,D}^1(\Omega)^d \times \bar{L}(\Omega)^{2d-3}$ with

$$\bar{L}^2(\Omega) = \begin{cases} L^2(\Omega) & \text{if } \Gamma_N \neq \emptyset \\ \{\boldsymbol{\varphi} \in L^2(\Omega) : \int_\Omega \boldsymbol{\varphi} d\mathbf{x} = 0\} & \text{if } \Gamma_N = \emptyset, \end{cases}$$

where

$$\underline{\boldsymbol{\chi}} = \begin{cases} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} & \text{if } d = 2 \\ (\underline{\boldsymbol{\chi}}_1, \underline{\boldsymbol{\chi}}_2, \underline{\boldsymbol{\chi}}_3) & \text{if } d = 3 \end{cases}$$

with

$$\underline{\boldsymbol{\chi}}_1 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix} \quad \underline{\boldsymbol{\chi}}_2 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix} \quad \underline{\boldsymbol{\chi}}_3 = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

and where $\operatorname{as} \underline{\boldsymbol{\tau}} = (\underline{\boldsymbol{\tau}} - \underline{\boldsymbol{\tau}}^\top) / 2$ is the skew-symmetric part of $\underline{\boldsymbol{\tau}}$.

We are interested in the eigenvalue problem corresponding to (1), that is, we are looking for $\omega \in \mathbb{R}$ such that for non vanishing \mathbf{u} and for some $\underline{\boldsymbol{\sigma}}$ it holds

$$(2) \quad \begin{aligned} \mathcal{A}\underline{\boldsymbol{\sigma}} - \underline{\boldsymbol{\varepsilon}}(\mathbf{u}) &= 0 && \text{in } \Omega \\ \mathbf{div} \underline{\boldsymbol{\sigma}} &= -\omega \mathbf{u} && \text{in } \Omega \\ \mathbf{u} &= \mathbf{0} && \text{on } \Gamma_D \\ \underline{\boldsymbol{\sigma}} \mathbf{n} &= \mathbf{0} && \text{on } \Gamma_N. \end{aligned}$$

Thanks to the regularity properties of the solution of (1), the eigenvalue problem (2) is compact so that its eigenvalues form an increasing sequence

$$0 < \omega_1 \leq \omega_2 \leq \dots \leq \omega_i \leq \dots$$

and the eigenspaces are finite dimensional. As usual, we repeat the eigenvalues according to their multiplicity, so that each eigenvalue ω_i corresponds to a one-dimensional eigenspace E_i .

In order to approximate the eigenmodes, we are going to generalize the ideas of [3] to the two- and three-field formulations presented above. In particular, we are not writing directly a least-squares formulation of the eigenvalue problem (which would lead to a non-linear problem), but we study the spectrum of the operators associated with the least-squares *source* formulations.

2.1. Eigenvalue problem associated with the two-field formulation. The minimization of the functional $\mathcal{F}(\boldsymbol{\tau}, \mathbf{v}, \mathbf{f})$ gives rise to the following variational formulation: find $\underline{\boldsymbol{\sigma}} \in \underline{\mathbf{X}}_N$ and $\mathbf{u} \in H_{0,D}^1(\Omega)^d$ such that

$$(3) \quad \begin{cases} (\mathcal{A}\underline{\boldsymbol{\sigma}}, \mathcal{A}\boldsymbol{\tau}) + (\mathbf{div} \underline{\boldsymbol{\sigma}}, \mathbf{div} \boldsymbol{\tau}) - (\mathcal{A}\boldsymbol{\tau}, \underline{\boldsymbol{\varepsilon}}(\mathbf{u})) = -(\mathbf{f}, \mathbf{div} \boldsymbol{\tau}) & \forall \boldsymbol{\tau} \in \underline{\mathbf{X}}_N \\ -(\mathcal{A}\underline{\boldsymbol{\sigma}}, \underline{\boldsymbol{\varepsilon}}(\mathbf{v})) + (\underline{\boldsymbol{\varepsilon}}(\mathbf{u}), \underline{\boldsymbol{\varepsilon}}(\mathbf{v})) = 0 & \forall \mathbf{v} \in H_{0,D}^1(\Omega)^d. \end{cases}$$

The corresponding eigenvalue problem is obtained after replacing \mathbf{f} with $\omega \mathbf{u}$, that is: find $\omega \in \mathbb{C}$ such that for a non vanishing $\mathbf{u} \in H_{0,D}^1(\Omega)^d$ and for some $\underline{\boldsymbol{\sigma}} \in \underline{\mathbf{X}}_N$ we have

$$(4) \quad \begin{cases} (\mathcal{A}\underline{\boldsymbol{\sigma}}, \mathcal{A}\boldsymbol{\tau}) + (\mathbf{div} \underline{\boldsymbol{\sigma}}, \mathbf{div} \boldsymbol{\tau}) - (\mathcal{A}\boldsymbol{\tau}, \underline{\boldsymbol{\varepsilon}}(\mathbf{u})) = -\omega(\mathbf{u}, \mathbf{div} \boldsymbol{\tau}) & \forall \boldsymbol{\tau} \in \underline{\mathbf{X}}_N \\ -(\mathcal{A}\underline{\boldsymbol{\sigma}}, \underline{\boldsymbol{\varepsilon}}(\mathbf{v})) + (\underline{\boldsymbol{\varepsilon}}(\mathbf{u}), \underline{\boldsymbol{\varepsilon}}(\mathbf{v})) = 0 & \forall \mathbf{v} \in H_{0,D}^1(\Omega)^d. \end{cases}$$

The structure of the eigenvalue problem (4) in terms of operators is similar to the one described in [3] in the case of the FOSLS formulation for the Poisson equation. Namely, by natural definition of the operators, we are led to the following form:

$$(5) \quad \begin{pmatrix} \mathbf{A} & \mathbf{B}^\top \\ \mathbf{B} & \mathbf{C} \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \omega \begin{pmatrix} 0 & \mathbf{D} \\ 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}.$$

One important difference between this formulation and the one presented in [3] for the Poisson equation is that in our case the operators \mathbf{B}^\top and $-\mathbf{D}$ are not the same. It follows that it is not possible to show in general that (5) corresponds to a symmetric problem. This fact has important consequences for the numerical approximation. We will see in Section 5 that most of the computed eigenvalues are real, but there are exceptions.

2.2. Eigenvalue problem associated with the three-field formulation. The variational formulation associated with the minimization of the functional \mathcal{G} is obtained by seeking $\underline{\sigma} \in \underline{\mathbf{X}}_N$, $\mathbf{u} \in H_{0,D}^1(\Omega)^d$, and $\psi \in \bar{L}^2(\Omega)$ such that

$$(6) \quad \begin{cases} (\mathcal{A}\underline{\sigma}, \mathcal{A}\underline{\tau}) + (\mathbf{div} \underline{\sigma}, \mathbf{div} \underline{\tau}) + (\text{as}(\underline{\sigma}), \text{as}(\underline{\tau})) \\ \quad - (\mathcal{A}\underline{\tau}, \underline{\varepsilon}(\mathbf{u})) + (-1)^d (\mathcal{A}\underline{\tau}, \underline{\chi}\psi) = -(\mathbf{f}, \mathbf{div} \underline{\tau}) \quad \forall \underline{\tau} \in \underline{\mathbf{X}}_N \\ - (\mathcal{A}\underline{\sigma}, \underline{\varepsilon}(\mathbf{v})) + (\underline{\varepsilon}(\mathbf{u}), \underline{\varepsilon}(\mathbf{v})) - (-1)^d (\underline{\chi}\psi, \underline{\nabla} \mathbf{v}) = 0 \quad \forall \mathbf{v} \in H_{0,D}^1(\Omega)^d \\ (-1)^d (\mathcal{A}\underline{\sigma}, \underline{\chi}\varphi) - (-1)^d (\underline{\chi}\varphi, \underline{\nabla} \mathbf{u}) + (\underline{\chi}\psi, \underline{\chi}\varphi) = 0 \quad \forall \varphi \in \bar{L}^2(\Omega). \end{cases}$$

Also in this case we consider the eigenvalue problem by replacing \mathbf{f} with $\omega \mathbf{u}$ in the right hand side. The problems reads: find $\omega \in \mathbb{C}$ such that for a non vanishing $\mathbf{u} \in H_{0,D}^1(\Omega)^d$ and for some $\underline{\sigma} \in \underline{\mathbf{X}}_N$ and $\psi \in \bar{L}^2(\Omega)$ it holds

$$(7) \quad \begin{cases} (\mathcal{A}\underline{\sigma}, \mathcal{A}\underline{\tau}) + (\mathbf{div} \underline{\sigma}, \mathbf{div} \underline{\tau}) + (\text{as}(\underline{\sigma}), \text{as}(\underline{\tau})) \\ \quad - (\mathcal{A}\underline{\tau}, \underline{\varepsilon}(\mathbf{u})) + (-1)^d (\mathcal{A}\underline{\tau}, \underline{\chi}\psi) = -\omega(\mathbf{u}, \mathbf{div} \underline{\tau}) \quad \forall \underline{\tau} \in \underline{\mathbf{X}}_N \\ - (\mathcal{A}\underline{\sigma}, \underline{\varepsilon}(\mathbf{v})) + (\underline{\varepsilon}(\mathbf{u}), \underline{\varepsilon}(\mathbf{v})) - (-1)^d (\underline{\chi}\psi, \underline{\nabla} \mathbf{v}) = 0 \quad \forall \mathbf{v} \in H_{0,D}^1(\Omega)^d \\ (-1)^d (\mathcal{A}\underline{\sigma}, \underline{\chi}\varphi) - (-1)^d (\underline{\chi}\varphi, \underline{\nabla} \mathbf{u}) + (\underline{\chi}\psi, \underline{\chi}\varphi) = 0 \quad \forall \varphi \in \bar{L}^2(\Omega). \end{cases}$$

The operator form of the eigenvalue problem (7) involves 3-by-3 block operators as follows:

$$(8) \quad \begin{pmatrix} \mathbf{A} & \mathbf{B}^\top & \mathbf{C}^\top \\ \mathbf{B} & \mathbf{D} & \mathbf{E}^\top \\ \mathbf{C} & \mathbf{E} & \mathbf{F} \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \omega \begin{pmatrix} 0 & \mathbf{F} & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix}$$

Also in this case the system is not symmetric and the numerical approximation presented in Section 5 will show that some computed eigenvalues may be complex.

3. NUMERICAL APPROXIMATION

As it is apparent from formulations (4) and (7), the numerical approximation will lead to generalized eigenvalue problems of the form $\mathcal{A}x = \omega \mathcal{B}y$ where the matrix \mathcal{B} is singular. From the algebraic point of view, as we observed in [3], the solution of this problem satisfies the following properties.

- (1) If the matrix \mathcal{B} is invertible then our system is equivalent to the standard eigenvalue problem $\mathcal{B}^{-1}\mathcal{A}x = \omega x$.
- (2) If $\mathcal{K} = \ker \mathcal{A} \cap \ker \mathcal{B}$ is not trivial then the eigenvalue problem is degenerate and vectors in \mathcal{K} do not correspond to any eigenvalue of our original system.
- (3) If the matrix \mathcal{B} has a non-trivial kernel which does not contain any nonzero vector of $\ker(\mathcal{A})$ then it is conventionally assumed that our system has an eigenvalue $\omega = \infty$ with eigenspace equal to $\ker(\mathcal{B})$.
- (4) If \mathcal{B} is singular and \mathcal{A} is not (which is the most common situation in our framework) then it may be convenient to switch the roles of the two matrices and to consider the problem

$$\mathcal{B}x = \gamma \mathcal{A}x$$

Then (γ, x) with $\gamma = 0$ corresponds to the eigenmode (∞, x) of the original system; the remaining eigenmodes are (ω, x) with $\omega = 1/\gamma$.

3.1. Approximation of the two-field formulation. Given finite dimensional subspaces $\Sigma_h \subset \underline{\mathbf{X}}_N$ and $U_h \subset H_{0,D}^1(\Omega)^d$, the Galerkin approximation of (4) reads: find $\omega_h \in \mathbb{C}$ such that for a non vanishing $\mathbf{u}_h \in U_h$ and for some $\underline{\boldsymbol{\sigma}}_h \in \Sigma_h$ we have

$$(9) \quad \begin{cases} (\mathcal{A}\underline{\boldsymbol{\sigma}}_h, \mathcal{A}\underline{\boldsymbol{\tau}}) + (\mathbf{div} \underline{\boldsymbol{\sigma}}_h, \mathbf{div} \underline{\boldsymbol{\tau}}) - (\mathcal{A}\underline{\boldsymbol{\tau}}, \underline{\boldsymbol{\varepsilon}}(\mathbf{u}_h)) = -\omega_h(\mathbf{u}_h, \mathbf{div} \underline{\boldsymbol{\tau}}) & \forall \underline{\boldsymbol{\tau}} \in \Sigma_h \\ -(\mathcal{A}\underline{\boldsymbol{\sigma}}_h, \underline{\boldsymbol{\varepsilon}}(\mathbf{v})) + (\underline{\boldsymbol{\varepsilon}}(\mathbf{u}_h), \underline{\boldsymbol{\varepsilon}}(\mathbf{v})) = 0 & \forall \mathbf{v} \in U_h. \end{cases}$$

The structure of this problem is analogous of (5) with the natural definition of the involved matrices. The following proposition is the analogue of [3, Prop. 6] and characterizes the number of significant eigenvalues of (9).

Proposition 1. *The solution of the generalized eigenvalue problem associated with the formulation (9) include $\omega = \infty$ with multiplicity equal to $\dim(\Sigma_h) + \dim(\ker(\mathbf{D}))$ and a number of complex eigenvalues (counted with their multiplicity) equal to $\text{rank}(\mathbf{D})$.*

Remark 2. Since the eigenvalue problem stated in (9) considers eigenfunctions with $\mathbf{u}_h \neq \mathbf{0}$, the total number of eigenvalues of the problem we are interested in, is equal to $\dim U_h$; the corresponding values are the $\text{rank}(\mathbf{D})$ complex solutions and possibly $\omega = \infty$ if \mathbf{D} is not full rank.

3.2. Approximation of the three-field formulation. Given finite dimensional subspaces $\Sigma_h \subset \underline{\mathbf{X}}_N$, $U_h \subset H_{0,D}^1(\Omega)^d$, and $\Phi_h \subset \bar{L}^2(\Omega)$, the Galerkin approximation of (7) is: find $\omega_h \in \mathbb{C}$ such that for a non vanishing $\mathbf{u}_h \in U_h$ and for some $\underline{\boldsymbol{\sigma}}_h \in \Sigma_h$ and $\psi_h \in \Phi_h$ we have

$$(10) \quad \begin{cases} (\mathcal{A}\underline{\boldsymbol{\sigma}}_h, \mathcal{A}\underline{\boldsymbol{\tau}}) + (\mathbf{div} \underline{\boldsymbol{\sigma}}_h, \mathbf{div} \underline{\boldsymbol{\tau}}) + (\text{as}(\underline{\boldsymbol{\sigma}}_h), \text{as}(\underline{\boldsymbol{\tau}})) \\ \quad - (\mathcal{A}\underline{\boldsymbol{\tau}}, \underline{\boldsymbol{\varepsilon}}(\mathbf{u}_h)) + (-1)^d (\mathcal{A}\underline{\boldsymbol{\tau}}, \underline{\boldsymbol{\chi}}\psi_h) = -\omega_h(\mathbf{u}_h, \mathbf{div} \underline{\boldsymbol{\tau}}) & \forall \underline{\boldsymbol{\tau}} \in \Sigma_h \\ -(\mathcal{A}\underline{\boldsymbol{\sigma}}_h, \underline{\boldsymbol{\varepsilon}}(\mathbf{v})) + (\underline{\boldsymbol{\varepsilon}}(\mathbf{u}_h), \underline{\boldsymbol{\varepsilon}}(\mathbf{v})) - (-1)^d (\underline{\boldsymbol{\chi}}\psi_h, \underline{\boldsymbol{\nabla}} \mathbf{v}) = 0 & \forall \mathbf{v} \in U_h \\ (-1)^d (\mathcal{A}\underline{\boldsymbol{\sigma}}_h, \underline{\boldsymbol{\chi}}\varphi) - (-1)^d (\underline{\boldsymbol{\chi}}\varphi, \underline{\boldsymbol{\nabla}} \mathbf{u}_h) + (\underline{\boldsymbol{\chi}}\psi_h, \underline{\boldsymbol{\chi}}\varphi) = 0 & \forall \varphi \in \Phi_h. \end{cases}$$

The matrix structure of this problem corresponds to (7) and the following proposition, in analogy of Proposition 1, gives the characterization of the discrete eigenvalues.

Proposition 2. *The solution of the generalized eigenvalue problem associated with the formulation (10) consists of $\omega = \infty$ and a number of complex eigenvalues (counted with their multiplicity) equal to $\text{rank}(\mathbf{F})$.*

Remark 3. As in Remark 2, the eigenvalues of formulation (10) are $\text{rank}(\mathbf{F})$ complex values and possibly $\omega = \infty$ if \mathbf{F} is not full rank.

4. A PRIORI ERROR ESTIMATES

It is quite standard to obtain a priori error estimates for eigenvalue problems by studying the convergence of the discrete solution operator towards the continuous one [6]. In our framework, this boils down to showing $L^2(\Omega)$ estimates for the discretization of (3) and (6), respectively, when the right hand side \mathbf{f} is in $L^2(\Omega)^d$. For brevity, we omit the details and we refer the interested reader to [3].

4.1. $L^2(\Omega)$ estimate for the two-field formulation. The $L^2(\Omega)$ estimate for the two-field formulation (3) is the natural generalization of what has been presented in [3] in the case of the Poisson problem. The original idea comes from [1, Sec. 7] (see also [7]).

Let $(\underline{\sigma}, \mathbf{u}) \in \underline{\mathbf{X}}_N \times H_{0,D}^1(\Omega)^2$ solve (3) and $(\underline{\sigma}_h, \mathbf{u}_h) \in \Sigma_h \times U_h$ solve the corresponding discrete problem; we consider the dual problem: find $\underline{\mathbf{s}} \in \underline{\mathbf{X}}_N$ and $\mathbf{p} \in H_{0,D}^1(\Omega)^d$ such that

$$(11) \quad \begin{cases} (\mathcal{A}\underline{\mathbf{s}}, \mathcal{A}\underline{\mathbf{t}}) + (\mathbf{div} \underline{\mathbf{s}}, \mathbf{div} \underline{\mathbf{t}}) - (\mathcal{A}\underline{\mathbf{t}}, \underline{\boldsymbol{\varepsilon}}(\mathbf{p})) = 0 & \forall \underline{\mathbf{t}} \in \underline{\mathbf{X}}_N \\ -(\mathcal{A}\underline{\mathbf{s}}, \underline{\boldsymbol{\varepsilon}}(\mathbf{q})) + (\underline{\boldsymbol{\varepsilon}}(\mathbf{p}), \underline{\boldsymbol{\varepsilon}}(\mathbf{q})) = (\mathbf{u} - \mathbf{u}_h, \mathbf{q}) & \forall \mathbf{q} \in H_{0,D}^1(\Omega)^d. \end{cases}$$

Taking as test functions $\underline{\mathbf{t}} = \underline{\sigma} - \underline{\sigma}_h$ and $\mathbf{q} = \mathbf{u} - \mathbf{u}_h$ we have

$$\begin{aligned} \|\mathbf{u} - \mathbf{u}_h\|_0^2 &= (\mathcal{A}\underline{\mathbf{s}}, \mathcal{A}(\underline{\sigma} - \underline{\sigma}_h)) + (\mathbf{div} \underline{\mathbf{s}}, \mathbf{div}(\underline{\sigma} - \underline{\sigma}_h)) \\ &\quad - (\underline{\boldsymbol{\varepsilon}}(\mathbf{p}), \mathcal{A}(\underline{\sigma} - \underline{\sigma}_h)) - (\mathcal{A}\underline{\mathbf{s}}, \underline{\boldsymbol{\varepsilon}}(\mathbf{u} - \mathbf{u}_h)) \\ &\quad + (\underline{\boldsymbol{\varepsilon}}(\mathbf{p}), \underline{\boldsymbol{\varepsilon}}(\mathbf{u} - \mathbf{u}_h)) \\ &= (\mathcal{A}(\underline{\mathbf{s}} - \underline{\boldsymbol{\tau}}_h), \mathcal{A}(\underline{\sigma} - \underline{\sigma}_h)) + (\mathbf{div}(\underline{\mathbf{s}} - \underline{\boldsymbol{\tau}}_h), \mathbf{div}(\underline{\sigma} - \underline{\sigma}_h)) \\ &\quad - (\underline{\boldsymbol{\varepsilon}}(\mathbf{p} - \mathbf{v}_h), \mathcal{A}(\underline{\sigma} - \underline{\sigma}_h)) - (\mathcal{A}(\underline{\mathbf{s}} - \underline{\boldsymbol{\tau}}_h), \underline{\boldsymbol{\varepsilon}}(\mathbf{u} - \mathbf{u}_h)) \\ &\quad + (\underline{\boldsymbol{\varepsilon}}(\mathbf{p} - \mathbf{v}_h), \underline{\boldsymbol{\varepsilon}}(\mathbf{u} - \mathbf{u}_h)) \end{aligned}$$

for all $\underline{\boldsymbol{\tau}}_h \in \Sigma_h$ and $\mathbf{v} \in U_h$, where we used the error equations corresponding to (3). It follows

$$\begin{aligned} \|\mathbf{u} - \mathbf{u}_h\|_0^2 &\leq C (\|\underline{\mathbf{s}} - \underline{\boldsymbol{\tau}}_h\|_{\mathbf{div}} + \|\mathbf{p} - \mathbf{v}_h\|_1) (\|\underline{\sigma} - \underline{\sigma}_h\|_{\mathbf{div}} + \|\mathbf{u} - \mathbf{u}_h\|_1) \\ &\leq C (\|\underline{\mathbf{s}} - \underline{\boldsymbol{\tau}}_h\|_{\mathbf{div}} + \|\mathbf{p} - \mathbf{v}_h\|_1) \|\mathbf{f}\|_0. \end{aligned}$$

We observe explicitly that we cannot obtain any rate of convergence out of the term $\|\underline{\sigma} - \underline{\sigma}_h\|_{\mathbf{div}}$ since $\mathbf{div} \underline{\sigma}$ is only in $L^2(\Omega)$ if \mathbf{f} is not more regular. On the other hand, the required uniform convergence comes from the (minimal) regularity of the dual problem (11) so that we get

$$\|\mathbf{u} - \mathbf{u}_h\|_0^2 \leq Ch^s \|\mathbf{u} - \mathbf{u}_h\|_0 \|\mathbf{f}\|_0$$

for some positive s as long as the finite element spaces Σ_h and U_h satisfies the following approximation properties

$$(12) \quad \begin{aligned} \inf_{\underline{\boldsymbol{\tau}} \in \Sigma_h} \|\underline{\mathbf{s}} - \underline{\boldsymbol{\tau}}\|_{\mathbf{div}} &\leq Ch^s \|\underline{\mathbf{s}}\|_s + \|\mathbf{div} \underline{\mathbf{s}}\|_s \\ \inf_{\mathbf{v} \in U_h} \|\mathbf{p} - \mathbf{v}_h\|_1 &\leq Ch^s \|\mathbf{p}\|_{1+s}. \end{aligned}$$

Remark 4. The power s appearing in the estimate of $\|\mathbf{u} - \mathbf{u}_h\|_0$, which is limited by $\mathbf{f} \in L^2(\Omega)^d$, is not related to the rate of convergence of the numerical scheme, but guarantees the uniform convergence that implies the correct approximation of the spectrum. The optimal convergence of the scheme is shown in the next theorem by using the full regularity of the eigenspaces.

Theorem 3. *Let $\omega_i = \omega_{i+1} = \dots = \omega_{i+m-1}$ be an eigenvalue of (4) and denote by $E = \bigoplus_{j=i}^{i+m-1} E_j$ the corresponding eigenspace. If the discrete spaces Σ_h and U_h satisfy the approximation properties (12) then for h small enough (so that $\dim U_h \geq i+m-1$) the m discrete eigenvalues $\omega_{i,h} = \omega_{i+1,h} = \dots = \omega_{i+m-1,h}$ of (9) converge*

to ω_i ; let us denote by E_h the sum of the discrete eigenspaces. Then the following error estimates hold true

$$\begin{aligned} \delta(E, E_h) &\leq C\rho(h) \\ |\omega_j - \omega_{j,h}| &\leq \rho(h)^2 \quad j = i, \dots, i + m - 1 \end{aligned}$$

with

$$\rho(h) = \sup_{\substack{\mathbf{u} \in E \\ \|\mathbf{u}\|=1}} \inf_{\substack{\boldsymbol{\tau} \in \Sigma_h \\ \mathbf{v} \in U_h}} (\|\underline{\boldsymbol{\varepsilon}}(\mathbf{u}) - \boldsymbol{\tau}\|_{\mathbf{div}} + \|\mathbf{u} - \mathbf{v}\|_1).$$

Proof. The proof is standard (see [3, 6]); the only delicate part consists in showing the double order convergence for the eigenvalues, since the discrete problem is not symmetric. The result can be obtained by considering the adjoint problem, performing the corresponding analysis for its approximation (with $\rho(h)$ replaced by $\rho^*(h)$) and by using the standard Babuška–Osborn theory [2] from which we can conclude that the error of the eigenvalues is bounded by $\rho(h)\rho^*(h)$ (note that the continuous eigenvalues have ascent multiplicity equal to one since the continuous problem is symmetric). \square

4.2. $L^2(\Omega)$ -estimate for the three-field formulation. The $L^2(\Omega)$ error estimate for the discretization of the three-field formulation (7) has been presented in [5, Theorem 3] in the case of a convex domain so that the H^2 regularity of a generalized Stokes problem could be used (see [7]). Since we are not interested in optimal estimates, but only in the uniform convergence in the spirit of Remark 4, the arguments of [7] and [5] could be adapted in order to get the estimate

$$\|\mathbf{u} - \mathbf{u}_h\|_0 \leq Ch^s \|\mathbf{f}\|_0$$

where \mathbf{u} solves (6) and \mathbf{u}_h is the corresponding discrete solution, provided the following approximation properties are satisfied

$$(13) \quad \begin{aligned} \inf_{\boldsymbol{\tau} \in \Sigma_h} \|\underline{\mathbf{s}} - \boldsymbol{\tau}\|_{\mathbf{div}} &\leq Ch^s \|\underline{\mathbf{s}}\|_s + \|\mathbf{div} \underline{\mathbf{s}}\|_s \\ \inf_{\mathbf{v} \in U_h} \|\mathbf{p} - \mathbf{v}_h\|_1 &\leq Ch^s \|\mathbf{p}\|_{1+s} \\ \inf_{\boldsymbol{\varphi} \in \Phi_h} \|\boldsymbol{\psi} - \boldsymbol{\varphi}\|_0 &\leq Ch^s \|\boldsymbol{\psi}\|_s. \end{aligned}$$

This allows to state the following convergence theorem which is the analogous of Theorem 3 in this situation. In analogy to (12) we make explicit the requirements on the approximation properties of our spaces.

Theorem 4. *Let $\omega_i = \omega_{i+1} = \dots = \omega_{i+m-1}$ be an eigenvalue of (7) and denote by $E = \bigoplus_{j=i}^{i+m-1} E_j$ the corresponding eigenspace. If the discrete spaces Σ_h , U_h , and Φ_h satisfy the approximation properties (13) then for h small enough (so that $\dim U_h \geq i + m - 1$) the m discrete eigenvalues $\omega_{i,h} = \omega_{i+1,h} = \dots = \omega_{i+m-1,h}$ of (10) converge to ω_i ; let us denote by E_h the sum of the discrete eigenspaces. Then the following error estimates hold true*

$$\begin{aligned} \delta(E, E_h) &\leq C\rho(h) \\ |\omega_j - \omega_{j,h}| &\leq \rho(h)^2 \quad j = i, \dots, i + m - 1 \end{aligned}$$




FIGURE 1. Meshes on the unit square

with

$$\rho(h) = \sup_{\substack{\mathbf{u} \in E \\ \|\mathbf{u}\|=1}} \inf_{\substack{\boldsymbol{\tau} \in \Sigma_h \\ \mathbf{v} \in U_h \\ \boldsymbol{\varphi} \in \Phi_h}} (\|\boldsymbol{\nabla}(\mathbf{u}) - \boldsymbol{\tau}\|_{\text{div}} + \|\mathbf{u} - \mathbf{v}\|_1 + \|\mathbf{curl}(\mathbf{u}) - \boldsymbol{\varphi}\|_0).$$

5. NUMERICAL RESULTS

Our numerical results confirm the theoretical investigations of the previous sections.

The numerical solution of a generalized eigenvalue problem such as those arising from the discretization of (4) and (7) can be challenging and in this paper we do not focus on the efficiency of the solver but rather on the accuracy of the obtained results.

5.1. Numerical results on the square. We start by taking the domain equal to the unit square $\Omega =]0, 1[^2$. In this case a pretty accurate estimate of the first eigenvalue is given by $\omega = 52.344691168$ if we consider

$$\mathcal{A}\boldsymbol{\tau} = \frac{1}{2} \left(\boldsymbol{\tau} - \frac{1}{2} \text{tr}(\boldsymbol{\tau}) \mathbf{I} \right),$$

which corresponds to a Stokes problem, and homogeneous Dirichlet boundary conditions everywhere, that is $\Gamma_N = \emptyset$ (see, for instance, [9]). Since we also want to investigate the symmetry of the formulation, we consider three different mesh sequences: a symmetric and uniform mesh (CROSSED), a non-symmetric and uniform mesh (RIGHT), and a non structured non symmetric mesh (NONUNIF). The meshes are indexed with respect to the number N of subdivisions of the square's sides. The three meshes for $N = 4$ are plotted in Figure 1.

In Table 1 we report the numerical results corresponding to computations performed with the two-field scheme (9) when using the three mesh sequences with the level of refinement varying from $N = 4$ to $N = 12$. We considered a second order scheme based on Raviart–Thomas elements for the approximation of Σ_h (satisfying assumption (12)). The estimate of Theorem 3 predicts fourth order of convergence for the eigenvalues which is confirmed by our tests. It is interesting to observe that the first computed eigenvalue is always real in our tests.

We performed the same computations for the three-field formulation presented in (10). Also in this case we use a second order scheme based on Raviart–Thomas elements (satisfying assumption (13)). The corresponding results are reported in Table 2. It turns out that also in this case the first computed eigenvalue is always real and that the convergence properties match with the theoretical results with

TABLE 1. Eigenvalues of the two-field formulation on the unit square: $\Sigma_h = RT_1$, $U_h = P_2^c$

Mesh	$N = 4$	$N = 6$ (rate)	$N = 8$ (rate)	$N = 10$ (rate)	$N = 12$ (rate)
CROSSED	52.618734	52.400609 (3.9)	52.362201 (4.0)	52.351749 (4.1)	52.348048 (4.1)
RIGHT	54.132943	52.751624 (3.7)	52.480276 (3.8)	52.401472 (3.9)	52.372369 (3.9)
NONUNIF	52.744298	52.435687 (3.6)	52.368139 (4.7)	52.354017 (4.1)	52.349733 (3.4)

TABLE 2. Eigenvalues of the three-field formulation on the unit square: $\Sigma_h = RT_1$, $U_h = P_2^c$, $\Phi_h = P_1^d$

Mesh	$N = 4$	$N = 6$ (rate)	$N = 8$ (rate)	$N = 10$ (rate)	$N = 12$ (rate)
CROSSED	52.523637	52.377459 (4.2)	52.353859 (4.4)	52.348025 (4.5)	52.346144 (4.6)
RIGHT	53.712947	52.621373 (3.9)	52.426543 (4.2)	52.375437 (4.4)	52.358317 (4.5)
NONUNIF	52.630390	52.398013 (4.1)	52.355912 (5.4)	52.348310 (5.1)	52.347239 (1.9)

TABLE 3. Eigenvalues of the three-field formulation on the unit square: one more refinement for the non uniform mesh

Mesh	$N = 6$	$N = 8$ (rate)	$N = 10$ (rate)	$N = 12$ (rate)	$N = 14$ (rate)
NONUNIF	52.398013	52.355912 (3.8)	52.348310 (3.9)	52.347239 (1.6)	52.345561 (5.9)

TABLE 4. First five eigenvalues computed with the three-field scheme on the non-uniform mesh for $N = 10$

#	Value
1	52.348309870785620
2	$92.163865261631784 - 0.000422328750065i$
3	$92.163865261631784 + 0.000422328750065i$
4	128.2902654382040
5	154.3710922938166

the exception of the convergence rate for $N = 12$ on the non-uniform mesh. In order to check better this phenomenon, we computed one more refinement which is reported in Table 3. It is clear that the overall convergence matches with the expected fourth order rate and that the non uniform behavior is due to the fact that the mesh sequence is not structured.

Since the eigenvalue problems corresponding to the considered formulations are not symmetric, it is interesting to investigate whether the computed eigenvalues are complex or real. From the results that we are going show, it is clear that in some cases the computed eigenvalues have a non vanishing imaginary part; this implies that in general the discrete formulations cannot be reduced to symmetric problems. On the other hand, in most cases the computed eigenvalues are real; this occurs, in particular on the symmetric meshes.

In Table 4 we report the first five eigenvalues computed with the three-field scheme on the non-uniform mesh for $N = 10$ (similar results could be presented for the two-field scheme as well). It is apparent that the second and the third eigenvalue are complex conjugate.

TABLE 5. First five eigenvalues computed with the three-field scheme on the non-uniform mesh for $N = 11$

#	Value
1	52.346475661045424
2	92.147313995882541
3	92.151062887227738
4	128.2536615472890
5	154.2967136612170




FIGURE 2. Mesh of the L-shaped domain

TABLE 6. Eigenvalues of the two-field formulation on the L-shaped domain: $\Sigma_h = RT_1$, $U_h = P_2^c$

Mesh	$N = 4$	$N = 8$ (rate)	$N = 16$ (rate)	$N = 32$ (rate)
Uniform	35.606285	31.937374 (4.2)	31.871123 (-0.4)	31.983573 (0.8)

TABLE 7. Eigenvalues of the three-field formulation on the L-shaped domain: $\Sigma_h = RT_1$, $U_h = P_2^c$, $\Phi_h = P_1^d$

Mesh	$N = 4$	$N = 8$ (rate)	$N = 16$ (rate)	$N = 32$ (rate)
Uniform	34.132843	31.491151 (1.6)	31.677105 (0.5)	31.888816 (0.9)

For the sake of comparison, Table 5 shows the same results on the mesh for $N = 11$ and we can see that in this case all the first five eigenvalues are real. It is also interesting to observe that the second and the third eigenvalues on the mesh for $N = 10$ have the same real part, while the corresponding eigenvalues on the mesh for $N = 11$ are real and different from each other. Clearly they are approximation of the double eigenvalue $\omega_2 = \omega_3$. In any case, this is perfectly compatible with the statements of Theorems 3 and 4 where the difference $|\omega_j - \omega_{j,h}|$ has to be understood in the sense of the distance in the complex plane.

5.2. Numerical results on the L-shaped domain. We conclude our numerical tests with computations on the L-shaped domain where the re-entrant corner causes singularities in the solution. We consider a uniform mesh (see Figure 2 for the case $N = 4$).

An estimate of the first eigenvalues in this case has been provided in [4] and corresponds to $\omega = 32.13269464746$. Tables 6 and 7 report on the computed approximation with the two- and three-field approximations (9) and (10), respectively. As expected, the singularity of the eigenfunction corresponding to the first eigenvalue causes a reduction of the order of convergence.

TABLE 8. Some eigenvalues computed with the three-field scheme on the L-shaped domain for $N = 8$

#	Value
38	339.9524318713583
39	346.0018703851194
40	$350.8454478342160 - 2.5574107928386i$
41	$350.8454478342160 + 2.5574107928386i$
42	359.0078935078376
43	378.8779264703741

Also in this case the first computed eigenvalue is real, however, also in presence of a symmetric mesh, it turns out that there might be complex eigenvalues. For instance, Table 8 reports some higher eigenvalues computed for $N = 8$. Also in this case the complex eigenvalues converge towards real number according to the statement of Theorems 3 and 4; moreover, the presence of complex eigenvalues depends on the mesh and on the level of refinements. The effect of the mesh on complex eigenvalues will be the object of further investigation.

REFERENCES

1. Douglas N. Arnold, Daniele Boffi, and Richard S. Falk, *Quadrilateral $H(\text{div})$ finite elements*, SIAM J. Numer. Anal. **42** (2005), no. 6, 2429–2451. MR 2139400
2. Ivo Babuška and John Osborn, *Eigenvalue problems*, Handbook of numerical analysis, Vol. II, Handb. Numer. Anal., II, North-Holland, Amsterdam, 1991, pp. 641–787.
3. Fleurianne Bertrand and Daniele Boffi, *First order least-squares formulations for eigenvalue problems*, arXiv:2002.08145 [math.NA], 2020.
4. Fleurianne Bertrand, Daniele Boffi, and Rui Ma, *An adaptive finite element scheme for the Hellinger–Reissner elasticity mixed eigenvalue problem*, submitted, 2020.
5. Fleurianne Bertrand, Zhiqiang Cai, and Eun Young Park, *Least-squares methods for elasticity and Stokes equations with weakly imposed symmetry*, Comput. Methods Appl. Math. **19** (2019), no. 3, 415–430. MR 3977481
6. Daniele Boffi, *Finite element approximation of eigenvalue problems*, Acta Numer. **19** (2010), 1–120. MR 2652780
7. Zhiqiang Cai and Jaeun Ku, *The L^2 norm error estimates for the div least-squares method*, SIAM J. Numer. Anal. **44** (2006), no. 4, 1721–1734. MR 2257124
8. Zhiqiang Cai and Gerhard Starke, *Least-squares methods for linear elasticity*, SIAM J. Numer. Anal. **42** (2004), no. 2, 826–842. MR 2084237
9. Joscha Gedicke and Arbaz Khan, *Arnold–Winther mixed finite elements for Stokes eigenvalue problems*, SIAM J. Sci. Comput. **40** (2018), no. 5, A3449–A3469. MR 3864690

HUMBOLDT-UNIVERSITÄT ZU BERLIN, GERMANY AND KING ABDULLAH UNIVERSITY OF SCIENCE AND TECHNOLOGY, SAUDI ARABIA

KING ABDULLAH UNIVERSITY OF SCIENCE AND TECHNOLOGY, SAUDI ARABIA, AND UNIVERSITY OF PAVIA, ITALY