



Temperature dependence of the relationship between pCO₂ and dissolved organic carbon in lakes

Item Type	Article
Authors	Pinho, L.;Duarte, Carlos M.;Marotta, H.;Enrich-Prast, A.
Citation	Temperature dependence of the relationship between pCO ₂ and dissolved organic carbon in lakes 2016, 13 (3):865 Biogeosciences
Eprint version	Publisher's Version/PDF
DOI	10.5194/bg-13-865-2016
Publisher	Copernicus GmbH
Journal	Biogeosciences
Rights	This work is distributed under the Creative Commons Attribution 3.0 License.
Download date	2023-12-06 08:25:19
Link to Item	http://hdl.handle.net/10754/600213



Temperature dependence of the relationship between $p\text{CO}_2$ and dissolved organic carbon in lakes

L. Pinho^{1,2,3}, C. M. Duarte^{2,4}, H. Marotta^{5,6}, and A. Enrich-Prast^{1,7}

¹Postgraduate Program in Ecology, Department of Ecology, Institute of Biology, Federal University of Rio de Janeiro, Av. Carlos Chagas Filho, 373, Cidade Universitária (Ilha do Fundão), P. O. Box 68020, Rio de Janeiro – RJ, 21941-901, Brazil

²Global Change Department, IMEDEA (CSIC-UIB), Mediterranean Institute for Advanced Studies, C. Miquel Marquès, 21, 07190 Esporles, Balearic Islands, Spain

³Department of Chemical Oceanography, Rio de Janeiro State University, Pavilhão João Lyra Filho, sala 4008 Bloco E, Rua São Francisco Xavier, 524, Maracanã – RJ, 20550-900, Brazil

⁴King Abdullah University of Science and Technology (KAUST), Red Sea Research Center (RSRC), Thuwal 23955-6900, Saudi Arabia

⁵Research Center on Biomass and Water Management (NAB/UFF), Sedimentary Environmental Processes Laboratory (LAPSA/UFF), International Laboratory of Global Change (LINC Global), Federal Fluminense University, Av. Litorânea s/n, Campus Praia Vermelha, Niterói – RJ, 24210-340, Brazil

⁶Postgraduate Program in Geography, Postgraduate Program in Geosciences (Geochemistry), Federal Fluminense University, Niterói – RJ, 24220-900, Brazil

⁷Department of Environmental Changes, Linköping University, 581 83 Linköping, Sweden

Correspondence to: L. Pinho (luana.pinho@uerj.br)

Received: 25 November 2014 – Published in Biogeosciences Discuss.: 6 February 2015

Revised: 11 January 2016 – Accepted: 24 January 2016 – Published: 15 February 2016

Abstract. The relationship between the partial pressure of carbon dioxide ($p\text{CO}_2$) and dissolved organic carbon (DOC) concentration in Brazilian lakes, encompassing 225 samples across a wide latitudinal range in the tropics, was tested. Unlike the positive relationship reported for lake waters, which was largely based on temperate lakes, we found no significant relationship for low-latitude lakes ($<33^\circ$), despite very broad ranges in both $p\text{CO}_2$ and DOC levels. These results suggest substantial differences in the carbon cycling of low-latitude lakes, which must be considered when upscaling limnetic carbon cycling to global scales.

1 Introduction

Lakes cover less than 2 % of the continent's surface (Downing et al., 2006; McDonald et al., 2012) but play a significant role in the global carbon (C) cycle (Cole et al., 1994, 2007; Tranvik et al., 2009), contributing significantly to C burial and emissions to the atmosphere (Cole et al., 2007;

Downing et al., 2008; Tranvik et al., 2009). Dissolved organic carbon (DOC) represents a major C pool in lakes, with both autochthonous and allochthonous contributions (Duarte and Prairie, 2005; Cole et al., 2007; Prairie, 2008; Tranvik et al., 2009), supporting heterotrophy (Sobek et al., 2007) and affecting key biological and physicochemical processes involved in C cycling (Steinberg et al., 2006). Large inputs of terrestrial organic C and its subsequent mineralization have been suggested to be a major driver of CO_2 supersaturation commonly encountered in lakes (Duarte and Prairie, 2005; Cole et al., 2007; Prairie, 2008; Marotta et al., 2009).

The mechanistic connection between DOC and heterotrophic CO_2 production is believed to underpin the significant positive relationship between $p\text{CO}_2$ and DOC reported in comparative analyses (Houle et al., 1995; Sobek et al., 2005; Larsen et al., 2012). However, recent analyses have revealed that the relationship between $p\text{CO}_2$ and DOC in lake waters is regionally variable and not universal (Lapierre and del Giorgio, 2012). Hence, the relationship between $p\text{CO}_2$ and DOC reported in comparative analyses, based on data

sets dominated by temperate and high-latitude lakes ($> 33^\circ$), may not be extrapolated for all types of lakes, mainly because the tropical low-latitude lakes ($< 33^\circ$) are generally underrepresented in global data sets (Raymond et al., 2013).

One priority of comparative studies is the latitudinal variance, where lake temperature, ice cover and mixing regime will differ and these climatically driven processes, in turn, should strongly influence organic carbon cycling (Hanson et al., 2015). At low latitudes, warm conditions over the whole year may increase the metabolic rates involved in the C cycling in terrestrial (Ometto et al., 2005) and aquatic (Marotta et al., 2009, 2010) ecosystems on an annual basis compared to the high-latitude lakes. High temperatures affect heterotrophic activity and the associated mineralization rates of organic matter in soils (Davidson and Janssens, 2006), waters (López-Urrutia and Morán, 2007; Wohlers et al., 2008; Regaudie-de-Gioux and Duarte, 2012) and aquatic sediments (Wadham et al., 2012; Gudas et al., 2010; Marotta et al., 2014). Enhanced heterotrophic activity in warm ecosystems would support high aquatic CO_2 production and subsidize high CO_2 evasion from global lake water to the atmosphere.

The largest previous comparative analysis already published in the literature for global lake waters (Sobek et al., 2005) reported a significant positive relationship between DOC and $p\text{CO}_2$ and a non-significant variation of $p\text{CO}_2$ among lakes with changing temperature. However, both analyses were characterized by a paucity of low-latitude data. A strong positive relationship between temperature and $p\text{CO}_2$ was observed when subtropical and tropical ecosystems were included in the data set (Marotta et al., 2009), likely caused by the potential increase in metabolic rates under warmer conditions (Brown et al., 2004; López-Urrutia et al., 2006). Hence, the relationship between lake $p\text{CO}_2$ and DOC could also be temperature-dependent and, therefore, may differ between temperate and tropical lakes. The extensive low-latitude territory of Brazil, which has a high density of lakes and ponds (Downing et al., 2006), is appropriate to examine general patterns in the tropics (e.g., Marotta et al., 2009; Kosten et al., 2010). Here, we test the applicability of the relationship between $p\text{CO}_2$ and DOC using inputs derived from a high-latitude data set (Sobek et al., 2005) with added tropical and subtropical data of low-latitude lakes from Brazil.

2 Methods

2.1 Study area and lakes

Brazil extends from $5^\circ 16' 20''$ N to $33^\circ 44' 42''$ S, covering an area of approximately $8\,547\,000\text{ km}^2$, constituting half of South America, and it encompasses a high diversity of low-latitude landscapes (Ab'Saber, 2003) that are predominantly located within tropical latitudes. We conducted a survey of pH, alkalinity and DOC between 2003 and 2011

in surface waters of 166 permanent lakes from 0 to 33° of south latitude across Brazil (Fig. 1), yielding a total of 225 water samples. The lakes were sampled in representative biomes of Brazil: (1) the Amazonia forest (Amazonia biome, $n = 65$), (2) the Pantanal floodplain (Pantanal biome, $n = 29$) and (3) the tropical ($< 24^\circ$ of latitude) and (4) subtropical ($> 24^\circ$ and $< 33^\circ$ of south latitude) coasts, both in the Atlantic Forest biome ($n = 35$ and $n = 37$ lakes, respectively; Fig. 1). These biomes follow the classification of the Brazilian Institute of Geography and Statistics for biomes (IBGE 2004, ftp://geofpt.ibge.gov.br/mapas_tematicos/mapas_murais/biomas.pdf). Our data set encompasses a broad inter-lake heterogeneity ($n = 166$) for pH, alkalinity and DOC, simultaneously sampled among Brazilian biomes and along the latitudinal gradient, independent of the year's season.

The Amazonian Forest biome is formed by the most extensive hydrographic network on the globe – the Amazon River basin – which occupies a total area of approximately 6.11 million km^2 from its headwaters in the Peruvian Andes to its mouth in the Atlantic Ocean (ANA – www.ana.gov.br). The Amazon Forest is the Brazilian biome with the highest mean annual precipitation (approximately 2200 mm) and has warm mean air temperatures, approximately 25°C , high cloud coverage and high humidity with low fluctuations over the whole year (Chambers, 1999). We sampled a wide variety of lakes, characteristic of different areas of the Amazonian Forest, encompassing “clear” (low DOC and suspended solids), “white” (low DOC and high suspended solids) and “dark” (high DOC and low suspended solids) lakes.

The Pantanal floodplain is the world's largest tropical freshwater wetland, extending across an area of approximately $150\,000\text{ km}^2$ between 16 and 20° S and 58 and 55° W (Por, 1995). The annual average temperature and precipitation are approximately 22°C and 1000 mm , respectively (Mariot et al., 2007), with a strong seasonality and subsequent variation in the flooded area (Junk and Nunes da Cunha, 2005). The high-water period occurs during the rainy summer (usually from September to December), and low waters typically occur during the dry winter (from March to July; Hamilton et al., 2002).

The Atlantic Forest biome extends along a broad latitudinal belt, between 5 and 30° S from the subtropics to tropics and a narrow longitudinal section between 55 and 56° W , and occupies an area of 1.11 million km^2 along the Brazilian coast (IBGE; www.ibge.gov.br). This biome is characterized by numerous shallow coastal lakes, receiving high inputs of refractory organic matter (Farjalla et al., 2009) derived from the typical open xerophytic vegetation on sandy soils, where water retention is low (Scarano, 2002). The mean air temperatures vary from 27°C in winter to 30°C in summer at the tropical coast ($< 24^\circ$ of latitude; Chellappa et al., 2009) and from 17 and 20°C at the subtropical coast ($> 24^\circ$ of latitude; Waechter, 1998). The mean annual precipitation reaches 1164 mm (Henriques et al., 1986)




Figure 1. Geographic location of Brazilian lakes sampled in different biomes (IBGE 2004, available at ftp://geoftp.ibge.gov.br/mapas_tematicos/mapas_murais/biomas.pdf): Amazonia forest (vertical lines), Pantanal floodplain (dark gray) and Atlantic Forest (gray; tropical and subtropical coastal lakes).

and 1700 mm (Waechter, 1998) in the tropical and subtropical Brazilian coast, respectively. This biome is also characterized by strong seasonality, with rainy summers and dry winters (Chellappa et al., 2009).

2.2 Sampling design and analytical methods

Our sampling design encompassed the most representative Brazilian biomes from tropical and subtropical coastal areas to tropical and subtropical forests (Amazon and Atlantic Forest) and inland wetlands (Pantanal), with the intra-lake heterogeneity and seasonal fluctuations randomly assessed and further integrated by means of each ecosystem. To analyze the relationship between $p\text{CO}_2$ and DOC in tropical lake waters, we joined data on 194 lakes ($< 33^\circ$ of latitude) with both variables sampled at the same time, including 166 data samples from our own survey and 28 from the literature compilation (Table S1 in the Supplement). The values reported here, gathered in an opportunistic manner, represent daily averages ($N = 4$ or 5 samples) for a given year's season or/and one sampling time over different seasons, which were also both integrated by means of each lake. To test the global importance of the relationship between $p\text{CO}_2$ and DOC, we added our low-latitude data (225) to the Sobek et al. (2005) data set (4902 lakes) as this data set had a paucity of tropical

ecosystem data (148 tropical lakes, but only one with $p\text{CO}_2$ and DOC sampled at the same time).

pH, salinity and temperature were measured in situ. pH was determined using a pH meter (Digimed – DM2) with reference standards certified by Mettler Toledo (4.00 ± 0.01 and 7.00 ± 0.01 units) before each sampling hour. Temperature and salinity were measured using a thermosalinometer (Mettler Toledo – SevenGo™ SG3) coupled to a probe in Lab 737 previously calibrated with 0.01 M KCl. Surface lake water was collected for total alkalinity and DOC analyses, taking care to avoid bubbles at approximately 0.5 m of depth using a 1 L Van Dorn bottle. Total alkalinity (TA) was determined in the field by the Gran's titration method with 0.0125 M HCl immediately after sampling (Stumm and Morgan, 1996). Water samples for DOC were pre-filtered ($0.7 \mu\text{m}$, Whatman GF/F) and preserved by acidification with 85 % H_3PO_4 to reach a $\text{pH} < 2.0$ in sealed glass vials (Spyres et al., 2000). In the lab, DOC was determined by high-temperature catalytic oxidation using a TOC-5000 Shimadzu Analyzer; quality control was checked with a calibration curve made with potassium hydrogen phthalate before each sample battery analysis. $p\text{CO}_2$ concentrations in surface waters were calculated from pH and alkalinity following Weiss (1974), after corrections for temperature, altitude and ionic strength according to Cole et al. (1994).




Figure 2. Values of (a) temperature ($^{\circ}\text{C}$), (b) DOC concentrations (mg CL^{-1}) and (c) $p\text{CO}_2$ concentration (μatm) of Brazilian lakes sampled from different biomes, as defined by subtropical coastal lakes ($n = 37$), tropical coastal lake ($n = 63$), Pantanal floodplain ($n = 58$) and Amazonia forest ($n = 67$). The line depicts the median. The boxes show the quartiles, and the whiskers mark the 10th and 90th percentiles. Different lowercase letters near the box plot indicate significant statistic differences between the groups (Kruskall–Wallis followed by Dunn’s multiple comparison post hoc test, $p < 0.05$).

In order to address the potential contribution of DOC to TA, which is especially important in DOC-enriched acid freshwaters, we used the data set from Abril et al. (2015) to correct $p\text{CO}_2$ values calculated from pH and TA after the corrections for temperature, altitude and ionic strength (Cole et al., 1994). Full details on fitted regression equations to correct $p\text{CO}_2$ in function of the DOC and pH are described in the Supplement (Fig. S3).

2.3 Statistical analyses

The variables $p\text{CO}_2$ and DOC did not meet the assumptions of parametric tests even after logarithmic transformations (Zar, 1996) as the data were not normally distributed




Figure 3. Comparisons of $p\text{CO}_2$ against DOC concentrations for lakes from this study (black circles) and from Sobek et al. (2005; gray circles). Each point in the plot represents one measurement. The dashed line represents the linear regression for all Brazilian data points (not significant; $p > 0.05$), and the solid line represents the linear regression from Sobek et al. (2005; $p < 0.05$, $R^2 = 0.26$, $\log p\text{CO}_2$ (μatm) = $2.67 + 0.414 \log \text{DOC}$; mg CL^{-1}).

(Kolmogorov–Smirnov, $p < 0.05$) and the variances were heterogeneous (Bartlett, $p > 0.05$). Therefore, we used medians and nonparametric tests to compare these variables among biomes (Kruskall–Wallis followed by Dunn’s multiple comparison post hoc test, $p < 0.05$). The linear regression equations were fitted to compare our results with those of previous studies from Sobek et al. (2005). Statistical analyses were performed using the software Graphpad Prism version 4.0 for Macintosh (GraphPad Software, San Diego, CA).

3 Results

The lake waters surveyed were warm across all biomes (median 25–75 %, interquartile range = 27.5°C , 25.2 – 30.1) but colder in subtropical coastal lakes (23.4°C , 20.0 – 26.2) than in Pantanal and Amazonian lakes (29.5°C , 27.7 – 31.4 and 29.4°C , 27.6 – 31.0 , respectively; Dunn’s test, $p < 0.05$, Fig. 2a). DOC concentrations were consistently high (6.3 mg CL^{-1} , 4.3 – 11.9) for all Brazilian biomes but significantly lower in the Amazonian Forest (3.8 mg CL^{-1} , 2.7 – 5.8) than in the tropical coast (13.4 mg CL^{-1} , 6.1 – 32.8 ; Fig. 2b; Dunn’s test, $p < 0.05$). Most lakes (approximately 83 % of raw data) showed surface waters supersaturated in CO_2 relative to the atmospheric equilibrium ($p\text{CO}_2$ in atmospheric equilibrium is $400.83 \mu\text{atm}$, 2015 annual mean; data available in www.esrl.noaa.gov/gmd/ccgg/trends/), with much higher $p\text{CO}_2$ values in Amazonian lakes ($7956 \mu\text{atm}$, 3033 – $11\,346$) than in subtropical coastal lakes ($900 \mu\text{atm}$, 391.3 – 3212 ; Fig. 2c; Dunn’s test, $p < 0.05$).

The $p\text{CO}_2$ in the surface waters of Brazilian lakes was independent of DOC concentrations (linear regression for raw data, $p > 0.05$, Fig. 3). The same absence of positive significance pattern was found in comparison with corrected data. A negative (linear regression, $p < 0.05$, $R^2 = 0.03$, $n = 194$, $p\text{CO}_2 = -98.76 (\pm 39.92) \times \text{DOC} + 6529 (\pm 641.1)$) or non-significant (linear regression, $p > 0.05$) DOC– $p\text{CO}_2$ relationship for tropical lakes ($N = 194$, DOC- and pH-corrected data, respectively (Fig. S3a and c)) contrasted with a significant positive relationship for those at other latitudes ($N = 4433$; linear regression, $p < 0.05$, $R^2 = 0.20$, $p\text{CO}_2 = 64.43 (\pm 2.04) \times \text{DOC} + 625.1 (\pm 20.87)$ and $R^2 = 0.12$, $p\text{CO}_2 = 45.70 (\pm 1.84) \times \text{DOC} + 623.7 (\pm 18.83)$) for DOC-corrected data and pH-corrected data, respectively (Fig. S3b and d, full details on corrections in the Supplement). The range of $p\text{CO}_2$ for a similar DOC range in Brazilian lakes was larger than that reported by Sobek et al. (2005) for the data set dominated by high-latitude cold lakes, despite the number of lakes in their data set being much larger (more details in the Supplement, Fig. S3).

4 Discussion

The Brazilian lakes sampled here were characterized by a prevalence of CO_2 supersaturation, consistent with general trends previously reported for global lakes (e.g., Raymond et al., 2013; Cole et al., 1994, 2007), including those at tropical latitudes (Marotta et al., 2009). The very high $p\text{CO}_2$ levels observed here, with a median of 900 and 8300 μatm for subtropical and Amazon lake waters, respectively, are consistent with those reported previously for the Amazon River and tributaries (2000–12 000 μatm ; Richey et al., 2002), Amazon floodplain lakes (3000–4898 μatm ; Rudorff et al., 2012), Pantanal lakes and wetlands (2732–10 620 μatm ; Hamilton et al., 1995), and coastal lakes (768–9866 μatm ; Kosten et al., 2010; 361–20 037 μatm ; Marotta et al., 2010) and for global values for tropical lakes (1255–35 278 μatm ; Marotta et al., 2009), reservoirs (1840 μatm ; Aufdenkampe et al., 2011) and wetlands (3080–6170 μatm ; Aufdenkampe et al., 2011).

The non-significant or weakly negative relationship (Fig. S3) between DOC and $p\text{CO}_2$ reported here for warm low-latitude lakes contrasted with significant positive relationships derived from previous data sets dominated by high-latitude lakes (Houle, 1995; Prairie, 2002; Jonsson et al., 2003; Sobek et al., 2005; Roehm et al., 2009; Lapiere and del Giorgio, 2012; Larsen et al., 2012). The results presented show that warm low-latitude lakes range widely in $p\text{CO}_2$, reaching very high and low values, but tend to have comparatively more uniform DOC concentrations (Fig. 3). More intense metabolic processes that uptake and release CO_2 in lake waters, autotrophy and heterotrophy, respectively, could determine an enhanced variability in lake $p\text{CO}_2$ with decreasing latitude (Marotta et al., 2009).

In this way, the inclusion of warm tropical data in our study revealed novel increases in the variability of the DOC– $p\text{CO}_2$ relationship in lakes over the latitudinal gradient. One explanation for this pattern is that even similar DOC concentrations, representing the total pool of DOC, may show different mixtures between origins from aquatic primary producers and terrestrial sources (Kritzberg et al., 2006). The autochthonous DOC (i.e., produced in the lake) is related to the net CO_2 uptake (Staehr and Sand Jansen, 2007), while the allochthonous DOC (i.e., produced in the catchment) is a resource to the net CO_2 release in lake waters (Sobek et al., 2007). The increased DOC release from aquatic primary producers into waters under tropical conditions, especially warmer annual conditions and higher solar incidence, can offset any positive relationship between $p\text{CO}_2$ and the terrestrial DOC that causes the net aquatic heterotrophy to subsidize (Marotta et al., 2010, 2012). This contributes to the explanation of non-significant relationships reported here (Fig. 3), suggesting a temperature dependence of the DOC– $p\text{CO}_2$ relationship in global lakes.

In conclusion, the finding that $p\text{CO}_2$ does not increase with DOC concentration in Brazilian tropical lakes rejects the hypothesis that DOC serves as a universal predictor for $p\text{CO}_2$ in lake waters (Larsen et al., 2012). Even discounting a possible artifact of the method that could be causing an overestimation in the values of $p\text{CO}_2$ or considering the contribution of organic acids to the alkalinity, the pattern of no relationship between DOC and $p\text{CO}_2$ in the tropical lakes was strongly confirmed (Fig. S3). Therefore, our results, contributing to the filling of a gap in the literature of tropical studies, suggest potentially important latitudinal differences for depositional aquatic environments, whose causes still need to be better addressed to improve accuracy of global C cycle models.

The Supplement related to this article is available online at doi:10.5194/bg-13-865-2016-supplement.

Author contributions. All authors contributed to the study design, data interpretation and preparation or refinement of the manuscript. L. Pinho and H. Marotta performed the sampling and sample analyses.

Acknowledgements. This research is a contribution to projects from Brazilian research agencies (FAPERJ, CAPES and CNPq). L. Pinho was supported by PhD scholarships from CAPES (period in Brazil) and FAPERJ (period in Spain). A. Enrich-Prast received postdoctoral and other CAPES and CNPq fellowships during studies at Linköping University. H. Marotta was supported by a research fellowship from FAPERJ (Programa Jovem Cientista do Nosso Estado), and a research grant from CNPq.

Edited by: M. Dai

References

- Abril, G., Bouillon, S., Darchambeau, F., Teodoru, C. R., Marwick, T. R., Tamooh, F., Ochieng Omengo, F., Geeraert, N., Deirmendjian, L., Polsenaere, P., and Borges, A. V.: Technical Note: Large overestimation of $p\text{CO}_2$ calculated from pH and alkalinity in acidic, organic-rich freshwaters, *Biogeosciences*, 12, 67–78, doi:10.5194/bg-12-67-2015, 2015.
- Ab'Saber, A.: Os Domínios de Natureza no Brasil: Potencialidades paisagísticas, Ateliê editorial Brazil, 2003.
- Aufdenkampe, A., Mayorga, E., Raymond, P., Melack, J., and Doney, S.: Riverine coupling of biogeochemical cycles between land, oceans, and atmosphere, *Front. Ecol. Environ.*, 9, 53–60, 2011.
- Brow, J., Gillooly, J., Allen, A., Savage, V., and West, G.: Toward a metabolic theory of ecology, *Ecology*, 85, 1771–1789, 2004.
- Chambers, J., Higuchi, N., Schimel, J., Ferreira, L., and Melack, J.: Decomposition and carbon cycling of dead trees in tropical forests of the central Amazon, *Oecologia*, 122, 380–388, 1999.
- Chellappa, N., Câmara, F., and Rocha, O.: Phytoplankton community: indicator of water quality in the Armando Ribeiro Gonçalves Reservoir and Pataxó Channel, Rio Grande do Norte, Brazil, *Braz. J. Biol.*, 69, 241–251, 2009.
- Cole, J. J., Caraco, N. F., Kling, G. W., and Kratz, T. K.: Carbon-Dioxide Supersaturation in the Surface Waters of Lakes, *Science*, 265, 1568–1570, 1994.
- Cole, J. J., Prairie, Y. T., Caraco, N. F., McDowell, W. H., Tranvik, L. J., Striegl, R. G., Duarte, C. M., Kortelainen, P., Downing, J. A., Middelburg, J. J., and Melack, J.: Plumbing the global carbon cycle: Integrating inland waters into the terrestrial carbon budget, *Ecosystems*, 10, 171–184, 2007.
- Davidson, E. and Janssens, I.: Temperature sensitivity of soil carbon decomposition and feedbacks to climate change, *Nature*, 440, 165–173, doi:10.1038/nature04514, 2006.
- Downing, J. A., Prairie, Y. T., Cole, J. J., Duarte, C. M., Tranvik, L. J., Striegl, R. G., McDowell, W. H., Kortelainen, P., Caraco, N. F., Melack, J. M., and Middelburg, J.: The global abundance and size distribution of lakes, ponds, and impoundments, *Limnol. Oceanogr.*, 51, 2388–2397, 2006.
- Downing, J. A., Cole, J. J., Middelburg, J. J., Striegl, R. G., and Duarte, C. M.: Sediment organic carbon burial in agriculturally eutrophic impoundments over the last century, *Glob. Biogeochem. Cy.*, 22, GB1018, doi:10.1029/2006GB002854, 2008.
- Duarte, C. M. and Prairie, Y. T.: Prevalence of heterotrophy and atmospheric CO_2 emissions from aquatic ecosystems, *Ecosystems*, 8, 862–870, 2005.
- Farjalla, V. F., Amado, A. M., Suhett, A. L., and Meirelles-Pereira, F.: DOC removal paradigms in highly humic aquatic ecosystems, *Environ. Sci. Pollut. R.* 16, 531–538, 2009.
- Gudasz, C., Bastviken, D., Steger, K., Premke, K., and Sobek, S.: Temperature-controlled organic carbon mineralization in lake sediments, *Nature*, 466, 478–481, 2010.
- Hamilton, S. K., Sippel, S. J., and Melack, J. M.: Oxygen depletion and carbon dioxide and methane production in waters of the Pantanal wetland of Brazil, *Biogeochemistry*, 30, 115–141, 1995.
- Hamilton, S. K., Sippel, S. J., and Melack, J. M.: Comparison of inundation patterns among major South American floodplains, *J. Geophys. Res.*, 107, D208038, doi:10.1029/2000JD000306, 2002.
- Hanson, P., Pace, M., Carpenter, C., Cole, J., and Stanley, E.: Integrating Landscape Carbon Cycling: Research Needs for Resolving Organic Carbon Budgets of Lakes, *Ecosystems*, 18, 363–375, doi:10.1007/s10021-014-9826-9, 2015.
- Henriques, R. P. B., Araújo, D. S. D., and Hay, J. D.: Descrição e classificação dos tipos de vegetação da restinga de Carapebús, Rio de Janeiro, *Rev. Bras. Bot.* 9, 173–189, 1986.
- Houle, D., Carigan, R., Lachance, M., and DuPont, J.: Dissolved organic carbon and sulfur in southwestern Québec lakes: Relationships with catchment and lake properties, *Limnol. Oceanogr.*, 40, 710–710, 1995.
- Jonsson, A., Karlsson, J., and Jansson, M.: Sources of carbon dioxide supersaturation in clearwater and humic lakes in Northern Sweden, *Ecosystems*, 6, 224–235, 2003.
- Junk, W. J. and Nunes de Cunha, C.: Pantanal: a large South American wetland at a crossroads, *Ecol. Eng.*, 24, 391–401, 2005.
- Kosten, S., Roland, F., Da Motta Marques, D., Van Nes, E., Mazzeo, N., Sternberg, L., Scheffer, M., and Cole, J.: Climate-dependent CO_2 emissions from lakes, *Global Biogeochem. Cy.*, 24, GB2007, doi:10.1029/2009GB003618, 2010.
- Kritzberg, E., Cole, J., Pace, M., and Granéli, W.: Bacterial growth on allochthonous carbon in humic and nutrient-enriched lakes: Results from whole-lake ^{13}C addition experiments, *Ecosystems*, 9, 489–499, 2006.
- Lapierre, J. F. and del Giorgio, P. A.: Geographical and environmental drivers of regional differences in the lake $p\text{CO}_2$ versus DOC relationship across northern landscapes, *J. Geophys. Res.-Biogeo.*, 117, G03015, doi:10.1029/2012JG001945, 2012.
- Larsen, S., Andersen, T., and Hessen, D.: The $p\text{CO}_2$ in boreal lakes: Organic carbon as a universal predictor?, *Glob. Biogeochem. Cy.*, 25, GB2012, doi:10.1029/2010GB003864, 2012.
- López-Urrutia, Á. and Morán, X. A. G.: Resource limitation of bacterial production distorts the temperature dependence of oceanic carbon cycling, *Ecology*, 88, 817–822, 2007.
- López-Urrutia, Á., San Martín, E., Harris, R., and Irigoien, X.: Scaling the metabolic balance of the oceans, *P. Natl. Acad. Sci. USA*, 103, 8739–8744, 2006.
- Mariot, M., Dudal, Y., Furian, S., Sakamoto, A., Valles, V., Valls, V., Fort, M., and Barbiero, L.: Dissolved organic matter fluorescence as a water-flow tracer in the tropical wetland of Pantanal of Nhecolândia, Brazil, *Sci. Total Environ.*, 388, 184–193, 2007.
- Marotta, H., Duarte, C. M., Sobek, S., and Enrich-Prast, A.: Large CO_2 disequilibria in tropical lakes, *Glob. Biogeochem. Cy.*, 23, GB4022, doi:10.1029/2008GB003434, 2009.
- Marotta, H., Duarte, C. M., Pinho, L., and Enrich-Prast, A.: Rainfall leads to increased $p\text{CO}_2$ in Brazilian coastal lakes, *Biogeosciences*, 7, 1607–1614, doi:10.5194/bg-7-1607-2010, 2010.
- Marotta, H., Duarte, C. M., Guimarães-Souza, B., and Enrich-Prast, A.: 2012 Synergistic control of CO_2 emissions by fish and nutrients), *Oecologia*, 168, 839–847, 2012.
- Marotta, H., Pinho, L., Gudasz, C., Bastviken, D., Tranvik, L., and Enrich-Prast, A.: Greenhouse gas production in low-latitude lake sediments responds strongly to warming, *Nature Climate Change*, 4, 467–470, 2014.

- McDonald, C., Rover, J., Stets, E., and Striegl, R.: The regional abundance and size distribution of lakes and reservoirs in the United States and implications for estimates of global lake extent, *Limnol. Oceanogr.*, 57, 597–606, 2012.
- Ometto, J. P. H. B., Nobre, A., Rocha, H., Artaxo, P., and Martinelli, L.: Amazonia and the modern carbon cycle: lessons learned, *Oecologia*, 143, 483–500, 2005.
- Por, F.: The Pantanal of Mato Grosso (Brazil) – World’s Largest Wetlands, Klume Academic Publishers, Brazil, 1995.
- Prairie, Y.: The summer metabolic balance in the epilimnion of southeastern Quebec lakes, *Limnol. Oceanogr.*, 47, 316–321, 2002.
- Prairie, Y. T.: Carbocentric limnology: looking back, looking forward, *Can. J. Fish. Aquat. Sci.*, 65, 543–548, 2008.
- Raymond, P., Hartmann, J., Lauerwald, R., Sobek, S., McDonald, C., Hoover, M., Butman, D., Striegl, R., Mayorga, E., Humborg, C., Kortelainen, P., Durr, H., Meybeck, M., Ciais, P., and Guth, P.: Global carbon dioxide emissions from inland waters, *Nature*, 503, 355–359, 2013.
- Regaudie-de-Gioux, A. and Duarte C. M.: Temperature dependence of planktonic metabolism in the ocean, *Global Biogeochem. Cy.*, 26, GB1015, doi:10.5194/bg-11-4529-2014, 2012.
- Richey, J. E., Melack, J. M., Aufdenkampe, A. K., Ballester, V. M., and Hess, L. L.: Outgassing from Amazonian rivers and wetlands as a large tropical source of atmospheric CO_2 , *Nature*, 416, 617–620, 2002.
- Roehm, C., Prairie, Y. T., and del Giorgio, P.: The $p\text{CO}_2$ dynamics in lakes in the boreal region of northern Québec, Canada, *Global Biogeochem. Cy.*, 23, GB3013, doi:10.1029/2008GB003297, 2009.
- Rudorff, C. M., Melack, J. M., MacIntyre, S., Barbosa, C. C. F., and Novo, E. M. L. M.: Seasonal and spatial variability of CO_2 emission from a large floodplain lake in the lower Amazon, *J. Geophys. Res.*, 117, G01002, doi:10.1029/2011JG001699, 2012.
- Scarano, F.: Structure, function and floristic relationships of plant communities in stressful habitats marginal to the Brazilian Atlantic rainforest, *Ann. Bot.*, 90, 517–524, 2002.
- Sobek, S., Tranvik, L. J., and Cole, J. J.: Temperature independence of carbon dioxide supersaturation in global lakes, *Global Biogeochem. Cy.*, 19, GB2003, doi:10.1029/2004GB002264, 2005.
- Sobek, S., Tranvik, L. J., and Cole, J. J.: Patterns and regulation of dissolved organic carbon: An analysis of 7500 widely distributed lakes, 2007.
- Spyres, G., Nimmo, M., Worsfold, P., Achterberg, E., and Miller, A.: Determination of dissolved organic carbon in seawater using high temperature catalytic oxidation techniques, *Trends Anal. Chem.*, 19, 498–506, 2000.
- Staeher, P. and Sand-Jansen, K.: Temporal dynamics and regulation of lake metabolism, *Limnol. Oceanogr.* 52, 108–110, 2007.
- Steinberg, C. E. W., Kamara, S., Prokhotskaya, V., Manusadzianas, L., Karasyova, T., Timofeyev, M., Jie, Z. Paul, A., Meinelt, T., Farjalla, V., Matsuo, A., Burnison, B., and Menzel, R.: Humic substances in the environment with an emphasis on freshwater systems, *Environ. Sci. Pollut. Res. Int.*, 15, 15–16, 2006.
- Stumm, W. and Morgan, J. J.: Aquatic chemistry: chemical equilibria and rates in natural waters, vol 3., Wiley, New York, 1996.
- Tranvik, L. J., Downing, J. A., Cotner, J. B., Loiselle, S. A., Striegl, R. G., Ballatore, T. J., Dillon, P., Finlay, K., Fortino, K., Knoll, L. B., Kortelainen, P. L., Kutser, T., Larsen, S., Laurion, I., Leech, D. M., McCallister, S. L., McKnight, D. M., Melack, J. M., Overholt, E., Porter, J. A., Prairie, Y., Renwick, W. H., Roland, F., Sherman, B. S., Schindler, D. W., Sobek, S., Tremblay, A., Vanni, M. J., Verschoor, A. M., von Wachenfeldt, E., and Weyhenmeyer, G. A.: Lakes and reservoirs as regulators of carbon cycling and climate, *Limnol. Oceanogr.*, 54, 2298–2314, 2009.
- Wadham, J., Arndt, S., Tulaczyk, S., Stibal, M., M. Tranter, M., Telling, J., Lis, G., Lawson, E., Ridgwell, A., Dubnick, A., Sharp, M., Anesio, A., and Butler, C.: Potential methane reservoirs beneath Antarctica, Potential methane reservoirs beneath Antarctica, *Nature*, 488, 633–637, 2012.
- Waechter, J. L.: Epiphytic orchids in eastern subtropical South America. Proc. 15th World Orchid Conference, *Naturalia Publications*, 332–341, 1998.
- Weiss, R. F.: Carbon Dioxide in water and seawater: The solubility of non-ideal gas, *Mar. Chem.*, 2, 203–215, 1974.
- Wohlers, J., Engel, A., Zollner, E., Breithaupt, P., Jurgens, K., Hoppe, H., Sommer, U., and Riebesell, U.: Changes in biogenic carbon flow in response to sea surface warming, *P. Natl. Acad. Sci.*, 106, 7067–7072, 2008.
- Zar, J. H.: Biostatistical analysis, 3 ed., Prentice Hall., New Jersey, 1996.